

CircARAP2* controls MICA-induced NK cell desensitization by erasing CTCF/PRC2-induced suppression in early endosome marker *RAB5A

Cellular and Molecular Life Sciences

Feifei Guo¹, Nawen Du¹, Xue Wen¹, Zhaozhi Li¹, Yantong Guo¹, Lei Zhou¹, Andrew R. Hoffman², Lingyu Li^{1#}, Ji-Fan Hu^{1,2#}, Jiuwei Cui^{1#}

¹ Cancer Center, The First Hospital of Jilin University, Changchun, 130021, China

² Stanford University School of Medicine, VA Palo Alto Health Care System, Palo Alto, CA 94304, USA

Correspondence to: Jiuwei Cui, M.D., Ph.D., Cancer Center, The First Hospital of Jilin University, 71 Xinmin Street, Changchun 130021, China, Tel: 86-43188782178, Fax: 86-43188786134, e-mail: cuijw@jlu.edu.cn; Ji-Fan Hu, M.D., Ph.D., Department of Medicine, PAVIR, VA Palo Alto Health Care System, Palo Alto, CA 94304, USA, Tel: 650-852-3275, Fax: 650-856-8024, e-mail: jifan@stanford.edu; jifanhu@jlu.edu.cn; Lingyu Li, M.D., Ph.D., Cancer Center, The First Hospital of Jilin University, 71 Xinmin Street, Changchun 130021, China, e-mail: lilingyu@jlu.edu.cn.

Supplementary Table S1 A list of siRNA sequences used in this manuscript

Target gene	Primers	Sequence	Name
CTCF	Sense	5'-GCCUGCCGUAGAAAUUGAATT-3'	CTCF-homo-2325
	Antisense	5'-UUCAAUUUCUACGGCAGGCTT-3'	
	Sense	5'-GGCAAGACAUGCUGAUAAUTT-3'	CTCF-homo-2154
	Antisense	5'-AUUAUCAGCAUGUCUUGCCTT-3'	
RAB5A	Sense	5'-GUCCUAUGCAGAUAGACAAUTT-3'	RAB5A-homo-980
	Antisense	5'-AUUGUCAUCUGCAUAGGACTT-3'	
	Sense	5'-CCAGUUCAAACUAGUACUUTT-3'	RAB5A-homo-592
	Antisense	5'-AAGUACUAGUUUGAACUGGTT-3'	
SRSF1	Sense	5'-GCCCAGAAGUCCAAGUUAUTT-3'	SRSF1-homo-797
	Antisense	5'-AUAACUUGGACUUCUGGGCTT-3'	
	Sense	5'-GGAACAACGAUUGCCGCAUTT-3'	SRSF1-homo-244
	Antisense	5'-AUGCGGCAAUCGUUGUUCCTT-3'	
NC	Sense	5'-UUCUCCGAACGUGUCACGUTT-3'	NC
	Antisense	5'-ACGUGACACGUUCGGAGAATT-3'	