

Electronic supplementary material (ESM)

ESM Table 1. List of medical treatments used by both groups

Medical Treatment	CAN N (% of group)	NO CAN N (% of group)	P value
Diabetes treatment			
- Insulin injection therapy	16 (67)	19 (79)	0.330
- Insulin pumps	8 (33)	5 (21)	0.330
- Concomitant treatment (GLP-1 RA, SGLT2i, metformin)	1 (4)	0 (0)	1.000
Antihypertensive treatment			
- Renal-angiotensin blockage	13 (54)	10 (42)	0.386
- Calcium channel blockers	8 (33)	5 (21)	0.330
- Beta blockers	2 (8)	0 (0)	0.489 [#]
- Diuretics	4 (17)	3 (13)	1.000 [#]
Lipid lowering treatment	17 (71)	16 (70)	0.386
Antithrombotic treatment	9 (38)	7 (29)	0.540
Antidepressive medication (tricyclic antidepressant (n=3) or serotonin reuptake inhibitors (n=4))	7 (29)	0 (0)	0.009 ^{*#}

Data are present as x(y) whereby x refers to the absolute number of participants using a particular treatment and y refers to that number when expressed as a percentage of the group. N=24 in both groups. * $p \leq 0.05$ between groups in the number of people using the treatment. # The fishers exact test p value (2-sided) was used as more than 20% of cells had expected frequencies < 5. GLP-1 RA: glucagon-like peptide-1 receptor agonists. SGLT2i: sodium-glucose cotransporter-2 inhibitors. CAN: Cardiovascular autonomic neuropathy.