

Article title

Major depressive disorders increase the susceptibility to self-reported infections in two German cohort studies

Journal name

Social Psychiatry and Psychiatric Epidemiology

Author names and affiliations

Henning Elpers¹, Henning Teismann, PhD¹, Jürgen Wellmann, PhD¹, Klaus Berger, MD¹, André Karch, MD¹, Nicole Rübsamen, PhD^{1,*}


¹ Institute of Epidemiology and Social Medicine, University of Münster, Germany.

* Corresponding author:

Nicole Rübsamen | Institute of Epidemiology and Social Medicine | University of Münster |
Albert-Schweitzer-Campus 1 | 48149 Münster | Germany
clinepi@uni-muenster.de


Online Resource 4: p value functions

P value function of the effect of MDD on URTI in the BiDirect (n=925) and HaBIDS (n=1,007) cohort study




The circle represents the counterfactual incidence rate ratio.

P value function of the effect of MDD on LRTI in the BiDirect (n=925) and HaBIDS (n=1,007) cohort study




The circle represents the counterfactual incidence rate ratio.

P value function of the effect of MDD on any RTI in the BiDirect (n=925) and HaBIDS (n=1,007) cohort study




The circle represents the counterfactual incidence rate ratio.

P value function of the effect of MDD on Cystitis in the BiDirect (n=925) and HaBIDS (n=1,007) cohort study



The circle represents the counter null incidence rate ratio.

P value function of the effect of MDD on fever in the BiDirect (n=925) and HaBIDS (n=1,007) cohort study



The circle represents the counter null incidence rate ratio.