

Supplementary Information

Tributyltin induces conjugation of ATG8s to single membranes via the V-ATPase–ATG16L1 axis, leading to transcription factor EB activation in human cell lines

Shunichi Hatamiya¹, Masatsugu Miyara^{1,†}, Nanako Takahashi¹, Ami Oguro¹, Yaichiro Kotake^{1,†}

¹Graduate School of Biomedical and Health Sciences, Hiroshima University, Hiroshima 734-8553, Japan

[†]Corresponding authors:

Masatsugu Miyara; e-mail address: miyara128@hiroshima-u.ac.jp

Yaichiro Kotake; e-mail address: yaichiro@hiroshima-u.ac.jp

Fig. S1




Fig. S1 Western blot images corresponding to Fig. 1a and 1c. **A–C** Three independent experiments corresponding to Fig. 1a. **D–F** Three independent experiments corresponding to Fig. 1c.

Fig. S2




Fig. S2 Western blot images corresponding to Fig. 2a and 2c. **A–C** Three independent experiments corresponding to Fig. 2a. **D–F** Three independent experiments corresponding to Fig. 2c.

Fig. S3




Fig. S3 Western blot images corresponding to Fig. 2e. **A–D** Four independent experiments corresponding to Fig. 2e.

Fig. S4




Fig. S4 Western blot images corresponding to Fig. 3a. Three independent experiments corresponding to Fig. 3a.

Fig. S5




Fig. S5 Detailed confocal images corresponding to Fig. 5c.

Fig. S6




Fig. S6 Western blot images corresponding to Fig. 5e. **A–C** Three independent experiments corresponding to Fig. 5e.