

Influence of larval prey availability on year class strength in Faroe Plateau cod (*Gadus morhua*)

Sólvá Jacobsen¹, Kristoffer Klitgaard Nielsen², Rune Kristiansen², Peter Grønkjær², Eilif Gaard¹ and Petur Steingrund¹.

¹Faroe Marine Research Institute, Nóatún 1, PO Box 3051, Tórshavn FO-100, Faroe Islands.

²Department of Bioscience, Aquatic Biology, Aarhus University, Ole Worms Allé 1, DK-8000 Aarhus C, Denmark.

*Correspondence:

Sólvá Jacobsen

solvaj@hav.fo

Supplementary material

Table S1 One-way ANOVA testing for interannual differences in mean zooplankton biomass ($\log \text{mg m}^{-3} + 1$) on the inner Faroe shelf 1998-2005.

			Df	Sum sq	Mean Sq	<i>F</i>	<i>P</i>
April	Copepod eggs	Year	6	0.05	0.01	2.33	0.054
		Station	35	0.13	0		
	Calanoid nauplii	Year	6	0.26	0.04	5.14	< 0.01
		Station	35	0.29	0.01		
	<i>Acartia</i> sp.	Year	6	2.26	0.38	17.25	< 0.01
		Station	35	0.76	0.02		
	<i>Pseudocalanus</i> sp.	Year	6	0.61	0.1	3.82	< 0.01
		Station	35	0.93	0.03		
	<i>C. fin.</i> < 1.5 mm	Year	6	1.29	0.21	11.28	< 0.01
		Station	35	0.67	0.02		
	<i>T. longicornis</i>	Year	6	0.32	0.05	1.55	0.193
		Station	35	1.2	0.03		
	<i>C. fin.</i> $\geq$ 1.5 mm	Year	6	0.7	0.12	1.55	0.19
		Station	35	2.65	0.08		
	Decapod larvae	Year	6	1.19	0.2	2.35	0.052
		Station	35	2.95	0.08		
June	<i>Acartia</i> sp.	Year	5	0.84	0.17	10.6	< 0.01
		Station	58	0.92	0.02		
	<i>Pseudocalanus</i> sp.	Year	5	0.24	0.05	4.62	< 0.01
		Station	58	0.61	0.01		
	<i>C. fin.</i> < 1.5 mm	Year	5	1.25	0.25	12.23	< 0.01
		Station	58	1.19	0.02		
	<i>T. longicornis</i>	Year	5	3.15	0.63	10.22	< 0.01
		Station	58	3.57	0.06		
	<i>C. fin.</i> $\geq$ 1.5 mm	Year	5	12.01	2.4	13.6	< 0.01
		Station	58	10.25	0.18		
	Decapod larvae	Year	5	3.2	0.64	6	< 0.01
		Station	58	6.18	0.11		