

Microbiome of zoophytophagous biological control agent *Nesidiocoris tenuis*

Yuta Owashi,¹ Toma Minami,² Taisei Kikuchi,^{3,4} Akemi Yoshida,³ Ryohei Nakano,^{2,5} Daisuke Kageyama,^{1#} Tetsuya Adachi-Hagimori^{2#}

¹ Institute of Agrobiological Sciences, National Agriculture and Food Research Organization (NARO), Tsukuba, Ibaraki, Japan

² Laboratory of Applied Entomology, University of Miyazaki, Miyazaki, Japan

³ Frontier Science Research Center, University of Miyazaki, Miyazaki, Japan

⁴ Department of Integrated Biosciences, Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan

⁵ Shizuoka Prefectural Research Institute of Agriculture and Forestry, Shizuoka, Japan

#Address correspondence to Tetsuya Adachi-Hagimori, tadachi@cc.miyazaki-u.ac.jp

#Address correspondence to Daisuke Kageyama, kagyamad@affrc.go.jp

Yuta Owashi and Toma Minami contributed equally to this work.




Fig. S1 Results of amplicon sequencing of the hypervariable V3/V4 region in 16S rRNA of the laboratory strain of *N. tenuis* derived from population No. 1 in Table S1. Assigned bacterial taxa are color-coded as blue for *wNtenA* and red for *Rickettsia*, consistent with Fig. 1. The next generation of an original female represents G1 and that of G1 represents G2. Under laboratory conditions, eggs of *Ephestia kuehniella* and leaves of *Crassula ovata* were offered as food and oviposition substrate, respectively