

Supplementary figures

Applied Microbiology and Biotechnology

Prevalence and Genomic Insights into *Yersinia enterocolitica* in Southeastern China (2008–2022)

Lei Fang¹, Shuxuan Li⁵, Jie Rong³, Shengkai Li⁴, Yuwen Zhang⁶, Huihuang Lou², Zhongbi Xie², Yuqin Hu², Yuejin Wu¹, Airong Xie^{2*}, Yi Li^{2*}

¹Department of Critical Care Medicine, Sir Run Run Shaw Hospital, College of Medicine, Zhejiang University, Hangzhou, China

²Wenzhou Center for Disease Control and Prevention, Wenzhou, China


³Lotabiome Biotechnology Inc., Suzhou, China

⁴Key Laboratory of Alkene-carbon Fibres-based Technology & Application for Detection of Major Infectious Diseases, MOE Key Laboratory of Geriatric Diseases and Immunology, Cancer Institute, Suzhou Medical College, Soochow University, Suzhou, China

⁵Nanfang College Guangzhou, Guangzhou, China

⁶School of Public Health, Hangzhou Medical College, Hangzhou, China

*Corresponding author: Dr. Yi Li, Wenzhou Center for Disease Control and Prevention, Wenzhou, China; zjwzliyi@126.com; 86-13858889839; Dr. Airong Xie, Wenzhou Center for Disease Control and Prevention, Wenzhou, China; sharonxar@aliyun.com; 86-13587428098.



Supplemental Fig. S1 The host transmission frequency of *Y. enterocolitica*. The arrows represent the direction of host transmission, with arrow darkness indicating transmission frequency.

