

Online Resource 4 Seventeen positive Moran's eigenvector mapping (MEM) variables identified in the analysis of presence data of 403 macrobenthic taxa (species or genera) identified in grab samples taken at a total of 341 stations in the Barents, Kara, and Laptev Seas during summers (May-October) between 1991 and 2014. MEM variables are visually subdivided into three categories representing distinct spatial scales

Unravelling the effects of environmental drivers and spatial structure on benthic species distribution patterns in Eurasian-Arctic seas (Barents, Kara and Laptev Seas)

Polar Biology

Miriam L. S. Hansen^{1, *}, Dieter Piepenburg^{1,2,3}, Dmitrii Pantiukhin⁴ and Casper Kraan^{2,3,5}

¹Institute of Ecosystem Research, Christian Albrecht University of Kiel, 24118 Kiel, Germany

²Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI), 27570 Bremerhaven, Germany

³Helmholtz Institute for Functional Marine Biodiversity at the University of Oldenburg, 23129 Oldenburg, Germany

⁴Institute of Earth Sciences, Saint Petersburg State University, St. Petersburg 199034, Russia

⁵Institute of Sea Fisheries, Johann Heinrich von Thünen Institute, 27572 Bremerhaven, Germany

*Corresponding author: mhansen@ecology.uni-kiel.de

