

Taxa	F. O. (%)	R. A. (%)	Max Ab. (org. 1000 m³)	Mean Ab (org. 1000 m³)	N
Hidrozoa					
N. I.	4	0.05	53	2.11	2
Rhopalonematidae	4	0.05	59	2.35	2
Siphonophora					
N. I.	4	0.02	36	1.43	32
<i>Diphyes antarctica</i> Moser, 1925	16	0.17	103	8.52	7
<i>Dimophyes arctica</i> (Chun, 1897)	52	0.89	265	43.57	36
Polychaeta					
<i>Vanadis antarctica</i> (McIntosh, 1885)	28	0.20	69	9.50	8
<i>Vanadis longissima</i> (Levinsen, 1885)	4	0.02	24	0.98	1
Larvae	24	0.45	156	22.74	18
Pteropoda					
<i>Spongiobranchaea australis</i> d'Orbigny, 1836	4	0.02	34	1.38	1
Ostracoda N.I.	60	7.08	2647	333.43	286
Copepoda					
<i>Calanoides acutus</i> (Giesbrecht, 1902)	60	2.43	375	113.10	98
<i>Calanus propinquus</i> Brady, 1883	8	0.05	31	2.39	2
Copepodite Calanidae	8	0.07	63	3.68	3
Copepodite <i>Calanoides</i> spp.	32	0.67	313	33.04	27
<i>Rhincalanus gigas</i> Brady, 1883	8	0.05	29	2.25	2
<i>Rhincalanus nasutus</i> Giesbrecht, 1888	4	0.02	29	1.18	1
<i>Ctenocalanus citer</i> Heron & Bowman, 1971	4	0.02	25	1.00	1
<i>Microcalanus pygmaeus</i> (Sars G.O., 1900)	8	0.10	103	5.31	4
Copepodite <i>Clausocalanus</i> sp.	4	0.07	100	4.00	3
<i>Aetideus arcuatus</i> (Vervoort, 1949)	4	0.02	29	1.14	1
<i>Aetideopsis antarctica</i> (Wolfenden, 1908)	8	0.10	88	4.64	4
<i>Chiridius gracilis</i> Farran, 1908	8	0.10	83	4.48	4
<i>Gaetanus tenuispinus</i> (Sars G.O., 1900)	8	0.05	32	2.47	2
<i>Paraeuchaeta biloba</i> Farran, 1929	4	0.05	63	2.50	2
<i>Paraeuchaeta exigua</i> (Wolfenden, 1911)	24	0.17	53	7.92	7
Copepodite <i>Paraeuchaeta</i> spp.	52	3.61	1367	185.60	146
<i>Scaphocalanus curtus</i> (Farran, 1926)	4	0.02	29	1.18	1
<i>Scaphocalanus elongatus</i> Scott A., 1909	4	0.05	59	2.35	2
<i>Scolecithricella minor minor</i> (Brady, 1883)	20	0.15	54	6.65	6
<i>Scolecithricella vittata</i> (Giesbrecht, 1893)	4	0.07	86	3.43	3
<i>Scottocalanus securifrons</i> (Scott T., 1894)	4	0.02	34	1.38	1
Copepodite <i>Scaphocalanus</i> sp.	4	0.02	29	1.18	1
Copepodite Scolecithricidae	12	0.10	69	5.34	4
<i>Haloptilus ocellatus</i> Wolfenden, 1905	4	0.02	29	1.18	1
<i>Haloptilus oxycephalus</i> (Giesbrecht, 1889)	4	0.02	29	1.18	1
<i>Heterorhabdus lobatus</i> Bradford, 1971	4	0.02	28	1.11	1
Heteroabdididae	4	0.02	29	1.18	1
<i>Metridia gerlachei</i> Giesbrecht, 1902	92	13.37	3500	615.07	540
<i>Metridia longa</i> (Lubbock, 1854)	28	0.25	63	11.80	10
Copepodite <i>Metridia</i> spp.	80	30.38	6063	1440.84	1227
Copepodite <i>Centropages</i> sp.	4	0.02	26	1.05	1
<i>Labidocera</i> sp.	4	0.02	32	1.29	1

Copepodite N.I.	28	0.64	342	29.53	26
<i>Oithona frigida</i> Giesbrecht, 1902	12	1.31	1125	64.42	53
<i>Oithona similis</i> Claus, 1866	8	1.16	1313	59.40	47
<i>Oithona</i> spp.	56	3.61	1667	177.58	246
<i>Triconia antarctica</i> (Heron, 1977)	28	0.45	353	21.08	18
<i>Agetus flaccus</i> (Giesbrecht, 1891)	4	0.07	88	3.53	3
Amphipoda					
<i>Eusirus microps</i> Walker, 1906	4	0.02	26	1.05	1
<i>Cylopus lucasi</i> Spence Bate, 1862	8	0.05	32	2.34	2
<i>Vibilia antarctica</i> Stebbing, 1888	16	0.12	65	5.81	5
<i>Vibilia armata</i> Bovallius, 1887	24	0.17	53	7.50	7
Manca Stage	8	0.05	32	2.34	2
Euphausiacea					
<i>Thysanoessa macrura</i> G.O. Sars, 1883	24	0.62	372	25.61	25
<i>Euphausia superba</i> Dana, 1850	4	0.02	23	0.93	1
Calyptopis	4	0.05	63	2.50	2
Furcilia	48	1.06	259	37.76	43
Chaetognata					
N. I.	52	2.01	541	91.76	81
<i>Eukrohnia hamata</i> (Möbius, 1875)	60	2.85	676	133.11	115
<i>Pseudosagitta maxima</i> (Conant, 1896)	32	0.40	88	18.12	16
<i>Pseudosagitta</i> spp.	4	0.15	158	6.32	6
Appendicularia					
<i>Oikopleura</i> sp.	8	0.07	59	3.73	3
<i>Fritillaria</i> sp.	4	0.05	69	2.76	2
Thaliacea					
<i>Salpa maxima</i> Forskål, 1775	8	0.05	31	2.43	2
<i>Salpa thompsoni</i> Foxton, 1961	84	21.49	5698	912.12	960
<i>Ihlea racovitzai</i> (Van Beneden & De Selys Longchamp, 1913)	8	0.07	54	3.41	3