

Perceptions of radiologists on structured reporting for cancer imaging – a survey by the European Society of Oncologic Imaging (ESOI)

Electronic Supplementary Material

CT CHEST, ABDOMEN, PELVIS

DATE:

CLINICAL INDICATION:

TECHNIQUE:

COMPARISON TO:

FINDINGS:

LUNGS AND PLEURA:

MEDIASTINUM / THORACIC NODES:

HEPATOBIILIARY:

PANCREAS:

SPLEEN:

KIDNEYS AND ADRENAL GLANDS:

ABDOMINOPELVIC NODES:

GASTROINTESTINAL TRACT / PERITONEUM:

PELVIC ORGANS:

BONES AND SOFT TISSUES:

CONCLUSION:

TUMOR STAGE (BASELINE):

TREATMENT RESPONSE:

☐ CR ☐ PR ☐ SD ☐ PD ☐ N/A ☐ Other:

Diagnostic certainty:

Unlikely:	<10%
Less likely:	~25%
Possibly:	~50%
Probably / suspicious for:	~75%
Consistent with:	>90%

Quantitative descriptors:

Few:	2-5
Multiple :	6-20
Numerous:	>20

CT CHEST, ABDOMEN, PELVIS

DATE:

CLINICAL INDICATION:

TECHNIQUE:

COMPARISON TO:

FINDINGS:

CHEST:

ABDOMEN:

PELVIS:

BONES / SOFT TISSUES:

CONCLUSION:

TUMOR STAGE (BASELINE):

TREATMENT RESPONSE:

☐ CR ☐ PR ☐ SD ☐ PD ☐ N/A ☐ Other:



Diagnostic certainty:

Unlikely:	<10%
Less likely:	~25%
Possibly:	~50%
Probably / suspicious for:	~75%
Consistent with:	>90%

Quantitative descriptors:

Few:	2-5
Multiple :	6-20
Numerous:	>20