

## Further Reading

Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine

J. Patijn

### Cervical Region

- Aartun, E., Aartun, E., Degerfalk, A., Degerfalk, A., Kentsdotter, L., Kentsdotter, L., & Hestbaek, L. (2014). Screening of the spine in adolescents: inter- and intra-rater reliability and measurement error of commonly used clinical tests. *BMC Musculoskeletal Disorders*, 15(1), 1–10. <http://doi.org/10.1186/1471-2474-15-37>
- Arumugam, A., Mani, R., & Raja, K. (2011). Interrater reliability of the craniocervical flexion test in asymptomatic individuals--a cross-sectional study. *Journal of Manipulative and Physiological Therapeutics*, 34(4), 247–253. <http://doi.org/10.1016/j.jmpt.2011.04.011>
- Bae, J.-M. (2014). An overview of systematic reviews of diagnostic tests accuracy. *Epidemiology and Health*, 36, e2014016. <http://doi.org/10.4178/epih/e2014016>
- Bakhtadze, M. A., Patijn, J., Galaguz, V. N., Bolotov, D. A., & Popov, A. A. (2011). Inter-examiner reproducibility of the segmental motion palpation springing test for side bending at level C2–C3. *International Musculoskeletal Medicine*, 33(1), 8–14.
- Bernhoff, G., Landén Ludvigsson, M., Peterson, G., Bertilson, B. C., Elf, M., & Peolsson, A. (2016). The pain drawing as an instrument for identifying cervical spine nerve involvement in chronic whiplash-associated disorders. *Journal of Pain Research*, 9, 397–404. <http://doi.org/10.2147/JPR.S104747>
- Bertilson, B. C., Grunnesjö, M., & Streder, L.-E. (2003). Reliability of Clinical Tests in the Assessment of Patients With Neck/Shoulder Problem—Impact of History. *Spine*, 28(19), 2222–2231. <http://doi.org/10.1097/01.BRS.0000089685.55629.2E>
- Beyer, B., Sobczak, S., Salem, W., Feipel, V., & Dugailly, P.-M. (2016). 3D motion reliability of occipital condylar glide testing: From concept to kinematics evidence. *Manual Therapy*, 21, 159–164. <http://doi.org/10.1016/j.math.2015.07.005>
- Billis, E. V., Foster, N. E., & Wright, C. C. (2003). Reproducibility and repeatability: errors of three groups of physiotherapists in locating spinal levels by palpation. *Manual Therapy*.
- Branney, J., & Breen, A. C. (2014). Does inter-vertebral range of motion increase after spinal manipulation? A prospective cohort study. *Chiropractic & Manual Therapies*, 22, 24. <http://doi.org/10.1186/s12998-014-0024-9>

Further Reading, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
Cervical Region

- Cattrysse, E., Swinkels, R. A., Oostendorp, R. A., & Duquet, W. (1997). Upper cervical instability: are clinical tests reliable? *Manual Therapy*, 2(2)91–97.  
<http://doi.org/10.1054/math.1997.0290>
- Chaves, T. C., Nagamine, H. M., de Sousa, L. M., de Oliveira, A. S., & Grossi, D. B. (2010). Comparison between the reliability levels of manual palpation and pressure pain threshold in children who reported orofacial pain. *Manual Therapy*, 15(5), 508–512.  
<http://doi.org/10.1016/j.math.2010.03.010>
- Cleland, J. A., Childs, J. D., Fritz, J. M., & Whitman, J. M. (2006). Interrater reliability of the history and physical examination in patients with mechanical neck pain. *Archives of Physical Medicine and Rehabilitation*, 87(10), 1388–1395. <http://doi.org/10.1016/j.apmr.2006.06.011>
- Cohen, S. P., Bajwa, Z. H., Kraemer, J. J., Dragovich, A., Williams, K. A., Stream, J., et al. (2007). Factors predicting success and failure for cervicofacet radiofrequency denervation: a multi-center analysis. *Regional Anesthesia and Pain Medicine*, 32(6), 495–503.  
<http://doi.org/10.1016/j.rapm.2007.05.009>
- Cook, C., & Hegedus, E. (2011). Diagnostic utility of clinical tests for spinal dysfunction. *Manual Therapy*, 16(1), 21–25. <http://doi.org/10.1016/j.math.2010.07.004>
- de Wit, M. P. T., Berlo, S. E., Aanerud, G. J., Aletaha, D., Bijlsma, J. W., Croucher, L., et al. (2005). Three- dimensional measurement of intervertebral range of motion in ossification of the posterior longitudinal ligament: are there mobile segments in the continuous type? *Physical Therapy*, 2013(1-2), 475–485.e10. <http://doi.org/10.1016/j.soard.2006.05.005>
- Deboer, K. F., Harmon, R., Tuttle, C. D., & Wallace, H. (1985). Reliability study of detection of somatic dysfunctions in the cervical spine. *Journal of Manipulative and Physiological Therapeutics*, 8(1), 9–16.
- Fedorak, C., Ashworth, N., Marshall, J., & Paull, H. (2003). Reliability of the visual assessment of cervical and lumbar lordosis: how good are we? *Spine*, 28(16), 1857–1859. <http://doi.org/10.1097/01.BRS.0000083281.48923.BD>
- Fjellner, A., Bexander, C., Faleij, R., & Strender, L. E. (1999a). Interexaminer reliability in physical examination of the cervical spine. *Journal of Manipulative and Physiological Therapeutics*, 22(8), 511– 516.
- Fjellner, A., Bexander, C., Faleij, R., & Strender, L. E. (1999b). Interexaminer reliability in physical examination of the cervical spine. *Journal of Manipulative and Physiological Therapeutics*, 22(8), 511– 516.
- Gadotti, I. C., & Magee, D. (2013). Validity of surface markers placement on the cervical spine for craniocervical posture assessment. *Manual Therapy*, 18(3), 243–247.  
<http://doi.org/10.1016/j.math.2012.10.012>
- Gemmell, H., & Miller, P. (2005). Interexaminer reliability of multidimensional examination regimens used for detecting spinal manipulable lesions: a systematic review. *Clin*

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
Cervical Region

Chiropractic, 8, 199–204.

- Gross, A. R., Haines, T., Thomson, M. A., Goldsmith, C., & McIntosh, J. (1996). Diagnostic tests for temporomandibular disorders: an assessment of the methodologic quality of research reviews. *Manual Therapy*, 1(5), 250–257. <http://doi.org/10.1054/math.1996.0275>
- Hall, T. M., Robinson, K. W., Fujinawa, O., Akasaka, K., & Pyne, E. A. (2008). Intertester Reliability and Diagnostic Validity of the Cervical Flexion- Rotation Test. *Journal of Manipulative and Physiological Therapeutics*, 31(4), 293–300. <http://doi.org/10.1016/j.jmpt.2008.03.012>
- Hall, T., Briffa, K., Hopper, D., & Robinson, K. (2010). Long-term stability and minimal detectable change of the cervical flexion-rotation test. *The Journal of Orthopaedic and Sports Physical Therapy*, 40(4), 225–229. <http://doi.org/10.2519/jospt.2010.3100>
- Haneline, M. T., & Young, M. (2009). A review of intraexaminer and interexaminer reliability of static spinal palpation: a literature synthesis. *Journal of Manipulative and Physiological Therapeutics*, 32(5), 379–386. <http://doi.org/10.1016/j.jmpt.2009.04.010>
- Haneline, M. T., Cooperstein, R., Young, M., & Birkeland, K. (2008). Spinal motion palpation: a comparison of studies that assessed intersegmental end feel vs excursion. *Journal of Manipulative and Physiological Therapeutics*, 31(8), 616–626. <http://doi.org/10.1016/j.jmpt.2008.09.007>
- Haneline, M., Cooperstein, R., Young, M., & Birkeland, K. (2009). An annotated bibliography of spinal motion palpation reliability studies. *J Can Chiropr Assoc*, 53(1), 40–58.
- Hanney, W. J., George, S. Z., Kolber, M. J., Young, I., Salamh, P. A., & Cleland, J. A. (2014). Inter-rater reliability of select physical examination procedures in patients with neck pain. *Physiotherapy Theory and Practice*, 30(5), 345–352. <http://doi.org/10.3109/09593985.2013.870267>
- Hanten, W., Olson, S., & Ludwig, G. (2002). Reliability of manual mobility testing of the upper cervical spine in subjects with cervicogenic headache. *Journal of Manual*, 10(2), 76–82.
- Harrison, D. E., Haas, J. W., Harrison, D. D., Holland, B., & Janik, T. (2005). Sagittal Skin Contour of the Cervical Spine: Interexaminer and
- Intraexaminer Reliability of the Flexicurve Instrument. *Journal of Manipulative and Physiological Therapeutics*, 28(7), 516–519. <http://doi.org/10.1016/j.jmpt.2005.07.009>

- Hollerwöger, D., & Hollerwöger, D. (2006). Methodological quality and outcomes of studies addressing manual cervical spine examinations: a review. *Manual Therapy*.
- Hoppenbrouwers, M., Eckhardt, M. M. E. M., Verkerk, K., & Verhagen, A. (2006). Reproducibility of the measurement of active and passive cervical range of motion. *Journal of Manipulative and Physiological Therapeutics*, 29(5), 363–367. <http://doi.org/10.1016/j.jmpt.2006.04.007>
- Hoving, J. L., Pool, J. J. M., van Mameren, H., Devillé, W. J. L. M., Assendelft, W. J. J., de Vet, H. C. W., et al. (2005). Reproducibility of cervical range of motion in patients with neck pain. *BMC Musculoskeletal Disorders*, 6, 59. <http://doi.org/10.1186/1471-2474-6-59>
- Hsu, W.-H., Chen, Y.-L., Lui, T.-N., Chen, T.-Y., Hsu, Y.-H., Lin, C.-L., & Ming-Lun, T. (2011). Comparison of the kinematic features between the in vivo active and passive flexion-extension of the subaxial cervical spine and their biomechanical implications. *Spine*, 36(8), 630–638. <http://doi.org/10.1097/BRS.0b013e3181da79af>
- Hubka, M. J., & Phelan, S. P. (1994). Interexaminer reliability of palpation for cervical spine tenderness. *Journal of Manipulative and Physiological Therapeutics*, 17(9), 591–595.
- Humphreys, B. K., Delahaye, M., & Peterson, C. K. (2004). An investigation into the validity of cervical spine motion palpation using subjects with congenital block vertebrae as a 'gold standard'. *BMC Musculoskeletal Disorders*, 5(1), 19. <http://doi.org/10.1186/1471-2474-5-19>
- James, G., & Doe, T. (2010). The craniocervical flexion test: intra-tester reliability in asymptomatic subjects. *Physiotherapy Research International : the Journal for Researchers and Clinicians in Physical Therapy*, 15(3), 144–149. <http://doi.org/10.1002/pri.456>
- Jorritsma, W., Dijkstra, P. U., De Vries, G. E., Geertzen, J. H. B., & Reneman, M. F. (2014). Physical dysfunction and nonorganic signs in patients with chronic neck pain: exploratory study into interobserver reliability and construct validity. *The Journal of Orthopaedic and Sports Physical Therapy*, 44(5), 366–376. <http://doi.org/10.2519/jospt.2014.4715>
- Jull, G., Bogduk, N., & Marsland, A. (1988). The accuracy of manual diagnosis for cervical zygapophysial joint pain syndromes. *The Medical Journal of Australia*, 148(5), 233–236.
- Jull, G., Zito, G., Trott, P., Potter, H., & Shirley, D. (1997). Inter-examiner reliability to detect painful upper cervical joint dysfunction. *The Australian Journal of Physiotherapy*, 43(2), 125–129.
- Jørgensen, R., Ris, I., Falla, D., & Juul-Kristensen, B. (2014). Reliability, construct and discriminative validity of clinical testing in subjects with and without chronic neck pain. *BMC Musculoskeletal Disorders*, 15, 408. <http://doi.org/10.1186/1471-2474-15-408>

- Kaale, B. R., Kråkenes, J., Albrektsen, G., & Wester, K. (2008). Clinical assessment techniques for detecting ligament and membrane injuries in the upper cervical spine region—A comparison with MRI results. *Manual Therapy*, 13(5), 397–403. <http://doi.org/10.1016/j.math.2007.03.007>
- King, W., Lau, P., Lees, R., & Bogduk, N. (2007). The validity of manual examination in assessing patients with neck pain. *The Spine Journal : Official Journal of the North American Spine Society*, 7(1), 22–26. <http://doi.org/10.1016/j.spinee.2006.07.009>
- Licht, G., Müller-Ehrenberg, H., Mathis, J., Berg, G., & Greitemann, G. (2007). Clinical examination of myofascial trigger points is reliable. *Manuelle Medizin*, 45(6), 402–408. <http://doi.org/10.1007/s00337-007-0559-0>
- Lijmer, J. G., Mol, B. W., Heisterkamp, S., Bonsel, G. J., Prins, M. H., van der Meulen, J. H., & Bossuyt, P.
- M. (1999). Empirical evidence of designrelated bias in studies of diagnostic tests. *JAMA: the Journal of the American Medical Association*, 282(11), 1061–1066.
- Lindgren, K. A., Leino, E., & Manninen, H. (1992). Cervical rotation lateral flexion test in brachialgia. *Archives of Physical Medicine and Rehabilitation*, 73(8), 735–737.
- Ljungquist, T., Harms-Ringdahl, K., Nygren, A., & Jensen, I. (1999). Intra- and inter-rater reliability of an 11-test package for assessing dysfunction due to back or neck pain. *Physiotherapy Research International : the Journal for Researchers and Clinicians in Physical Therapy*, 4(3), 214–232.
- Maigne, J.-Y., Chantelot, F., & Chatellier, G. (2009). Interexaminer agreement of clinical examination of the neck in manual medicine. *Annals of Physical and Rehabilitation Medicine*, 52(1), 41–48. <http://doi.org/10.1016/j.rehab.2008.11.001>
- Malanga, G. A., Landes, P., & Nadler, S. F. (2003). Provocative tests in cervical spine examination: historical basis and scientific analyses. *Pain Physician*, 6(2), 199–205.
- Manning, D. M., Dedrick, G. S., Sizer, P. S., & Brismée, J.-M. (2012). Reliability of a seated three-dimensional passive intervertebral motion test for mobility, end-feel, and pain provocation in patients with cervicalgia. *The Journal of Manual & Manipulative Therapy*, 20(3), 135–141. <http://doi.org/10.1179/2042618611Y.0000000023>
- Marcotte, J., Normand, M. C., & Black, P. (2002). The kinematics of motion palpation and its effect on the reliability for cervical spine rotation. *Journal of Manipulative and Physiological Therapeutics*, 25(7), E7.
- Marcotte, J., Normand, M. C., & Black, P. (2005). Measurement of the pressure applied during motion palpation and reliability for cervical spine rotation. *Journal of Manipulative and Physiological Therapeutics*, 28(8), 591–596. <http://doi.org/10.1016/j.jmpt.2005.08.003>

- Matteucci, M. J., Moszyk, D., & Migliore, S. A. (2015). Agreement between resident and faculty emergency physicians in the application of NEXUS criteria for suspected cervical spine injuries., 48(4), 445–449. <http://doi.org/10.1016/j.jemermed.2014.11.006>
- May, S., Littlewood, C., & Bishop, A. (2006). Reliability of procedures used in the physical examination of non-specific low back pain: A systematic review. *Australian Journal of Physiotherapy*, 52(2), 91–102. [http://doi.org/10.1016/S0004-9514\(06\)70044-7](http://doi.org/10.1016/S0004-9514(06)70044-7)
- Mior, S. A., King, R. S., McGregor, M., & Bernard, M. (1985). Intra and interexaminer reliability of motion palpation in the cervical spine. *The Journal of the CCA*, 29(4), 195–198.
- Moran, R. (2001). Intraexaminer and interexaminer reliability for palpation of the cranial rhythmic impulse at the head and sacrum. *Journal of Manipulative and Physiological Therapeutics*, 24(3), 183–190. <http://doi.org/10.1067/mmt.2001.113773>
- Myburgh, C., Lauridsen, H. H., Larsen, A. H., & Hartvigsen, J. (2011). Standardized manual palpation of myofascial trigger points in relation to neck/shoulder pain; the influence of clinical experience on inter-examiner reproducibility. *Manual Therapy*, 16(2), 136–140. <http://doi.org/10.1016/j.math.2010.08.002>
- Nansel, D. D., Peneff, A. L., Jansen, R. D., & Cooperstein, R. (1989). Interexaminer concordance in detecting joint-play asymmetries in the cervical spines of otherwise asymptomatic subjects. *Journal of Manipulative and Physiological Therapeutics*, 12(6), 428–433.
- Olson, L. E., Millar, A. L., Dunker, J., Hicks, J., & Glanz, D. (2006). Reliability of a Clinical Test for Deep Cervical Flexor Endurance. *Journal of Manipulative and Physiological Therapeutics*, 29(2), 134–138. <http://doi.org/10.1016/j.jmpt.2005.12.009>
- Patijn, J., & Ellis, R. (2001a). Low back pain: reproducibility of diagnostic procedures in manual/musculoskeletal medicine. *Journal of Orthopaedic Medicine*, 23(1), 36–42. Retrieved from <http://www.google.nl/search?>
- Patijn, J., & Ellis, R. F. (2001b). Low Back Pain: Reproducibility of diagnostic procedures in Manual/Musculoskeletal Medicine. *J Orthop Med*, 23(1), 36–42.
- Patroncini, M., Hannig, S., Meichtry, A., & Luomajoki, H. (2014). Reliability of movement control tests on the cervical spine. *BMC Musculoskeletal Disorders*, 15, 402. <http://doi.org/10.1186/1471-2474-15-402>
- Perreault, N., Brisson, C., Dionne, C. E., Montreuil, S., & Punnett, L. (2008). Agreement between a self-administered questionnaire on musculoskeletal disorders of the neck-shoulder region and a physical examination. *BMC Musculoskeletal Disorders*, 9, 34. <http://doi.org/10.1186/1471-2474-9-34>

- Piva, S. R., Erhard, R. E., Childs, J. D., & Browder, D. A. (2006). Inter-tester reliability of passive intervertebral and active movements of the cervical spine. *Manual Therapy*, 11(4), 321–330. <http://doi.org/10.1016/j.math.2005.09.001>
- Pool, J. J., Hoving, J. L., de Vet, H. C., van Mameren, H., & Bouter, L. M. (2004). The interexaminer reproducibility of physical examination of the cervical spine. *Journal of Manipulative and Physiological Therapeutics*, 27(2), 84–90. <http://doi.org/10.1016/j.jmpt.2003.12.002>
- Póvoa, L. C., Ferreira, A. P. A., & Silva, J. G. (2015). Validation of palpatory methods for evaluating anatomical bone landmarks of the cervical spine: a systematic review. *Journal of Manipulative and Physiological Therapeutics*, 38(4), 302–310. <http://doi.org/10.1016/j.jmpt.2015.04.002>
- Rey-Eiriz, G., Alburquerque-Sendín, F., Barrera-Mellado, I., Martín-Vallejo, F. J., & Fernández-de-las-Peñas, C. (2010). Validity of the posterioranterior middle cervical spine gliding test for the examination of intervertebral joint hypomobility in mechanical neck pain. *Journal of Manipulative and Physiological Therapeutics*, 33(4), 279–285. <http://doi.org/10.1016/j.jmpt.2010.03.005>
- Robinson, R., Robinson, H. S., Bjørke, G., & Kvale, A. (2009). Reliability and validity of a palpation technique for identifying the spinous processes of C7 and L5. *Manual Therapy*, 14(4), 409–414. <http://doi.org/10.1016/j.math.2008.06.002>
- Rossetini, G., Rondoni, A., Lovato, T., Strobe, M., Verzè, E., Vicentini, M., & Testa, M. (2016). Intra- and inter-rater reliability of 3D passive intervertebral motion in subjects with nonspecific neck pain assessed by physical therapy students: A pilot study. *Journal of Back and Musculoskeletal Rehabilitation*. <http://doi.org/10.3233/BMR-160718>
- Rubinstein, S. M., & van Tulder, M. (2008). A best-evidence review of diagnostic procedures for neck and low-back pain. *Best Practice & Research. Clinical Rheumatology*, 22(3), 471–482. <http://doi.org/10.1016/j.berh.2007.12.003>
- Schneider, G. M., Jull, G., Thomas, K., & Salo, P. (2011). Screening of patients suitable for diagnostic cervical facet joint blocks - A role for physiotherapists. *Manual Therapy*, 17(2), 180–183. <http://doi.org/10.1016/j.math.2011.11.004>
- Schneider, G. M., Jull, G., Thomas, K., Smith, A., Emery, C., Faris, P., et al. (2013). Intrarater and interrater reliability of select clinical tests in patients referred for diagnostic facet joint blocks in the cervical spine. *Archives of Physical Medicine and Rehabilitation*, 94(8), 1628–1634. <http://doi.org/10.1016/j.apmr.2013.02.015>
- Schöps, P., Pflingsten, M., & Siebert, U. (2000). Reliability of manual medical examination techniques of the cervical spine. Study of quality assurance in manual diagnosis]. *Zeitschrift Für Orthopädie Und Ihre Grenzgebiete*, 138(1), 2–7. <http://doi.org/10.1055/s-2000-10105>

- Sebastian, D., Chovvath, R., & Malladi, R. (2015). Cervical extensor endurance test: a reliability study. *Journal of Bodywork and Movement Therapies*, 19(2), 213–216. <http://doi.org/10.1016/j.jbmt.2014.04.014>
- Seffinger, M. A., Najm, W. I., Mishra, S. I., Adams, A., Dickerson, V. M., Murphy, L. S., & Reinsch, S. (2004). Reliability of Spinal Palpation for Diagnosis of Back and Neck Pain. *Spine*, 29(19), E413–E425. <http://doi.org/10.1097/01.brs.0000141178.98157.8e>
- Segarra, V., Dueñas, L., Torres, R., Falla, D., Jull, G., & Lluch, E. (2015). Inter-and intra-tester reliability of a battery of cervical movement control dysfunction tests. *Manual Therapy*, 20(4), 570–579. <http://doi.org/10.1016/j.math.2015.01.007>
- Shin, S., Yoon, D. M., & Yoon, K. B. (2011). Identification of the correct cervical level by palpation of spinous processes. *Anesthesia and Analgesia*, 112(5), 1232–1235. <http://doi.org/10.1213/ANE.0b013e3182110f9f>
- Smedmark, V., Wallin, M., & Arvidsson, I. (2000). Inter-examiner reliability in assessing passive intervertebral motion of the cervical spine. *Manual Therapy*, 5(2), 97–101. <http://doi.org/10.1054/math.2000.0234>
- Southerst, D., Stupar, M., Côté, P., Mior, S., & Stern, P. (2013). The reliability of measuring pain distribution and location using body pain diagrams in patients with acute whiplash-associated disorders. *Journal of Manipulative and Physiological Therapeutics*, 36(7), 395–402. <http://doi.org/10.1016/j.jmpt.2013.05.023>
- Stamos, I., Heneghan, N. R., McCarthy, C., & Wright, C. (2012). Inter-examiner reliability of active combined movements assessment of subjects with a history of mechanical neck problems. *Manual Therapy*, 17(5), 438–444. <http://doi.org/10.1016/j.math.2012.04.009>
- Stochkendahl, M. J., Christensen, H. W., Hartvigsen, J., Vach, W., Haas, M., Hestbaek, L., et al. (2006). Manual examination of the spine: a systematic critical literature review of reproducibility. *Journal of Manipulative and Physiological Therapeutics*, 29(6), 475–85–485.e1–10. <http://doi.org/10.1016/j.jmpt.2006.06.011>
- Stovall, B. A., & Kumar, S. (2010). Reliability of bony anatomic landmark asymmetry assessment in the lumbopelvic region: application to osteopathic medical education. *The Journal of the American Osteopathic Association*, 110(11), 667–674. <http://doi.org/10.1016/j.apmr.2010.12.042>
- Strender, L. E., Lundin, M., & Nell, K. (1997). Interexaminer reliability in physical examination of the neck. *Journal of Manipulative and Physiological Therapeutics*, 20(8), 516–520.
- Ughetto, J., & Lempereur, J. J. (2013). Étude de la reproductibilité intraet inter-observateurs des mesures centimétriques de la mobilité du rachis cervical et du rachis thoracolombaire. *Kinésithérapie*,

- Van Suijlekom, H. A., de Vet, H. C., Van Den Berg, S. G., & Weber, W. E. (2000). Interobserver reliability in physical examination of the cervical spine in patients with headache. *Headache*, 40(7), 581–586.
- Van Suijlekom, J. A., de Vet, H. C., Van Den Berg, S. G., & Weber, W. E. (1999). Interobserver reliability of diagnostic criteria for cervicogenicheadache. *Cephalalgia : an International Journal of Headache*, 19(9), 817–823.
- Van Trijffel, E., Anderegg, Q., Bossuyt, P. M. M., & Lucas, C. (2005). Inter-examiner reliability of passive assessment of intervertebral motion in the cervical and lumbar spine: A systematic review. *Manual Therapy*, 10(4), 256–269. <http://doi.org/10.1016/j.math.2005.04.008>
- van Trijffel, E., Lindeboom, R., Bossuyt, P. M., Schmitt, M. A., Lucas, C., Koes, B. W., & Oostendorp, R. (2014). Indicating spinal joint mobilisations or manipulations in patients with neck or low-back pain: protocol of an inter-examiner reliability study among manualtherapists. *Chiropractic & Manual Therapies*, 22, 22. <http://doi.org/10.1186/2045-709X-22-22>
- Viikari-Juntura, E. (1987). Interexaminer reliability of observations in physical examinations of the neck. *Physical Therapy*, 67(10), 1526–1532.
- Viitanen, J. V., Kokko, M. L., Heikkilä, S., & Kautiainen, H. (1998). Neck mobility assessment in ankylosing spondylitis: a clinical study of nine measurements including new tape methods for cervical rotation and lateral flexion. *British Journal of Rheumatology*, 37(4), 377–381.
- Wainner, R. S., Fritz, J. M., Irrgang, J. J., Boninger, M. L., Delitto, A., & Allison, S. (2003). Reliability and diagnostic accuracy of the clinical examination and patient self-report measures for cervical radiculopathy. *Spine*, 28(1), 52–62.
- <http://doi.org/10.1097/01.BRS.0000038873.01855.50>
- Williams, M. A., McCarthy, C. J., Chorti, A., Cooke, M. W., & Gates, S. (2010). A systematic review of reliability and validity studies of methods or measuring active and passive cervical range of motion. *Journal of Manipulative and Physiological Therapeutics*, 33(2), 138–155. <http://doi.org/10.1016/j.jmpt.2009.12.009>

## Lumbar Region

- Aalto, T. J., Airaksinen, O., Härkönen, T. M., & Arokoski, J. P. (2005). Effect of passive stretch on reproducibility of hip range of motion measurements. *Archives of Physical Medicine and Rehabilitation*, 86(3), 549–557. <http://doi.org/10.1016/j.apmr.2004.04.041>
- Aartun, E., Aartun, E., Degerfalk, A., Degerfalk, A., Kentsdotter, L., Kentsdotter, L., & Hestbaek, L. (2014). Screening of the spine in adolescents: inter- and intra-rater reliability and measurement error of commonly used clinical tests. *BMC Musculoskeletal Disorders*, 15(1), 1–10. <http://doi.org/10.1186/1471-2474-15-37>
- Abbott, J. H., McCane, B., Herbison, P., Moginie, G., Chapple, C., & Hogarty, T. (2005). Lumbar segmental instability: a criterion-related validity study of manual therapy assessment. *BMC Musculoskeletal Disorders*, 6, 56. <http://doi.org/10.1186/1471-2474-6-56>
- Adams, M. A., & Dolan, P. (1991). A technique for quantifying the bending moment acting on the lumbar spine in vivo. *Journal of Biomechanics*, 24(2), 117–126.
- Adhia, D. B., Tumilty, S., Mani, R., Milosavljevic, S., & Bussey, M. D. (2016). Can hip abduction and external rotation discriminate sacroiliac joint pain? *Manual Therapy*, 21, 191–197. <http://doi.org/10.1016/j.math.2015.08.002>
- Alviso, D. J., Dong, G. T., & Lentell, G. L. (1988). Intertester reliability for measuring pelvic tilt in standing. *Physical Therapy*, 68(9), 1347–1351.
- Alyazedi, F. M., Lohman, E. B., Wesley Swen, R., & Bahjri, K. (2015). The inter-rater reliability of clinical tests that best predict the subclassification of lumbar segmental instability: structural, functional and combined instability. *The Journal of Manual & Manipulative Therapy*, 23(4), 197–204. <http://doi.org/10.1179/2042618615Y.0000000002>
- Arab, A. M., Abdollahi, I., Joghataei, M. T., Golafshani, Z., & Kazemnejad, A. (2009). Inter- and intra- examiner reliability of single and composites of selected motion palpation and pain provocation tests for sacroiliac joint. *Manual Therapy*, 14(2), 213–221. <http://doi.org/10.1016/j.math.2008.02.004>
- Arab, A. M., Carmichael, J. P., Abdollahi, I., Joghataei, M. T., Golafshani, Z., & Kazemnejad, A. (1987). Inter- and intra-examiner reliability of palpation for sacroiliac joint dysfunction. *Journal of Manipulative and Physiological Therapeutics*, 10(4), 164–171.
- Arana, E., Royuela, A., Kovacs, F. M., Estremera, A., Sarasibar, H., Amengual, G., et al. (2010). Lumbar Spine: Agreement in the Interpretation of 1.5-T MR Images by Using the Nordic Modic Consensus Group Classification Form1. *Dx.Doi.org*, 254(3), 809–817. <http://doi.org/10.1148/radiol.09090706>

- Ayala, F., Sainz de Baranda, P., De Ste Croix, M., & Santonja, F. (2012). Reproducibility and concurrent validity of hip joint angle test for estimating hamstring flexibility in recreationally active young men. *Journal of Strength and Conditioning Research / National Strength & Conditioning Association*, 26(9), 2372–2382. <http://doi.org/10.1519/JSC.0b013e31823db1e2>
- Azevedo, D. C., Lauria, A. C., manipulative, A. P. J. O., 2013. (n.d.). Intraexaminer and interexaminer reliability of pressure biofeedback unit for assessing lumbopelvic stability during 6 lower limb movement tests. *International Journal of Osteopatic Medicine*. <http://doi.org/10.1016/j.jmpt.2012.12.008>
- Azevedo, D. C., Santos, H., Carneiro, R. L., & Andrade, G. T. (2014). Reliability of sagittal pelvic position assessments in standing, sitting and during hip flexion using palpation meter. *Journal of Bodywork and Movement Therapies*, 18(2), 210–214. <http://doi.org/10.1016/j.jbmt.2013.05.017>
- Barakatt, E. T., Romano, P. S., Riddle, D. L., & Beckett, L. A. (2009a). The Reliability of Maitland's Irritability Judgments in Patients with Low Back Pain. *The Journal of Manual & Manipulative Therapy*, 17(3), 135–140.
- Barakatt, E. T., Romano, P. S., Riddle, D. L., Beckett, L. A., & Kravitz, R. (2009b). An Exploration of Maitland's Concept of Pain Irritability in Patients with Low Back Pain. *The Journal of Manual & Manipulative Therapy*, 17(4), 196–205.
- Barakatt, E., Smidt, G. L., Dawson, J. D., Wei, S.-H., & Heiss, D. G. (1996). Interinnominate Motion and Symmetry: Comparison Between Gymnasts and Nongymnasts. *The Journal of Orthopaedic and Sports Physical Therapy*, 23(5), 309–319. <http://doi.org/10.2519/jospt.1996.23.5.309>
- Billis, E., Billis, E., McCarthy, C. J., McCarthy, C., Gliatis, J., Gliatis, J., et al. (2012). Inter-tester reliability of discriminatory examination items for sub-classifying non-specific low back pain. *Journal of Rehabilitation Medicine*, 44(10), 851–857. <http://doi.org/10.2340/16501977-0950>
- Binkley, J., Stratford, P. W., & Gill, C. (1995). Interrater reliability of lumbar accessory motion mobility testing. *Physical Therapy*, 75(9), 786–92; discussion 793–5.
- Blower, P. W., & Griffin, A. J. (1984). Clinical sacroiliac tests in ankylosing spondylitis and other causes of low back pain--2 studies. *Annals of the Rheumatic Diseases*, 43(2), 192–195.
- Boline, P. D., Haas, M., Meyer, J. J., Kassak, K., Nelson, C., & Keating, J. C. (1993). Interexaminer reliability of eight evaluative dimensions of lumbar segmental abnormality: Part II. *Journal of Manipulative and Physiological Therapeutics*, 16(6), 363–374.

- Brismée, J.-M., Atwood, K., Fain, M., Hodges, J., Sperle, A., Swaney, M., et al. (2005). Interrater Reliability of Palpation of Three-Dimensional Segmental Motion of the Lumbar Spine. *Journal of Manual & Manipulative Therapy*, 13(4), 215–220. <http://doi.org/10.1179/106698105790824761>
- Burdett, R. G., Brown, K. E., & Fall, M. P. (1986). Reliability and validity of four instruments for measuring lumbar spine and pelvic positions. *Physical Therapy*, 66(5), 677–684.
- Cakir, B., Richter, M., Käfer, W., Wieser, M., Puhl, W., & Schmidt, R. (2006). Evaluation of lumbar spine motion with dynamic X-ray--a reliability analysis. *Spine*, 31(11), 1258–1264. <http://doi.org/10.1097/01.brs.0000217763.80593.50>
- Carlsson, H., & Rasmussen-Barr, E. (2013). Clinical screening tests for assessing movement control in non-specific low-back pain. A systematic review of intra- and inter-observer reliability studies. *Manual Therapy*, 18(2), 103–110. <http://doi.org/10.1016/j.math.2012.08.004>
- Cibere, J., Thorne, A., Bellamy, N., Greidanus, N., Chalmers, A., Mahomed, N., et al. (2008). Reliability of the hip examination in osteoarthritis: Effect of standardization. *Arthritis and Rheumatism*, 59(3), 373–381. <http://doi.org/10.1002/art.23310>
- Cibulka, M. T., Delitto, A., & Koldehoff, R. M. (1988). Changes in innominate tilt after manipulation of the sacroiliac joint in patients with low back pain. An experimental study. *Physical Therapy*, 68(9), 1359–1363.
- Clare, H. A., Adams, R., & Maher, C. G. (2003). Reliability of detection of lumbar lateral shift. *Journal of Manipulative and Physiological Therapeutics*, 26(8), 476–480. [http://doi.org/10.1016/S0161-4754\(03\)00104-0](http://doi.org/10.1016/S0161-4754(03)00104-0)
- Cleary, C., Sanders, A. K., & Nick, T. G. (2007). Reliability of the Skin Compliance Device in the Assessment of Scar Pliability. *Journal of Hand Therapy*, 20(3), 232–238. <http://doi.org/10.1197/j.jht.2007.04.016>
- Cook, C., Brown, C., Michael, K., Isaacs, R., Howes, C., Richardson, W., et al. (2011). The clinical value of a cluster of patient history and observational findings as a diagnostic support tool for lumbar spine stenosis. *Physiotherapy Research International: the Journal for Researchers and Clinicians in Physical Therapy*, 16(3), 170–178. <http://doi.org/10.1002/pri.500>
- Cott, A., Parkinson, W., Bell, M. J., Adachi, J., Bédard, M., Cividino, A., & Bensen, W. (1992). Interrater reliability of the tender point criterion for fibromyalgia. *The Journal of Rheumatology*, 19(12), 1955–1959.
- Dankaerts, W., O'Sullivan, P. B., Straker, L. M., Burnett, A. F., & Skouen, J. S. (2006). The inter-examiner reliability of a classification method for non-specific chronic low back pain patients with motor control impairment. *Manual Therapy*, 11(1), 28–39. <http://doi.org/10.1016/j.math.2005.02.001>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/musculoskeletal medicine  
**Lumbar Region**

- Davis, A. M., Bridge, P., Miller, J., & Nelson-Wong, E. (2011). Interrater and intrarater reliability of the active hip abduction test. *The Journal of Orthopaedic and Sports Physical Therapy*, 41(12), 953–960. <http://doi.org/10.2519/jospt.2011.3684>
- Degenhardt, B. F., Johnson, J. C., Snider, K. T., & Snider, E. J. (2010). Maintenance and improvement of interobserver reliability of osteopathic palpatory tests over a 4-month period. *The Journal of the American Osteopathic Association*, 110(10), 579–586.
- Degenhardt, B. F., Snider, K. T., Snider, E. J., & Johnson, J. C. (2005a). Interobserver reliability of osteopathic palpatory diagnostic tests of the lumbar spine: improvements from consensus training. *The Journal of the American Osteopathic Association*, 105(10), 465–473.
- Degenhardt, B. F., Snider, K. T., Snider, E. J., & Johnson, J. C. (2005b). Interobserver reliability of osteopathic palpatory diagnostic tests of the lumbar spine: improvements from consensus training. *The Journal of the American Osteopathic Association*, 105(10), 465–473.
- Dennis, R. J., Finch, C. F., Elliott, B. C., & Farhart, P. J. (2008). The reliability of musculoskeletal screening tests used in cricket. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 9(1), 25–33. <http://doi.org/10.1016/j.ptsp.2007.09.004>
- Denteneer, L., Stassijns, G., De Hertogh, W., Truijen, S., & Van Daele, U. (2017). Inter- and Intrarater Reliability of Clinical Tests Associated With Functional Lumbar Segmental Instability and Motor Control Impairment in Patients With Low Back Pain: A Systematic Review. *Archives of Physical Medicine and Rehabilitation*, 98(1), 151–164.e6. <http://doi.org/10.1016/j.apmr.2016.07.020>
- Donahue, M. S., Riddle, D. L., & Sullivan, M. S. (1996). Intertester reliability of a modified version of McKenzie's lateral shift assessments obtained on patients with low back pain. *Physical Therapy*, 76(7), 706–16; discussion717–26.
- Dreyfuss, P., Dryer, S., Griffin, J., Hoffman, J., & Walsh, N. (1994). Positive sacroiliac screening tests in asymptomatic adults. *Spine*, 19(10), 1138–1143.
- Dreyfuss, P., Michaelsen, M., Pauza, K., McLarty, J., & Bogduk, N. (1996). The value of medical history and physical examination in diagnosing sacroiliac joint pain. *Spine*, 21(22), 2594–2602.
- Ekedahl, H., Jönsson, B., & Frobell, R. B. (2012). Fingertip-to-floor test and straight leg raising test: validity, responsiveness, and predictive value in patients with acute/subacute low back pain. *Archives of Physical Medicine and Rehabilitation*, 93(12), 2210–2215. <http://doi.org/10.1016/j.apmr.2012.04.020>

- Ekedahl, K. H., Jönsson, B., & Frobell, R. B. (2010). Validity of the fingertip-to-floor test and straight leg raising test in patients with acute and subacute low back pain: a comparison by sex and radicular pain. *Archives of Physical Medicine and Rehabilitation*, 91(8), 1243–1247. <http://doi.org/10.1016/j.apmr.2010.05.002>
- Elgueta-Cancino, E., Schabrun, S., Danneels, L., & Hodges, P. (2014). A clinical test of lumbopelvic control: development and reliability of a clinical test of dissociation of lumbopelvic and thoracolumbar motion. *Manual Therapy*, 19(5), 418–424. <http://doi.org/10.1016/j.math.2014.03.009>
- Enoch, F., Kjaer, P., Elkjaer, A., Remvig, L., & Juul-Kristensen, B. (2011). Inter-examiner reproducibility of tests for lumbar motor control. *BMC Musculoskeletal Disorders*, 12(1), 114. <http://doi.org/10.1186/1471-2474-12-114>
- Farshad-Amacker, N. A., Lurie, B., Herzog, R. J., & Farshad, M. (2014). Interreader and intermodality reliability of standard anteroposterior radiograph and magnetic resonance imaging in detection and classification of lumbosacral transitional vertebra. *The Spine Journal : Official Journal of the North American Spine Society*, 14(8), 1470–1475. <http://doi.org/10.1016/j.spinee.2013.08.048>
- Fedorak, C., Ashworth, N., Marshall, J., & Paull, H. (2003). Reliability of the visual assessment of cervical and lumbar lordosis: how good are we? *Spine*, 28(16), 1857–1859. <http://doi.org/10.1097/01.BRS.0000083281.48923.BD>
- Fortin, J. D., Dwyer, A. P., West, S., & Pier, J. (1994). Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part I: Asymptomatic volunteers. *Spine*, 19(13), 1475–1482.
- Freburger, J. K., & Riddle, D. L. (1999). Measurement of sacroiliac joint dysfunction: a multicenter intertester reliability study. *Physical Therapy*, 79(12), 1134–1141.
- French, S. D., Green, S., & Forbes, A. (2000). Reliability of chiropractic methods commonly used to detect manipulable lesions in patients with chronic low-back pain. *Journal of Manipulative and Physiological Therapeutics*, 23(4), 231–238.
- Fritz, J. M., & Piva, S. R. (2003). Physical impairment index: reliability, validity, and responsiveness in patients with acute low back pain. *Spine*, 28(11), 1189–1194. <http://doi.org/10.1097/01.BRS.0000067270.50897.DB>
- Fritz, J. M., Whitman, J. M., & Childs, J. D. (2005). Lumbar spine segmental mobility assessment: an examination of validity for determining intervention strategies in patients with low back pain. ... *Of Physical Medicine and Rehabilitation*.
- Gauvin, M. G., Riddle, D. L., & Rothstein, J. M. (1990). Reliability of clinical measurements of forward bending using the modified fingertip-to-floor method. *Physical Therapy*, 70(7), 443–447.

- Gerwin, R. D., Shannon, S., Hong, C. Z., Hubbard, D., & Gevirtz, R. (1997). Interrater reliability in myofascial trigger point examination. *Pain*, 69(1-2), 65–73.
- Gonnella, C., Paris, S. V., & Kutner, M. (1982). Reliability in evaluating passive intervertebral motion. *Physical Therapy*, 62(4), 436–444.
- H Charles Woodfield RPh, D. C., Woodfield, H. C., Gerstman, B. B., Burt Gerstman DVM, M. P., Olaisen, R. H., Renate Henry Olaisen DC, M., et al. (2011). Interexaminer Reliability of Supine Leg Checks for Discriminating Leg-Length Inequality. *Journal of Manipulative and Physiological Therapeutics*, 34(4), 239–246. <http://doi.org/10.1016/j.jmpt.2011.04.009>
- Habets, B., van Cingel, R. E. H., & Ostelo, R. W. J. G. (2015). Reproducibility of a battery of commonly used clinical tests to evaluate lumbopelvic motor control. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 16(4), 331–339. <http://doi.org/10.1016/j.ptsp.2015.02.004>
- Haig, A. J., Haig, A. J., Park, P., Henke, P. K., Henke, P. K., Yamakawa, K. S. J., et al. (2013). Reliability of the clinical examination in the diagnosis of neurogenic versus vascular claudication. *The Spine Journal : Official Journal of the North American Spine Society*, 13(12), 1826–1834. <http://doi.org/10.1016/j.spinee.2013.06.021>
- Haneline, M. T., & Young, M. (2009). A review of intraexaminer and interexaminer reliability of static spinal palpation: a literature synthesis. *Journal of Manipulative and Physiological Therapeutics*, 32(5), 379–386. <http://doi.org/10.1016/j.jmpt.2009.04.010>
- Harris-Hayes, M., & Van Dillen, L. R. (2009). The inter-tester reliability of physical therapists classifying low back pain problems based on the movement system impairment classification system. *PM & R : the Journal of Injury, Function, and Rehabilitation*, 1(2), 117–126. <http://doi.org/10.1016/j.pmrj.2008.08.001>
- Hawk, C., Phongphua, C., Bleecker, J., Swank, L., Lopez, D., & Rubley, T. (1999). Preliminary study of the reliability of assessment procedures for indications for chiropractic adjustments of the lumbar spine. *Journal of Manipulative and Physiological Therapeutics*, 22(6), 382–389.
- Hebert, J. J., Koppenhaver, S. L., Teyhen, D. S., Walker, B. F., & Fritz, J. M. (2013). The evaluation of lumbar multifidus muscle function via palpation: reliability and validity of a new clinical test. *The Spine Journal : Official Journal of the North American Spine Society*. <http://doi.org/10.1016/j.spinee.2013.08.056>
- Henriksen, M., Lund, H., Bliddal, H., & Danneskiold-Samsøe, B. (2006). Dynamic control of the lumbopelvic complex; lack of reliability of established test procedures. *European Spine Journal*, 16(6), 733–740. <http://doi.org/10.1007/s00586-006-0198-5>
- Heymann von, W., Moll, H., & Rauch, G. (2018). Study on sacroiliac joint diagnostics. *Manuelle Medizin*, (56), 239–248. <http://doi.org/10.1007/s00337-018-0405-6>

- Hicks, G. E., Fritz, J. M., Delitto, A., & Mishock, J. (2003). Interrater reliability of clinical examination measures for identification of lumbar segmental instability. *Archives of Physical Medicine and Rehabilitation*, 84(12), 1858–1864.
- Hidalgo, B., Benjamin Hidalgo PE, P. M. D., Hall, T., Toby Hall PT, M. P., Nielens, H., PhD, H. N. M., et al. (2014). Intertester agreement and validity of identifying lumbar pain provocative movement patterns using active and passive accessory movement tests. *Journal of Manipulative and Physiological Therapeutics*, 37(2), 105–115.  
<http://doi.org/10.1016/j.jmpt.2013.09.006>
- Holt, A. E., Shaw, N. J., Shetty, A., & Greenough, C. G. (2002). The reliability of the Low Back Outcome Score for back pain. *Spine*, 27(2), 206–210.
- Holtzman, G. (2011). Clinical Examination Procedures to Determine the Effect of Axial Decompression on Low Back Pain Symptoms in People With Chronic Low Back Pain. *The Journal of Orthopaedic and Sports Physical Therapy*.  
<http://doi.org/10.2519/jospt.2012.3724>
- Horneij, E., Hemborg, B., Johnsson, B., & Ekdahl, C. (2002). Clinical tests on impairment level related to low back pain: a study of test reliability. *Journal of Rehabilitation Medicine*, 34(4), 176–182.
- Hölmich, P., Hölmich, L. R., & Bjerg, A. M. (2004). Clinical examination of athletes with groin pain: an intraobserver and interobserver reliability study. *British Journal of Sports Medicine*, 38(4), 446–451. <http://doi.org/10.1136/bjsm.2003.004754>
- Hsieh, C. Y., Hong, C. Z., Adams, A. H., Platt, K. J., Danielson, C. D., Hoehler, F. K., & Tobis, J. S. (2000). Interexaminer reliability of the palpation of trigger points in the trunk and lower limb muscles. *Archives of Physical Medicine and Rehabilitation*, 81(3), 258–264.
- Hungerford, B. A., Gilleard, W., Moran, M., & Emmerson, C. (2007a). Evaluation of the Ability of Physical Therapists to Palpate Intrapelvic Motion With the Stork Test on the Support Side. *Physical Therapy*, 87(7), 879–887. <http://doi.org/10.2522/ptj.20060014>
- Hungerford, B. A., Gilleard, W., Moran, M., & Emmerson, C. (2007b). Evaluation of the ability of physical therapists to palpate intrapelvic motion with the Stork test on the support side. *Physical Therapy*, 87(7), 879–887. <http://doi.org/10.2522/ptj.20060014>
- Hunt, D. G., Zuberbier, O. A., Kozlowski, A. J., Robinson, J., Berkowitz, J., Schultz, I. Z., et al. (2001). Reliability of the lumbar flexion, lumbar extension, and passive straight leg raise test in normal populations embedded within a complete physical examination. *Spine*, 26(24), 2714–2718.

- Johansson, F. (2006). Interexaminer reliability of lumbar segmental mobility tests. *Manual Therapy*, 11(4), 331–336. <http://doi.org/10.1016/j.math.2005.06.014>
- Keating, J. C., Bergmann, T. F., Jacobs, G. E., Finer, B. A., & Larson, K. (1990). Interexaminer reliability of eight evaluative dimensions of lumbar segmental abnormality. *Journal of Manipulative and Physiological Therapeutics*, 13(8), 463–470.
- Kilpikoski, S., Airaksinen, O., Kankaanpää, M., Leminen, P., Videman, T., & Alen, M. (2002). Interexaminer reliability of low back pain assessment using the McKenzie method. *Spine*, 27(8), E207–14.
- Kim, H. W., Ko, Y. J., Rhee, W. I., Lee, J. S., Lim, J. E., Lee, S. J., et al. (2007). Interexaminer reliability and accuracy of posterior superior iliac spine and iliac crest palpation for spinal level estimations. *Journal of Manipulative and Physiological Therapeutics*, 30(5), 386–389. <http://doi.org/10.1016/j.jmpt.2007.04.005>
- Kokmeyer, D. J., Van der Wurff, P., Aufdemkampe, G., & Fickenscher, T. C. M. (2002). The reliability of multitest regimens with sacroiliac pain provocation tests. *Journal of Manipulative and Physiological Therapeutics*, 25(1), 42–48.
- Kwong, E. H., Virani, N., Robert, M., Gerry, K., Harding, A., Rose, M. S., et al. (2013). Inter-rater reliability of the Active Straight-Leg Raise and One-Leg Standing tests in non-pregnant women. *Journal of Rehabilitation Medicine : Official Journal of the UEMS European Board of Physical and Rehabilitation Medicine*, 45(10), 1058–1064. <http://doi.org/10.2340/16501977-1213>
- Landel, R., Kulig, K., Fredericson, M., Li, B., & Powers, C. M. (2008). Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing. *Physical Therapy*, 88(1), 43–49. <http://doi.org/10.2522/ptj.20060179>
- Laslett, M. (2008). Evidence-based diagnosis and treatment of the painful sacroiliac joint. *The Journal of Manual & Manipulative Therapy*.
- Laslett, M., & Williams, M. (1994). The reliability of selected pain provocation tests for sacroiliac joint pathology. *Spine*, 19(11), 1243–1249.
- Laslett, M., Aprill, C. N., McDonald, B., & Young, S. B. (2005). Diagnosis of Sacroiliac Joint Pain: Validity of individual provocation tests and composites of tests. *Manual Therapy*, 10(3), 207–218. <http://doi.org/10.1016/j.math.2005.01.003>
- Laslett, M., Öberg, B., Aprill, C. N., & McDonald, B. (2004). Zygapophysial joint blocks in chronic low back pain: a test of Revel's model as a screening test. *BMC Musculoskeletal Disorders*, 5, 43. <http://doi.org/10.1186/1471-2474-5-43>

- Laslett, M., Van der Wurff, P., Buijs, E., & Aprill, C. N. (2007). Comments on Berthelot et al. review: "Provocative sacroiliac joint maneuvers and sacroiliac joint block are unreliable for diagnosing sacroiliac joint pain." *Joint, Bone, Spine : Revue Du Rhumatisme*, 74(3), 306–7; author reply 307–8. <http://doi.org/10.1016/j.jbspin.2006.11.006>
- Laslett, M., Young, S. B., Aprill, C. N., & McDonald, B. (2003). Diagnosing painful sacroiliac joints: A validity study of a McKenzie evaluation and sacroiliac provocation tests. *The Australian Journal of Physiotherapy*, 49(2), 89–97.
- Latimer, J., Maher, C. G., Refshauge, K., & Colaco, I. (1999). The reliability and validity of the Biering- Sorensen test in asymptomatic subjects and subjects reporting current or previous nonspecific low back pain. *Spine*, 24(20), 2085–9— discussion 2090.
- Leard, J. S., EdD, J. S. L., PhD, B. A. C., Crane, B. A., PhD, K. A. B., & Ball, K. A. (2009). Intrarater and interrater reliability of 22 clinical measures associated with lower quarter malalignment. *Journal of Manipulative and Physiological Therapeutics*, 32(4), 270–276. <http://doi.org/10.1016/j.jmpt.2009.03.009>
- Levangie, P. K. (1999). Four clinical tests of sacroiliac joint dysfunction: the association of test results with innominate torsion among patients with and without low back pain. *Physical Therapy*, 79(11), 1043–1057.
- Levin, U., Nilsson-Wikmar, L., & Stenström, C. H. (2005). Variability Within and Between Evaluations of Sacroiliac Pain With the Use of Distraction Testing. *Journal of Manipulative and Physiological Therapeutics*, 28(9), 688–695. <http://doi.org/10.1016/j.jmpt.2005.09.016>
- Levin, U., Nilsson-Wikmar, L., Stenström, C. H., & Lundeberg, T. (1998). Reproducibility of manual pressure force on provocation of the sacroiliac joint. *Physiotherapy Research International : the Journal for Researchers and Clinicians in Physical Therapy*, 3(1), 1–14.
- Lewit, K., & Rosina, A. (1999). Why yet another diagnostic sign of sacroiliac movement restriction? *Journal of Manipulative and Physiological Therapeutics*, 22(3), 154–160.
- Ljungquist, T., Harms-Ringdahl, K., Nygren, A., & Jensen, I. (1999). Intra- and inter-rater reliability of an 11-testpackage for assessing dysfunction due to back or neck pain. *Physiotherapy Research International : the Journal for Researchers and Clinicians in Physical Therapy*, 4(3), 214–232.
- Luomajoki, H., Kool, J., de Bruin, E. D., & Airaksinen, O. (2007). Reliability of movement control tests in the lumbar spine. *BMC Musculoskeletal Disorders*, 8, 90. <http://doi.org/10.1186/1471-2474-8-90>
- Mac-Thiong, J.-M., Labelle, H., Parent, S., Poitras, B., Jodoin, A., Ouellet, J., & Duong, L. (2007). Assessment of sacral doming in lumbosacral spondylolisthesis. *Spine*, 32(17), 1888–1895. <http://doi.org/10.1097/BRS.0b013e31811ebaa1>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
**Lumbar Region**

- Maher, C. G., Latimer, J., & Adams, R. (1998). An investigation of the reliability and validity of posteroanterior spinal stiffness judgments made using a reference-based protocol. *Physical Therapy*, 78(8), 829–837.
- Martin, R. L., & Sekiya, J. K. (2008). The interrater reliability of 4 clinical tests used to assess individuals with musculoskeletal hip pain. *The Journal of Orthopaedic and Sports Physical Therapy*, 38(2), 71–77. <http://doi.org/10.2519/jospt.2008.38.1.A71>
- Meijne, W., van Neerbos, K., Aufdemkampe, G., & van der Wurff, P. (1999). Intraexaminer and interexaminer reliability of the Gillet test. *Journal of Manipulative and Physiological Therapeutics*, 22(1), 4–9.
- Monnier, A., Heuer, J., Norman, K., & Äng, B. O. (2012). Inter- and intra-observer reliability of clinical movement control tests for marines. *BMC Musculoskeletal Disorders*, 13(1), 263. <http://doi.org/10.1186/1471-2474-13-263>
- Murphy, D. R., Byfield, D., McCarthy, P., Humphreys, K., Gregory, A. A., & Rochon, R. (2006). Interexaminer reliability of the hip extension test for suspected impaired motor control of the lumbar spine. *Journal of Manipulative and Physiological Therapeutics*, 29(5), 374–377. <http://doi.org/10.1016/j.jmpt.2006.04.012>
- O'Connor, S., McCaffrey, N., Whyte, E., & Moran, K. (2016). The development and reliability of a simple field based screening tool to assess core stability in athletes. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 20, 40–44. <http://doi.org/10.1016/j.ptsp.2015.12.003>
- O'Haire, C., & Gibbons, P. (2000). Inter-examiner and intra-examiner agreement for assessing sacroiliac anatomical landmarks using palpation and observation: pilot study. *Manual Therapy*, 5(1), 13–20. <http://doi.org/10.1054/math.1999.0203>
- Paatelma, M., Karvonen, E., & Heinonen, A. (2010). Inter- and intra-tester reliability of selected clinical tests in examining patients with early phase lumbar spine and sacroiliac joint pain and dysfunction. *Advances in Physiotherapy*, 12(2), 74–80. <http://doi.org/10.3109/14038190903582154>
- Pal, P., Milosavljevic, S., Sole, G., & Johnson, G. (2007). Hip and lumbar continuous motion characteristics during flexion and return in young healthy males. *European Spine Journal*, 16(6), 741–747. <http://doi.org/10.1007/s00586-006-0200-2>
- Perret, C., Poiraudau, S., Fermanian, J., & Revel, M. (2001a). Pelvic mobility when bending forward in standing position: validity and reliability of 2 motion analysis devices. *Archives of Physical Medicine and Rehabilitation*, 82(2), 221–226. <http://doi.org/10.1053/apmr.2001.18217>
- Perret, C., Poiraudau, S., Fermanian, J., Colau, M. M., Benhamou, M. A., & Revel, M. (2001b). Validity, reliability, and responsiveness of the fingertip-to-floor test. *Archives of Physical Medicine and Rehabilitation*, 82(11), 1566–1570. <http://doi.org/10.1053/apmr.2001.26064>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/musculoskeletal medicine  
**Lumbar Region**

- Petersen, T., Olsen, S., Laslett, M., Thorsen, H., Manniche, C., Ekdahl, C., & Jacobsen, S. (2004). Inter- tester reliability of a new diagnostic classification system for patients with non-specific low back pain. *The Australian Journal of Physiotherapy*, 50(2), 85–94.
- Peul, W. C., Brand, R., Thomeer, R. T. W. M., & Koes, B. W. (2008). Influence of gender and other prognostic factors on outcome of sciatica. *Pain*, 138(1), 180–191. <http://doi.org/10.1016/j.pain.2007.12.014>
- Potter, L., McCarthy, C., & Oldham, J. (2006). Intraexaminer reliability of identifying a dysfunctional segment in the thoracic and lumbar spine. *Journal of Manipulative and Physiological Therapeutics*, 29(3), 203–207. <http://doi.org/10.1016/j.jmpt.2006.01.005>
- Prather, H., Harris-Hayes, M., Hunt, D. M., Steger-May, K., Mathew, V., & Clohisey, J. C. (2010a). Reliability and Agreement of Hip Range of Motion and Provocative Physical Examination Tests in Asymptomatic Volunteers. *PMRJ*, 2(10), 888–895. <http://doi.org/10.1016/j.pmrj.2010.05.005>
- Prather, H., Harris-Hayes, M., Hunt, D. M., Steger-May, K., Mathew, V., & Clohisey, J. C. (2010b). Reliability and agreement of hip range of motion and provocative physical examination tests in asymptomatic volunteers. *PMRJ*, 2(10), 888–895. <http://doi.org/10.1016/j.pmrj.2010.05.005>
- Qvistgaard, E., Rasmussen, J., Lætgaard, J., Hecksher-Sørensen, S., & Bliddal, H. (2006). Intra-observer and interobserver agreement of the manual examination of the lumbar spine in chronic low-back pain. *European Spine Journal*, 16(2), 277–282. <http://doi.org/10.1007/s00586-006-0134-8>
- Riddle, D. L., & Freburger, J. K. (2002a). Evaluation of the presence of sacroiliac joint region dysfunction using a combination of tests: a multicenter intertester reliability study. *Physical Therapy*, 82(8), 772–781.
- Robinson, H. S., & Mengshoel, A. M. (2014). Assessments of lumbar flexion range of motion: intertester reliability and concurrent validity of 2 commonly used clinical tests. *Spine*, 39(4), E270–5. <http://doi.org/10.1097/BRS.0000000000000131>
- Robinson, H. S., Brox, J. I., Robinson, R., Bjelland, E., Solem, S., & Telje, T. (2007). The reliability of selected motion and pain provocation tests for the sacroiliac joint. *Manual Therapy*, 12(1), 72–79. <http://doi.org/10.1016/j.math.2005.09.004>
- Robinson, R., Robinson, H. S., Bjørke, G., & Kvale, A. (2009). Reliability and validity of a palpation technique for identifying the spinous processes of C7 and L5. *Manual Therapy*, 14(4), 409–414. <http://doi.org/10.1016/j.math.2008.06.002>
- Rose, du, A., & Breen, A. (2016). Relationships between lumbar inter-vertebral motion and lordosis in healthy adult males: a cross sectional cohort study. *BMC Musculoskeletal Disorders*, 17, 121. <http://doi.org/10.1186/s12891-016-0975-1>

- Roussel, N. A., Nijs, J., Truijen, S., Smeuninx, L., & Stassijns, G. (2007). Low back pain: clinimetric properties of the Trendelenburg test, active straight leg raise test, and breathing pattern during active straight leg raising. *Journal of Manipulative and Physiological Therapeutics*, 30(4), 270–278. <http://doi.org/10.1016/j.jmpt.2007.03.001>
- Sc, M. D. M., Deore, M., PhD, S. M., & May, S. (2012). The inter-rater and intra-rater reliability of passive physiological accessory movement assessment of lumbar spine in novice manual therapists. *Journal of Bodywork and Movement Therapies*, 16(3), 289–293. <http://doi.org/10.1016/j.jbmt.2011.12.006>
- Schäfer, A., Hall, T. M., Lüdtke, K., Mallwitz, J., & Briffa, N. K. (2009). Interrater reliability of a new classification system for patients with neural low back-related leg pain. *The Journal of Manual & Manipulative Therapy*, 17(2), 109–117.
- Schneider, M., Erhard, R., Brach, J., Tellin, W., Imbarlina, F., & Delitto, A. (2008). Spinal palpation for lumbar segmental mobility and pain provocation: an interexaminer reliability study. *Journal of Manipulative and Physiological Therapeutics*, 31(6), 465–473. <http://doi.org/10.1016/j.jmpt.2008.06.004>
- Schultz, S. M., & Jacobs, M. M. (2015). Reliability of Entry-Level Athletic Trainers' Palpation Skills of BonyAnatomical Landmarks in the Lumbopelvic Region. *Athletic Training Education Journal*, 10(4), 296–301. <http://doi.org/10.4085/1004296>
- Sedaghat, N., Latimer, J., Maher, C., & Wisbey-Roth, T. (2007). The reproducibility of a clinical grading system of motor control in patients with low back pain. *Journal of Manipulative and Physiological Therapeutics*, 30(7), 501–508. <http://doi.org/10.1016/j.jmpt.2007.07.008>
- Seffinger, M. A., Najm, W. I., Mishra, S. I., Adams, A., Dickerson, V. M., Murphy, L. S., & Reinsch, S. (2004). Reliability of Spinal Palpation for Diagnosis of Back and Neck Pain. *Spine*, 29(19), E413–E425. <http://doi.org/10.1097/01.brs.0000141178.98157.8e>
- Snider, K. T., DO, E. J. S., DO, B. F. D., MA, J. C. J., & DO, J. W. K. (2011). Palpatory Accuracy of Lumbar Spinous Processes Using Multiple Bony Landmarks. *Journal of Manipulative and Physiological Therapeutics*, 34(5), 306–313. <http://doi.org/10.1016/j.jmpt.2011.04.006>
- Snider, K. T., Johnson, J. C., Degenhardt, B. F., Snider, E. J., & Burton, D. C. (2014). Association of low back pain, somatic dysfunction, and lumbar bone mineral density: reproducibility of findings. *The Journal of the American Osteopathic Association*, 114(5), 356–367. <http://doi.org/10.7556/jaoa.2014.073>
- Stovall, B. A., & Kumar, S. (2010a). Reliability of bony anatomic landmark asymmetry assessment in the lumbopelvic region: application to osteopathic medical education. *The Journal of the American Osteopathic Association*, 110(11), 667–674.

- Stovall, B. A., & Kumar, S. (2010b). Reliability of bony anatomic landmark asymmetry assessment in the lumbopelvic region: application to osteopathic medical education. *The Journal of the American Osteopathic Association*, 110(11), 667–674.  
<http://doi.org/10.1016/j.apmr.2010.12.042>
- Stovall, B. A., Bae, S., & Kumar, S. (2010). Anterior superior iliac spine asymmetry assessment on a novel pelvic model: an investigation of accuracy and reliability. *Journal of Manipulative and Physiological Therapeutics*, 33(5), 378–385.  
<http://doi.org/10.1016/j.jmpt.2010.05.005>
- Sutlive, T. G., Lopez, H. P., Schnitker, D. E., Yawn, S. E., Halle, R. J., Mansfield, L. T., et al. (2008). Development of a clinical prediction rule for diagnosing hip osteoarthritis in individuals with unilateral hip pain. *The Journal of Orthopaedic and Sports Physical Therapy*, 38(9), 542–550. <http://doi.org/10.2519/jospt.2008.2753>
- Tidstrand, J., & Horneij, E. (2009). Inter-rater reliability of three standardized functional tests in patients with low back pain. *BMC Musculoskeletal Disorders*, 10, 58.  
<http://doi.org/10.1186/1471-2474-10-58>
- Tong, H. C., Heyman, O. G., Lado, D. A., & Isser, M. M. (2006). Interexaminer reliability of three methods of combining test results to determine side of sacral restriction, sacral base position, and innominate bone position. *The Journal of the American Osteopathic Association*, 106(8), 464–468.
- Trudelle-Jackson, E., Sarvaiya-Shah, S. A., & Wang, S. S. (2008). Interrater reliability of a movement impairmentbased classification system for lumbar spine syndromes in patients with chronic low back pain. *Journal of Orthopaedic & ...*, 38(6), 371–376.  
<http://doi.org/10.2519/jospt.2008.2760>
- van den Hoogen, H. J., Koes, B. W., Devillé, W., van Eijk, J. T., & Bouter, L. M. (1996). The inter-observer reproducibility of Lasègue's sign in patients with low back pain in general practice. *British Journal of General Practice*, 46(413), 727–730.
- Van Deursen, L., Patijn, J., & Ockhuysen, A. (1990). The value of some clinical tests of the sacroiliac joint. *Journal of Manual Medicine*.
- Van Dillen, L. R., Sahrman, S. A., Norton, B. J., Caldwell, C. A., Fleming, D. A., McDonnell, M. K., & Woolsey, N. B. (1998). Reliability of physical examination items used for classification of patients with low back pain. *Physical Therapy*, 78(9), 979–988.
- van Kessel-Cobelens, A. M., Verhagen, A. P., Mens, J. M., Snijders, C. J., & Koes, B. W. (2008a). Pregnancy-Related Pelvic Girdle Pain: Interrater Reliability of 3 Tests to Determine Asymmetric Mobility of the Sacroiliac Joints. *Journal of Manipulative and Physiological Therapeutics*, 31(2), 130–136. <http://doi.org/10.1016/j.jmpt.2007.12.003>

- van Kessel-Cobelens, A. M., Verhagen, A. P., Mens, J. M., Snijders, C. J., & Koes, B. W. (2008b). Pregnancy-related pelvic girdle pain: intertester reliability of 3 tests to determine asymmetric mobility of the sacroiliac joints. *Journal of Manipulative and Physiological Therapeutics*, 31(2), 130–136. <http://doi.org/10.1016/j.jmpt.2007.12.003>
- van Middendorp, J. J., Patel, A. A., Schuetz, M., & Joaquim, A. F. (2013). The precision, accuracy and validity of detecting posterior ligamentous complex injuries of the thoracic and lumbar spine: a critical appraisal of the literature. *European Spine Journal*, 22(3), 461–474. <http://doi.org/10.1007/s00586-012-2602-7>
- Vincent-Smith, B., & Gibbons, P. (1999). Inter-examiner and intra-examiner reliability of the standing flexion test. *Manual Therapy*, 4(2), 87–93. <http://doi.org/10.1054/math.1999.0173>
- Walsh, J., & Hall, T. (2009). Reliability, validity and diagnostic accuracy of palpation of the sciatic, tibial and common peroneal nerves in the examination of low back related leg pain. *Manual Therapy*, 14(6), 623–629. <http://doi.org/10.1016/j.math.2008.12.007>

## Shoulder Region

- Al-Shenqiti, A. M., & Oldham, J. A. (2005). Test-retest reliability of myofascial trigger point detection in patients with rotator cuff tendonitis. *Clinical Rehabilitation*, 19(5), 482–487.
- Ambrad-Chalela, E., Thomas, G. I., & Johansen, K. H. (2004). Recurrent neurogenic thoracic outlet syndrome. *American Journal of Surgery*, 187(4), 505–510. <http://doi.org/10.1016/j.amjsurg.2003.12.050>
- Atasoy, E. (1997). Thoracic outlet compression syndrome caused by a schwannoma of the C7 nerve root. *Journal of Hand Surgery (Edinburgh, Scotland)*, 22(5), 662–663.
- Atasoy, E. (2010). A hand surgeon's further experience with thoracic outlet compression syndrome. *The Journal of Hand Surgery*, 35(9), 1528–1538. <http://doi.org/10.1016/j.jhsa.2010.06.025>
- Athanassiadi, K., Kalavrouziotis, G., Karydakis, K., & Bellenis, I. (2001). Treatment of thoracic outlet syndrome: long-term results. *World Journal of Surgery*, 25(5), 553–557. <http://doi.org/10.1007/s002680020079>
- Baertschi, E., Swanenburg, J., Brunner, F., & Kool, J. (2013). Interrater reliability of clinical tests to evaluate scapulothoracic motion. *BMC Musculoskeletal Disorders*, 14, 315. <http://doi.org/10.1186/1471-2474-14-315>
- Bailey, P. (1985). Surfer's rib: isolated first rib fracture secondary to indirect trauma. *Annals of Emergency Medicine*, 14(4), 346–349.
- Bateman, J. E. (1983). Neurologic painful conditions affecting the shoulder. *Clinical Orthopaedics and Related Research*, (173), 44–54.
- Ben Kibler, W., Sciascia, A. D., Hester, P., Dome, D., & Jacobs, C. (2009). Clinical utility of traditional and new tests in the diagnosis of biceps tendon injuries and superior labrum anterior and posterior lesions in the shoulder. *The American Journal of Sports Medicine*, 37(9), 1840–1847. <http://doi.org/10.1177/0363546509332505>
- Bertilson, B. C., Grunnesjö, M., & Strender, L.-E. (2003). Reliability of Clinical Tests in the Assessment of Patients With Neck/Shoulder Problems—Impact of History. *Spine*, 28(19), 2222–2231. <http://doi.org/10.1097/01.BRS.0000089685.55629.2E>
- Bialocerkowski, A. E., & Galea, M. (2006). Comparison of visual and objective quantification of elbow and shoulder movement in children with obstetric brachial plexus palsy. *Journal of Brachial Plexus and Peripheral Nerve Injury*, 1, 5. <http://doi.org/10.1186/1749-7221-1-5>
- Boezaart, A. P., Haller, A., Laduzenski, S., Koyyalamudi, V. B., Ihnatsenka, B., & Wright, T. (2010). Neurogenic thoracic outlet syndrome: A case report and review of the literature. *International Journal of Shoulder Surgery*, 4(2), 27–35. <http://doi.org/10.4103/0973-6042.70817>

- Booth, C. M., & Roper, B. A. (1979). Chronic dislocation of the sternoclavicular joint: an operative repair. *Clinical Orthopaedics and Related Research*, (140), 17–20.
- Borstad, J. D., Dashottar, A., & Stoughton, T. (2015). Validity and reliability of the Low Flexion measurement for posterior glenohumeral joint capsule tightness. *Manual Therapy*, 20(6), 875–878. <http://doi.org/10.1016/j.math.2015.08.007>
- Borstad, J. D., Mathiowetz, K. M., Minday, L. E., Prabhu, B., Christopherson, D. E., & Ludewig, P. M. (2007). Clinical measurement of posterior shoulder flexibility. *Manual Therapy*, 12(4), 386–389. <http://doi.org/10.1016/j.math.2006.07.014>
- BRANDT, C., SOLE, G., KRAUSE, M., & NEL, M. (2007). An evidence-based review on the validity of the Kaltenborn rule as applied to the glenohumeral joint. *Manual Therapy*, 12(1), 3–11. <http://doi.org/10.1016/j.math.2006.02.011>
- Bron, C., Franssen, J., Wensing, M., & Oostendorp, R. A. B. (2007). Interrater reliability of palpation of myofascial trigger points in three shoulder muscles. *The Journal of Manual & Manipulative Therapy*, 15(4), 203–215.
- Brown, C. (1983). Compressive, invasive referred pain to the shoulder. *Clinical Orthopaedics and Related Research*, (173), 55–62.
- Bruyn, G. A. W., Pineda, C., Hernandez-Díaz, C., Ventura-Rios, L., Moya, C., Garrido, J., et al. (2010). Validity of ultrasonography and measures of adult shoulder function and reliability of ultrasonography in detecting shoulder synovitis in patients with rheumatoid arthritis using magnetic resonance imaging as a gold standard. *Arthritis Care & Research*, 62(8), 1079–1086. <http://doi.org/10.1002/acr.20175>
- Burns, S. A., Cleland, J. A., Carpenter, K., & Mintken, P. E. (2016). Interrater reliability of the cervicothoracic and shoulder physical examination in patients with a primary complaint of shoulder pain. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 18, 46–55. <http://doi.org/10.1016/j.ptsp.2015.07.001>
- Cadogan, A., Laslett, M., Hing, W., McNair, P., & Williams, M. (2011). Interexaminer reliability of orthopaedic special tests used in the assessment of shoulder pain. *Manual Therapy*, 16(2), 131–135. <http://doi.org/10.1016/j.math.2010.07.009>
- Cadogan, A., McNair, P., Laslett, M., & Hing, W. (2013). Shoulder pain in primary care: diagnostic accuracy of clinical examination tests for non-traumatic acromioclavicular joint pain. *BMC Musculoskeletal Disorders*, 14, 156. <http://doi.org/10.1186/1471-2474-14-156>
- Caliş, M., Akgün, K., Birtane, M., Karacan, I., Caliş, H., & Tüzün, F. (2000). Diagnostic values of clinical diagnostic tests in subacromial impingement syndrome. *Annals of the Rheumatic Diseases*, 59(1), 44–47.

- Calvert, E., Chambers, G. K., Regan, W., Hawkins, R. H., & Leith, J. M. (2009). Special physical examination tests for superior labrum anterior posterior shoulder tears are clinically limited and invalid: a diagnostic systematic review. *Journal of Clinical Epidemiology*, 62(5), 558–563. <http://doi.org/10.1016/j.jclinepi.2008.04.010>
- Carbone, S., Moroder, P., Runer, A., Resch, H., Gumina, S., & Hertel, R. (2016). Scapular dyskinesis after Latarjet procedure. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 25(3), 422–427. <http://doi.org/10.1016/j.jse.2015.08.001>
- Carter, T., Hall, H., McIntosh, G., Murphy, J., MacDougall, J., & Boyle, C. (2012). Intertester reliability of a classification system for shoulder pain. *Physiotherapy*, 98(1), 40–46. <http://doi.org/10.1016/j.physio.2010.12.003>
- Chesworth, B. M., MacDermid, J. C., Roth, J. H., & Patterson, S. D. (1998). Movement diagram and “end-feel” reliability when measuring passive lateral rotation of the shoulder in patients with shoulder pathology. *Physical Therapy*, 78(6), 593–601.
- Chronopoulos, E., Kim, T. K., Park, H. B., Ashenbrenner, D., & McFarland, E. G. (2004). Diagnostic value of physical tests for isolated chronic acromioclavicular lesions. *The American Journal of Sports Medicine*, 32(3), 655–661.
- Colli, B. O., Carlotti, C. G., Assirati, J. A., & Marques, W. (2006). Neurogenic thoracic outlet syndromes: a comparison of true and nonspecific syndromes after surgical treatment. *Surgical Neurology*, 65(3), 262–71– discussion 271–2. <http://doi.org/10.1016/j.surneu.2005.06.037>
- Collin, P., Treseder, T., Denard, P. J., Neyton, L., Walch, G., & Lädermann, A. (2015). What is the Best Clinical Test for Assessment of the Teres Minor in Massive Rotator Cuff Tears? *Clinical Orthopaedics and Related Research*, 473(9), 2959–2966. <http://doi.org/10.1007/s11999-015-4392-9>
- Colosimo, A. J., Byrne, E., Heidt, R. S., Carlonas, R. L., & Wyatt, H. (2004). Acute traumatic first-rib fracture in the contact athlete: a case report. *The American Journal of Sports Medicine*, 32(5), 1310–1312. <http://doi.org/10.1177/0363546503261352>
- De Beer, L., & Kroon, H. M. (2012). [A man with progressive neck and shoulder pain]. *Nederlands Tijdschrift Voor Geneeskunde*, 156(39), A3978.
- De Wilde, L., Plasschaert, F., Berghs, B., Van Hoecke, M., Verstraete, K., & Verdonk, R. (2003). Intra- and interexaminer reliability of four manual shoulder maneuvers used to identify subacromial pain. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 12(4), 346–349. <http://doi.org/10.1016/mse.2003.S1058274603000387>
- De Winter, A. F., Jans, M. P., Scholten, R. J., Devillé, W., van Schaardenburg, D., & Bouter, L. M. (1999). Diagnostic classification of shoulder disorders: interobserver agreement and determinants of disagreement. *Annals of the Rheumatic Diseases*, 58(5), 272–277.

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
Shoulder Region

- Dinnes, J., Loveman, E., McIntyre, L., & Waugh, N. (2003). The effectiveness of diagnostic tests for the assessment of shoulder pain due to soft tissue disorders: a systematic review. *Health Technology Assessment (Winchester, England)*, 7(29), iii–1–166.
- DROMERICK, A., Kumar, A., VOLSHTEYN, O., & EDWARDS, D. (2006). Hemiplegic Shoulder Pain Syndrome: Interrater Reliability of Physical Diagnosis Signs. *Archives of Physical Medicine and Rehabilitation*, 87(2), 294–295. <http://doi.org/10.1016/j.apmr.2005.10.004>
- Edwards, T. (2002). Interobserver and intraobserver reliability of the measurement of shoulder internal rotation by vertebral level. *Journal of Shoulder and Elbow Surgery*, 11(1), 40–42. <http://doi.org/10.1067/mse.2002.119853>
- Ellenbecker, T. S., Bailie, D. S., Mattalino, A. J., Carfagno, D. G., Wolff, M. W., Brown, S. W., & Kulikowich, J. M. (2002). Intrarater and interrater reliability of a manual technique to assess anterior humeral head translation of the glenohumeral joint. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 11(5), 470–475.
- Farber, A. J. (2006). Clinical Assessment of Three Common Tests for Traumatic Anterior Shoulder Instability. *The Journal of Bone and Joint Surgery*, 88(7), 1467–1474. <http://doi.org/10.2106/BJJS.E.00594>
- Gill, H. S., Rassi, El, G., Bahk, M. S., Castillo, R. C., & McFarland, E. G. (2007). Physical examination for partial tears of the biceps tendon. *The American Journal of Sports Medicine*, 35(8), 1334–1340. <http://doi.org/10.1177/0363546507300058>
- Gross, M. L., & Distefano, M. C. (1997). Anterior release test. A new test for occult shoulder instability. *Clinical Orthopaedics and Related Research*, (339), 105–108.
- Guanche, C. A., & Jones, D. C. (2003). Clinical testing for tears of the glenoid labrum. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 19(5), 517–523. <http://doi.org/10.1053/jars.2003.50104>
- Guerra-Soriano, F., Ruiz-Suárez, M., Encalada-Díaz, M. I., Pérez-Domenech, J., Moscona-Mishy, L., & Valero-González, F. S. (2010). [The Seebauer classification for the staging of arthropathy due to rotator cuff massive tear: intra- and interobserver concordance analysis]. *Acta Ortopédica Mexicana*, 24(6), 390–394.
- Hanchard, N. C. A., Howe, T. E., & Gilbert, M. M. (2005). Diagnosis of shoulder pain by history and selective tissue tension: agreement between assessors. *The Journal of Orthopaedic and Sports Physical Therapy*, 35(3), 147–153.
- Hansson, G. A., Balogh, I., Byström, J. U., Ohlsson, K., Nordander, C., Asterland, P., et al. (2001). Questionnaire versus direct technical measurements in assessing postures and movements of the head, upper back, arms and hands. *Scandinavian Journal of Work, Environment & Health*, 27(1), 30–40.

- Hayes, K. (2002). Reliability of 3 methods for assessing shoulder strength. *Journal of Shoulder and Elbow Surgery*, 11(1), 33–39. <http://doi.org/10.1067/mse.2002.119852>
- Hayes, K. W., & Petersen, C. M. (2001). Reliability of assessing end-feel and pain and resistance sequence in subjects with painful shoulders and knees. *The Journal of Orthopaedic and Sports Physical Therapy*, 31(8), 432–445.
- Hayes, K. W., & Petersen, C. M. (2003). Reliability of classifications derived from Cyriax's resisted testing in subjects with painful shoulders and knees. *The Journal of Orthopaedic and Sports Physical Therapy*, 33(5), 235–246.
- Hayes, K., Walton, J. R., Szomor, Z. L., & Murrell, G. A. C. (2002). Reliability of 3 methods for assessing shoulder strength. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 11(1), 33–39. <http://doi.org/10.1067/mse.2002.119852>
- Hayes, K., Walton, J. R., Szomor, Z. R., & Murrell, G. A. (2001). Reliability of five methods for assessing shoulder range of motion. *The Australian Journal of Physiotherapy*, 47(4), 289–294.
- Hegedus, E. J., Cook, C., Lewis, J., Wright, A., & Park, J. Y. (2015). Combining orthopedic special tests to improve diagnosis of shoulder pathology. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 16(2), 87–92. <http://doi.org/10.1016/j.ptsp.2014.08.001>
- Hegedus, E. J., Goode, A. P., Cook, C. E., Michener, L., Myer, C. A., Myer, D. M., & Wright, A. A. (2012). Which physical examination tests provide clinicians with the most value when examining the shoulder? Update of a systematic review with meta-analysis of individual tests. *British Journal of Sports Medicine*, 46(14), 964–978. <http://doi.org/10.1136/bjsports-2012-091066>
- Hertel, R., Ballmer, F. T., Lombert, S. M., & Gerber, C. (1996). Lag signs in the diagnosis of rotator cuff rupture. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 5(4), 307–313.
- Hertel, R., Lambert, S. M., & Ballmer, F. T. (1998). The deltoid extension lag sign for diagnosis and grading of axillary nerve palsy. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 7(2), 97–99.
- HICKEY, B., MILOSAVLJEVIC, S., BELL, M., & MILBURN, P. (2007). Accuracy and reliability of observational motion analysis in identifying shoulder symptoms. *Manual Therapy*, 12(3), 263–270. <http://doi.org/10.1016/j.math.2006.05.005>
- Hodler, J. (1996). [Diagnosis of shoulder impingement syndrome]. *Der Radiologe*, 36(12), 944–950.

- Holtby, R., & Razmjou, H. (2004a). Accuracy of the Speed's and Yergason's tests in detecting biceps pathology and SLAP lesions: comparison with arthroscopic findings. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 20(3), 231–236. <http://doi.org/10.1016/j.arthro.2004.01.008>
- Holtby, R., & Razmjou, H. (2004b). Validity of the supraspinatus test as a single clinical test in diagnosing patients with rotator cuff pathology. *The Journal of Orthopaedic and Sports Physical Therapy*, 34(4), 194–200.
- Huang, T.-S., Huang, H.-Y., Wang, T.-G., Tsai, Y.-S., & Lin, J.-J. (2015). Comprehensive classification test of scapular dyskinesis: A reliability study. *Manual Therapy*, 20(3), 427–432. <http://doi.org/10.1016/j.math.2014.10.017>
- Ilg, A., Bankes, M. J., & Emery, R. J. (2001). The intra- and inter-observer reliability of the Samilson and Prieto grading system of glenohumeral arthropathy. *Knee Surgery, Sports Traumatology, Arthroscopy : Official Journal of the ESSKA*, 9(3), 187–190.
- Jia, X., Yokota, A., McCarty, E. C., Nicholson, G. P., Weber, S. C., McMahon, P. J., et al. (2011). Reproducibility and Reliability of the Snyder Classification of Superior Labral Anterior Posterior Lesions Among Shoulder Surgeons. *The American Journal of Sports Medicine*, 39(5), 986–991. <http://doi.org/10.1177/0363546510392332>
- Johansson, K. M., & Adolfsson, L. E. (2005). Intraobserver and interobserver reliability for the strength test in the Constant-Murley shoulder assessment. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 14(3), 273–278. <http://doi.org/10.1016/j.jse.2004.08.001>
- Johansson, K., & Ivarson, S. (2009). Intra- and interexaminer reliability of four manual shoulder maneuvers used to identify subacromial pain. *Manual Therapy*, 14(2), 231–239. <http://doi.org/10.1016/j.math.2008.03.003>
- Juul-Kristensen, B., Hilt, K., Enoch, F., Remvig, L., & Sjøgaard, G. (2011). Scapular dyskinesis in trapezius myalgia and intraexaminer reproducibility of clinical tests. *Physiotherapy Theory and Practice*, 27(7), 492–502. <http://doi.org/10.3109/09593985.2010.528548>
- Juul-Kristensen, B., Røgind, H., Jensen, D. V., & Remvig, L. (2007). Inter-examiner reproducibility of tests and criteria for generalized joint hypermobility and benign joint hypermobility syndrome.
- *Rheumatology (Oxford, England)*, 46(12), 1835–1841. <http://doi.org/10.1093/rheumatology/kem290>
- Kappe, T., Cakir, B., Reichel, H., & Elsharkawi, M. (2011). Reliability of radiologic classification for cuff tear arthropathy. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 20(4), 543–547. <http://doi.org/10.1016/j.jse.2011.01.012>

- Kappe, T., Knappe, K., Elsharkawi, M., Reichel, H., & Cakir, B. (2015). Predictive value of preoperative clinical examination for subacromial decompression in impingement syndrome. *Knee Surgery, Sports Traumatology, Arthroscopy : Official Journal of the ESSKA*, 23(2), 443–448. <http://doi.org/10.1007/s00167-013-2386-2>
- Karlsson, J. (2010). Physical examination tests are not valid for diagnosing SLAP tears: a review. *Clinical Journal of Sport Medicine : Official Journal of the Canadian Academy of Sport Medicine*, 20(2), 134– 135. <http://doi.org/10.1097/01.jsm.0000369406.54312.0b>
- Kibler, W. B. (1995). Specificity and sensitivity of the anterior slide test in throwing athletes with superior glenoid labral tears. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 11(3), 296–300.
- Kibler, W. B., Sciascia, A., & Dome, D. (2006). Evaluation of apparent and absolute supraspinatus strength in patients with shoulder injury using the scapular retraction test. *The American Journal of Sports Medicine*, 34(10), 1643–1647. <http://doi.org/10.1177/0363546506288728>
- Kibler, W. B., Uhl, T. L., Maddux, J. W. Q., Brooks, P. V., Zeller, B., & McMullen, J. (2002b). Qualitative clinical evaluation of scapular dysfunction: a reliability study. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 11(6), 550–556. <http://doi.org/10.1067/mse.2002.126766>
- Kim, S H, Ha, K. I., & Han, K. Y. (1999). Biceps load test: a clinical test for superior labrum anterior and posterior lesions in shoulders with recurrent anterior dislocations. *The American Journal of Sports Medicine*, 27(3), 300–303.
- Kim, Seung-Ho, Ha, K.-I., Ahn, J.-H., Kim, S.-H., & Choi, H.-J. (2001). Biceps load test II: A clinical test for SLAP lesions of the shoulder. *Arthroscopy: the Journal of Arthroscopic & Related Surgery*, 17(2), 160– 164. <http://doi.org/10.1053/jars.2001.20665>
- Kim, Y S, Kim, J. M., Ha, K. Y., Choy, S., Joo, M. W., & Chung, Y. G. (2007). The Passive Compression Test: A New Clinical Test for Superior Labral Tears of the Shoulder. *The American Journal of Sports Medicine*, 35(9), 1489–1494. <http://doi.org/10.1177/0363546507301884>
- King, J. J., & Wright, T. W. (2014). Physical examination of the shoulder. *The Journal of Hand Surgery*, 39(10), 2103–2112. <http://doi.org/10.1016/j.jhsa.2014.04.024>
- Kopkow, C., Lange, T., Schmitt, J., & Kasten, P. (2015). Interrater reliability of the modified scapular assistance test with and without handheld weights. *Manual Therapy*, 20(6), 868–874. <http://doi.org/10.1016/j.math.2015.04.012>
- Kryger, A. I., Lassen, C. F., & Andersen, J. H. (2007). The role of physical examinations in studies of musculoskeletal disorders of the elbow. *Occupational and Environmental Medicine*, 64(11), 776–781. <http://doi.org/10.1136/oem.2005.026260>

- Kuhn, J. E., Dunn, W. R., Ma, B., Wright, R. W., Jones, G., Spencer, E. E., et al. (2007). Interobserver agreement in the classification of rotator cuff tears. *The American Journal of Sports Medicine*, 35(3), 437–441. <http://doi.org/10.1177/0363546506298108>
- Kuhn, J. E., Helmer, T. T., Dunn, W. R., & Throckmorton, T. W., V. (2011). Development and reliability testing of the frequency, etiology, direction, and severity (FEDS) system for classifying glenohumeral instability. *Journal of Shoulder and Elbow Surgery*, 20(4), 548–556. <http://doi.org/10.1016/j.jse.2010.10.027>
- Lange, T., Struyf, F., Schmitt, J., Lützner, J., & Kopkow, C. (2016). The reliability of physical examination tests for the clinical assessment of scapular dyskinesis in subjects with shoulder complaints: a systematic review. *Physical Therapy in Sport*, 1–64. <http://doi.org/10.1016/j.ptsp.2016.10.006>
- Leroux, J. L., Thomas, E., Bonnel, F., & Blotman, F. (1995). Diagnostic value of clinical tests for shoulder impingement syndrome. *Revue Du Rhumatisme (English Ed.)*, 62(6), 423–428.
- Levy, A. S., Lintner, S., Kenter, K., & Speer, K. P. (1999). Intra- and interobserver reproducibility of the shoulder laxity examination. *The American Journal of Sports Medicine*, 27(4), 460–463.
- Lewis, J. S., & Valentine, R. E. (2007). The pectoralis minor length test: a study of the intra-rater reliability and diagnostic accuracy in subjects with and without shoulder symptoms. *BMC Musculoskeletal Disorders*, 8, 64. <http://doi.org/10.1186/1471-2474-8-64>
- Lewis, J. S., & Valentine, R. E. (2008). Intraobserver Reliability of Angular and Linear Measurements of Scapular Position in Subjects With and Without Symptoms. *Archives of Physical Medicine and Rehabilitation*, 89(9), 1795–1802. <http://doi.org/10.1016/j.apmr.2008.01.028>
- LIN, J., & YANG, J. (2006). Reliability and validity of shoulder tightness measurement in patients with stiff shoulders. *Manual Therapy*, 11(2), 146–152. <http://doi.org/10.1016/j.math.2005.05.002>
- Lindgren, K. A., & Leino, E. (1988). Subluxation of the first rib: a possible thoracic outlet syndrome mechanism. *Archives of Physical Medicine and Rehabilitation*, 69(9), 692–695.
- Lindinger, A., Hoffmann, W., Röder, B., & Stapenhorst, K. (1984). [Compression syndrome of the upper thoracic outlet (thoracic outlet syndrome)–report on a 13-year-old girl]. *Klinische Pädiatrie*, 196(5), 315–318. <http://doi.org/10.1055/s-2008-1034087>
- Livain, T., Pichon, H., Vermeulen, J., Vaillant, J., Saragaglia, D., Poisson, M.-F., & Monnet, S. (2007). [Intra- and interobserver reproducibility of the French version of the Constant-Murley shoulder assessment during rehabilitation after rotator cuff surgery]. *Revue De Chirurgie Orthopédique Et Réparatrice De L'appareil Moteur*, 93(2), 142–149.

- Lluch, E., Benítez, J., Dueñas, L., Casaña, J., Alakhdar, Y., Nijs, J., & Struyf, F. (2014). The shoulder medial rotation test: an intertester and intratester reliability study in overhead athletes with chronic shoulder pain. *Journal of Manipulative and Physiological Therapeutics*, 37(3), 198–205. <http://doi.org/10.1016/j.jmpt.2013.12.012>
- Lo, I. K. Y., Nonweiler, B., Woolfrey, M., Litchfield, R., & Kirkley, A. (2004). An evaluation of the apprehension, relocation, and surprise tests for anterior shoulder instability. *The American Journal of Sports Medicine*, 32(2), 301–307.
- Longo, U. G., Berton, A., Ahrens, P. M., Maffulli, N., & Denaro, V. (2011). Clinical tests for the diagnosis of rotator cuff disease. *Sports Medicine and Arthroscopy Review*, 19(3), 266–278. <http://doi.org/10.1097/JSA.0b013e3182250c8b>
- Lyons, A. R., & Tomlinson, J. E. (1992). Clinical diagnosis of tears of the rotator cuff. *The Journal of Bone and Joint Surgery. British Volume*, 74(3), 414–415.
- Macdermid, J. C., Ghobrial, M., Quirion, K. B., St-Amour, M., Tsui, T., Humphreys, D., et al. (2007). Validation of a new test that assesses functional performance of the upper extremity and neck (FIT- HaNSA) in patients with shoulder pathology. *BMC Musculoskeletal Disorders*, 8, 42. <http://doi.org/10.1186/1471-2474-8-42>
- Madsen, P. H., Bak, K., Jensen, S., & Welter, U. (2011). Training Induces Scapular Dyskinesia in Pain- Free Competitive Swimmers: A Reliability and Observational Study. *Clinical Journal of Sport Medicine*, 21(2), 109–113. <http://doi.org/10.1097/JSM.0b013e3182041de0>
- Maier, M., Maier-Bosse, T., Schulz, C. U., Steinborn, M., Schmitt, C. A., Kleen, M., & Stäbler, A. (2003). Inter and intraobserver variability in DePalma's classification of shoulder calcific tendinitis. *The Journal of Rheumatology*, 30(5), 1029–1031.
- May, S., Chance-Larsen, K., Littlewood, C., Lomas, D., & Saad, M. (2010a). Reliability of physical examination tests used in the assessment of patients with shoulder problems: a systematic review. *Physiotherapy*, 96(3), 179–190. <http://doi.org/10.1016/j.physio.2009.12.002>
- McClure, P., Tate, A. R., Kareha, S., Irwin, D., & Zlupko, E. (2009). A clinical method for identifying scapular dyskinesis, part 1: reliability. *Journal of Athletic Training*, 44(2), 160–164. <http://doi.org/10.4085/1062-6050-44.2.160>
- McFarland, E. G., Garzon-Muvdi, J., Jia, X., Desai, P., & Petersen, S. A. (2010). Clinical and diagnostic tests for shoulder disorders: a critical review. *British Journal of Sports Medicine*, 44(5), 328–332. <http://doi.org/10.1136/bjsm.2009.067314>
- McFarland, E. G., Kim, T. K., & Savino, R. M. (2002). Clinical assessment of three common tests for superior labral anterior-posterior lesions. *The American Journal of Sports Medicine*, 30(6), 810–815.

- McFarland, E. G., Tanaka, M. J., Garzon-Muvdi, J., Jia, X., & Petersen, S. A. (2009). Clinical and imaging assessment for superior labrum anterior and posterior lesions. *Current Sports Medicine Reports*, 8(5), 234–239. <http://doi.org/10.1249/JSR.0b013e3181b7f042>
- McKenna, L., Straker, L., & Smith, A. (2009). The validity and intra-tester reliability of a clinical measure of humeral head position. *Manual Therapy*, 14(4), 397–403. <http://doi.org/10.1016/j.math.2008.06.004>
- McPhail, S. M., Dalland, L. N., Naess, K., Lund, M., Ytre-Hauge, T. J., Holm, S. R., & O'Leary, S. (2012). Intratherapist Reliability in the Rating of Scapula Posture in Multiple Planes of Reference. *ISRN Rehabilitation*, 2012(285), 1–8. <http://doi.org/10.1186/1477-7525-7-48>
- Mehlum, I. S., Veiersted, K. B., Waersted, M., Wergeland, E., & Kjuus, H. (2009). Self-reported versus expert-assessed work-relatedness of pain in the neck, shoulder, and arm. *Scandinavian Journal of Work, Environment & Health*, 35(3), 222–232.
- Michener, L. A., Walsworth, M. K., Doukas, W. C., & Murphy, K. P. (2009). Reliability and Diagnostic Accuracy of 5 Physical Examination Tests and Combination of Tests for Subacromial Impingement. *Archives of Physical Medicine and Rehabilitation*, 90(11), 1898–1903. <http://doi.org/10.1016/j.apmr.2009.05.015>
- Miller, C. A., Forrester, G. A., & Lewis, J. S. (2008). The Validity of the Lag Signs in Diagnosing Full- Thickness Tears of the Rotator Cuff: A Preliminary Investigation. *Archives of Physical Medicine and Rehabilitation*, 89(6), 1162–1168. <http://doi.org/10.1016/j.apmr.2007.10.046>
- Mimori, K., Muneta, T., Nakagawa, T., & Shinomiya, K. (1999). A new pain provocation test for superior labral tears of the shoulder. *The American Journal of Sports Medicine*, 27(2), 137–142.
- Munro, W., & Healy, R. (2009). The validity and accuracy of clinical tests used to detect labral pathology of the shoulder – A systematic review. *Manual Therapy*, 14(2), 119–130. <http://doi.org/10.1016/j.math.2008.08.008>
- Murrell, G. A., & Walton, J. R. (2001). Diagnosis of rotator cuff tears. *Lancet*, 357(9258), 769–770. [http://doi.org/10.1016/S0140-6736\(00\)04161-1](http://doi.org/10.1016/S0140-6736(00)04161-1)
- Myers, J. B., Oyama, S., Wassinger, C. A., Ricci, R. D., Abt, J. P., Conley, K. M., & Lephart, S. M. (2007). Reliability, precision, accuracy, and validity of posterior shoulder tightness assessment in overhead athletes. *The American Journal of Sports Medicine*, 35(11), 1922–1930. <http://doi.org/10.1177/0363546507304142>
- Nijs, J., Roussel, N., Vermeulen, K., & Souvereinys, G. (2005). Scapular positioning in patients with shoulder pain: a study examining the reliability and clinical importance of 3 clinical tests. *Archives of Physical Medicine and Rehabilitation*, 86(7), 1349–1355.

- Nomden, J. G., Slagers, A. J., Bergman, G. J. D., Winters, J. C., Kropmans, T. J. B., & Dijkstra, P. U. (2009). Interobserver reliability of physical examination of shoulder girdle. *Manual Therapy*, 14(2), 152–159. <http://doi.org/10.1016/j.math.2008.01.005>
- Nowak, D. D., Gardner, T. R., Bigliani, L. U., Levine, W. N., & Ahmad, C. S. (2010). Interobserver and intraobserver reliability of the Walch classification in primary glenohumeral arthritis. *Journal of Shoulder and Elbow Surgery*, 19(2), 180–183. <http://doi.org/10.1016/j.jse.2009.08.003>
- Nørregaard, J., Krogsgaard, M. R., Lorenzen, T., & Jensen, E. M. (2002). Diagnosing patients with longstanding shoulder joint pain. *Annals of the Rheumatic Diseases*, 61(7), 646–649.
- O'Brien, S. J., Pagnani, M. J., Fealy, S., McGlynn, S. R., & Wilson, J. B. (1998). The active compression test: a new and effective test for diagnosing labral tears and acromioclavicular joint abnormality. *The American Journal of Sports Medicine*, 26(5), 610–613.
- O'Leary, S., Christensen, S. W., Verouhis, A., Pape, M., Nilsen, O., & McPhail, S. M. (2015). Agreement between physiotherapists rating scapular posture in multiple planes in patients with neck pain: Reliability study. *Physiotherapy*, 101(4), 381–388. <http://doi.org/10.1016/j.physio.2015.01.005>
- Odom, C. J., Taylor, A. B., Hurd, C. E., & Denegar, C. R. (2001). Measurement of scapular asymmetry and assessment of shoulder dysfunction using the Lateral Scapular Slide Test: a reliability and validity study. *Physical Therapy*, 81(2), 799–809.
- Ostor, A. J. K., Richards, C. A., Prevost, A. T., Hazleman, B. L., & Speed, C. A. (2004). Interrater reproducibility of clinical tests for rotator cuff lesions. *Annals of the Rheumatic Diseases*, 63(10), 1288–1292. <http://doi.org/10.1136/ard.2003.014712>
- Palmer, K. T., Palmer, K., Walker-Bone, K., Linaker, C., Reading, I., Kellingray, S., et al. (2007). The Southampton Examination Schedule for the diagnosis of musculoskeletal disorders of the upper limb. *La Medicina Del Lavoro*, 98(2), 111–117.
- Park, H. B. (2005). Diagnostic Accuracy of Clinical Tests for the Different Degrees of Subacromial Impingement Syndrome. *The Journal of Bone and Joint Surgery*, 87(7), 1446–1455. <http://doi.org/10.2106/JBJS.D.02335>
- Park, H. B., Yokota, A., Gill, H. S., Rassi, El, G., & McFarland, E. G. (2005). Diagnostic accuracy of clinical tests for the different degrees of subacromial impingement syndrome. *The Journal of Bone and Joint Surgery. American Volume*, 87(7), 1446–1455. <http://doi.org/10.2106/JBJS.D.02335>
- Patel, S., Hossain, F. S., Colaco, H. B., El-Husseiny, M., & Lee, M. H. (2010). The accuracy of primary care teams in diagnosing disorders of the shoulder. *Journal of Evaluation in Clinical Practice*, 17(1), 118–122. <http://doi.org/10.1111/j.1365-2753.2010.01377.x>

- Pellecchia, G. L., Paolino, J., & Connell, J. (1996). Intertester reliability of the cyriax evaluation in assessing patients with shoulder pain. *The Journal of Orthopaedic and Sports Physical Therapy*, 23(1), 34–38.
- Pollard, H., Lakay, B., Tucker, F., Watson, B., & Bablis, P. (2005). Interexaminer reliability of the deltoid and psoas muscle test. *Journal of Manipulative and Physiological Therapeutics*, 28(1), 52–56. <http://doi.org/10.1016/j.jmpt.2004.12.008>
- Proud, E. L., Miller, K. J., Bilney, B., Balachandran, S., McGinley, J. L., & Morris, M. E. (2015). Evaluation of measures of upper limb functioning and disability in people with Parkinson disease: a systematic review. *Archives of Physical Medicine and Rehabilitation*, 96(3), 540–551.e1. <http://doi.org/10.1016/j.apmr.2014.09.016>
- Rabin, A., Irrgang, J. J., Fitzgerald, G. K., & Eubanks, A. (2006). The intertester reliability of the Scapular Assistance Test. *The Journal of Orthopaedic and Sports Physical Therapy*, 36(9), 653–660.
- Razmjou, H., Bean, A., Macdermid, J. C., van Osnabrugge, V., Travers, N., & Holtby, R. (2008). Convergent Validity of the Constant-Murley Outcome Measure in Patients with Rotator Cuff Disease. *Physiotherapy Canada. Physiothérapie Canada*, 60(1), 72–79. <http://doi.org/10.3138/physio/60/1/72>
- Rocourt, M. H. H., Radlinger, L., Kalberer, F., Sanavi, S., Schmid, N. S., Leunig, M., & Hertel, R. (2008). Evaluation of intratester and intertester reliability of the Constant-Murley shoulder assessment. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 17(2), 364– 369. <http://doi.org/10.1016/j.jse.2007.06.024>
- Ropars, M., Fournier, A., Campillo, B., Bonan, I., Delamarche, P., Crétual, A., & Thomazeau, H. (2010). Clinical assessment of external rotation for the diagnosis of anterior shoulder hyperlaxity. *Orthopaedics & Traumatology: Surgery & Research*, 96(8), S84–S87. <http://doi.org/10.1016/j.otsr.2010.09.002>
- Rutten, M. J. C. M., Jager, G. J., & Kiemeny, L. A. L. M. (2010). Ultrasound Detection of Rotator Cuff Tears: Observer Agreement Related to Increasing Experience. *AJR. American Journal of Roentgenology*, 195(6), W440–W446. <http://doi.org/10.2214/AJR.10.4526>
- Shadmehr, A., Bagheri, H., Ansari, N. N., & Sarafraz, H. (2010). The reliability measurements of lateral scapular slide test at three different degrees of shoulder joint abduction. *British Journal of Sports Medicine*, 44(4), 289–293. <http://doi.org/10.1136/bjsm.2008.050872>
- Sharma, S. P., Bærheim, A., & Kvale, A. (2015). Passive range of motion in patients with adhesive shoulder capsulitis, an intertester reliability study over eight weeks. *BMC Musculoskeletal Disorders*, 16, 37. <http://doi.org/10.1186/s12891-015-0495-4>
- Shin, S. H., Ro, D. H., Lee, O.-S., Oh, J. H., & Kim, S. H. (2012). Within-day reliability of shoulder range of motion measurement with a smartphone. *Manual Therapy*, 17(4), 298–304. <http://doi.org/10.1016/j.math.2012.02.010>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/musculoskeletal medicine  
Shoulder Region

- Smith, C. K., Bonauto, D. K., Silverstein, B. A., & Wilcox, D. (2010). Inter-Rater Reliability of Physical Examinations in a Prospective Study of Upper Extremity Musculoskeletal Disorders. *Journal of Occupational and Environmental Medicine*, 52(10), 1014–1018. <http://doi.org/10.1097/JOM.0b013e3181f4396b>
- Snels, I. A., Beckerman, H., Kate, ten, J. J., Lankhorst, G. J., & Bouter, L. M. (2001). Measuring subluxation of the hemiplegic shoulder: reliability of a method. *Neurorehabilitation and Neural Repair*, 15(3), 249–254.
- Stetson, W. B., & Templin, K. (2002). The crank test, the O'Brien test, and routine magnetic resonance imaging scans in the diagnosis of labral tears. *The American Journal of Sports Medicine*, 30(6), 806–809.
- Stovall, B. A., & Kumar, S. (2010). Reliability of bony anatomic landmark asymmetry assessment in the lumbopelvic region: application to osteopathic medical education. *The Journal of the American Osteopathic Association*, 110(11), 667–674. <http://doi.org/10.1016/j.apmr.2010.12.042>
- Takala, E. P. (1990). Pressure pain threshold on upper trapezius and levator scapulae muscles. Repeatability and relation to subjective symptoms in a working population., 22(2), 63–68.
- Tate, A. R., McClure, P., Kareha, S., Irwin, D., & Barbe, M. F. (2009). A Clinical Method for Identifying Scapular Dyskinesis, Part 2: Validity. *Journal of Athletic Training*, 44(2), 165–173. <http://doi.org/10.4085/1062-6050-44.2.165>
- Terwee, C. B., de Winter, A. F., Scholten, R. J., Jans, M. P., Devillé, W., van Schaardenburg, D., & Bouter, L. M. (2005). Interobserver reproducibility of the visual estimation of range of motion of the shoulder. *Archives of Physical Medicine and Rehabilitation*, 86(7), 1356–1361.
- Throckmorton, T. W., Dunn, W., Holmes, T., & Kuhn, J. E. (2009). Intraobserver and interobserver agreement of International Classification of Diseases, Ninth Revision codes in classifying shoulder instability. *Journal of Shoulder and Elbow Surgery*, 18(2), 199–203. <http://doi.org/10.1016/j.jse.2008.10.005>
- Tucci, H. T., Martins, J., Sposito, G. de C., Camarini, P. M. F., & de Oliveira, A. S. (2014). Closed Kinetic Chain Upper Extremity Stability test (CKCUES test): a reliability study in persons with and without shoulder impingement syndrome. *BMC Musculoskeletal Disorders*, 15, 1. <http://doi.org/10.1186/1471-2474-15-1>
- Tucker, S., Taylor, N. F., & Green, R. A. (2011). Anatomical validity of the Hawkins-Kennedy test - A pilot study. *Manual Therapy*, 16(4), 399–402. <http://doi.org/10.1016/j.math.2011.02.002>
- Tzannes, A., & Murrell, G. A. C. (2002). Clinical examination of the unstable shoulder. *Sports Medicine (Auckland, N.Z.)*, 32(7), 447–457.

- Tzannes, A., Paxinos, A., Callanan, M., & Murrell, G. A. C. (2004). An assessment of the interexaminer reliability of tests for shoulder instability. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 13(1), 18–23. <http://doi.org/10.1016/S1058274603002453>
- Uhl, T. L., Kibler, W. B., Gecewich, B., & Tripp, B. L. (2009). Evaluation of clinical assessment methods for scapular dyskinesis. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 25(11), 1240–1248. <http://doi.org/10.1016/j.arthro.2009.06.007>
- Valentine, R. E., & Lewis, J. S. (2006). Intraobserver reliability of 4 physiologic movements of the shoulder in subjects with and without symptoms. *Archives of Physical Medicine and Rehabilitation*, 87(9), 1242–1249. <http://doi.org/10.1016/j.apmr.2006.05.008>
- van Den Ende, C. H., Rozing, P. M., Dijkmans, B. A., Verhoef, J. A., Voogt-van der Harst, E. M., & Hazes, J. M. (1996). Assessment of shoulder function in rheumatoid arthritis. *The Journal of Rheumatology*, 23(12), 2043–2048.
- van der Sluijs, J. A., van Doorn-Loogman, M. H., Ritt, M. J. P. F., & Wuisman, P. I. J. M. (2006). Interobserver reliability of the Mallet score. *Journal of Pediatric Orthopaedics. Part B / European Paediatric Orthopaedic Society, Pediatric Orthopaedic Society of North America*, 15(5), 324–327.
- Vind, M., Bogh, S. B., Larsen, C. M., Knudsen, H. K., Sogaard, K., & Juul-Kristensen, B. (2011). Inter-examiner reproducibility of clinical tests and criteria used to identify subacromial impingement syndrome. *BMJ Open*, 1(1), e000042. <http://doi.org/10.1136/bmjopen-2010-000042>
- Vivekananda-Schmidt, P., Lewis, M., Coady, D., Morley, C., Kay, L., Walker, D., & Hassell, A. B. (2007). Exploring the use of videotaped objective structured clinical examination in the assessment of joint examination skills of medical students. *Arthritis and Rheumatism*, 57(5), 869–876. <http://doi.org/10.1002/art.22763>
- Walch, G., Boulahia, A., Calderone, S., & Robinson, A. H. (1998). The 'dropping' and 'hornblower's' signs in evaluation of rotator-cuff tears. *The Journal of Bone and Joint Surgery. British Volume*, 80(4), 624–628.
- Walsworth, M. K., Doukas, W. C., Murphy, K. P., Mielcarek, B. J., & Michener, L. A. (2007). Reliability and Diagnostic Accuracy of History and Physical Examination for Diagnosing Glenoid Labral Tears. *The American Journal of Sports Medicine*, 36(1), 162–168. <http://doi.org/10.1177/0363546507307508>
- Warner, J. J., & McMahon, P. J. (1995). The role of the long head of the biceps brachii in superior stability of the glenohumeral joint. *The Journal of Bone and Joint Surgery. American Volume*, 77(3), 366–372.

- Wolf, E. M., & Agrawal, V. (2001). Transdeltoid palpation (the rent test) in the diagnosis of rotator cuff tears. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 10(5), 470–473. <http://doi.org/10.1067/mse.2001.117126>
- Yang, J. S., Keener, J. D., Yamaguchi, K., Chen, J., Stobbs-Cucchi, G., Patton, R., & Galatz, L. M. (2015). Reliability of patient self-assessment of shoulder range of motion and strength after shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 24(1), 1–10. <http://doi.org/10.1016/j.jse.2014.08.025>
- Yang, J.-L., & Lin, J.-J. (2006). Reliability of function-related tests in patients with shoulder pathologies. *The Journal of Orthopaedic and Sports Physical Therapy*, 36(8), 572–576.
- Yonenobu, K., Abumi, K., Nagata, K., Taketomi, E., & Ueyama, K. (2001). Interobserver and intraobserver reliability of the japanese orthopaedic association scoring system for evaluation of cervical compression myelopathy. *Spine*, 26(17), 1890–4; discussion 1895.
- Zaslav, K. R. (2001). Internal rotation resistance strength test: a new diagnostic test to differentiate intra-articular pathology from outlet (Neer) impingement syndrome in the shoulder. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 10(1), 23–27. <http://doi.org/10.1067/mse.2001.111960>
- Zuckerman, J. D., Kummer, F. J., Cuomo, F., & Greller, M. (1997). Interobserver reliability of acromial morphology classification: an anatomic study. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 6(3), 286–287.

## Thoracic Region

- Aartun, E., Aartun, E., Degerfalk, A., Degerfalk, A., Kentsdotter, L., Kentsdotter, L., & Hestbaek, L. (2014). Screening of the spine in adolescents: inter- and intra-rater reliability and measurement error of commonly used clinical tests. *BMC Musculoskeletal Disorders*, 15(1), 1–10. <http://doi.org/10.1186/1471-2474-15-37>
- Anderson, V. B. (2011). The intra-rater reliability of measured thoracic spine mobility in chronic rotator cuff pathology. *Journal of Musculoskeletal & Neuronal Interactions*, 11(4), 314–319.
- Bertilson, B. C., Grunnesjö, M., & Strender, L.-E. (2003). Reliability of Clinical Tests in the Assessment of Patients With Neck/Shoulder Problems—Impact of History. *Spine*, 28(19), 2222–2231. <http://doi.org/10.1097/01.BRS.0000089685.55629.2E>
- Bock, S. L., Centeno, C. J., & Elliott, J. M. (2005). The presence and interrater reliability of thoracic allodynia in a whiplash cohort. *Pain Physician*, 8(3), 267–270.
- Boezaart, A. P., Haller, A., Laduzenski, S., Koyyalamudi, V. B., Ihnatsenka, B., & Wright, T. (2010). Neurogenic thoracic outlet syndrome: A case report and review of the literature. *International Journal of Shoulder Surgery*, 4(2), 27–35. <http://doi.org/10.4103/0973-6042.70817>
- Brismée, J.-M., Gipson, D., Ivie, D., Lopez, A., Moore, M., Matthijs, O., et al. (2006). Interrater reliability of a passive physiological intervertebral motion test in the mid-thoracic spine. *Journal of Manipulative and Physiological Therapeutics*, 29(5), 368–373. <http://doi.org/10.1016/j.jmpt.2006.04.009>
- Brunse, M. H., Stochkendahl, M. J., Vach, W., Kongsted, A., Poulsen, E., Hartvigsen, J., & Christensen,
- H. W. (2010a). Examination of musculoskeletal chest pain - an inter-observer reliability study. *Manual Therapy*, 15(2), 167–172. <http://doi.org/10.1016/j.math.2009.10.003>
- Brunse, M. H., Stochkendahl, M. J., Vach, W., Kongsted, A., Poulsen, E., Hartvigsen, J., & Christensen,
- H. W. (2010b). Examination of musculoskeletal chest pain - an inter-observer reliability study. *Manual Therapy*, 15(2), 167–172. <http://doi.org/10.1016/j.math.2009.10.003>
- Christensen, H. W., Vach, W., Manniche, C., Haghfelt, T., Hartvigsen, L., & Høilund-Carlsen, P. F. (2003). Palpation for muscular tenderness in the anterior chest wall: an observer reliability study. *Journal of Manipulative and Physiological Therapeutics*, 26(8), 469–475. [http://doi.org/10.1016/S0161-4754\(03\)00103-9](http://doi.org/10.1016/S0161-4754(03)00103-9)

- Christensen, H. W., Vach, W., Vach, K., Manniche, C., Haghfelt, T., Hartvigsen, L., & Flemming Højilund-Carlsen, P. (2002a). Palpation of the upper thoracic spine: An observer reliability study. *Journal of Manipulative and Physiological Therapeutics*, 25(5), 285–292. <http://doi.org/10.1067/mmt.2002.124424>
- Cooperstein, R. (2012). Interexaminer reliability of the Johnston and Friedman percussion scan of the thoracic spine: secondary data analysis using modified methods. *Journal of Chiropractic Medicine*, 11(3), 154–159. <http://doi.org/10.1016/j.jcm.2012.06.001>
- Cooperstein, R., DC, R. C. M., DC, M. H. M., Haneline, M., Young, M., & DC, M. Y. (2010). Interexaminer reliability of thoracic motion palpation using confidence ratings and continuous analysis. *Journal of Chiropractic Medicine*, 9(3), 99–106. <http://doi.org/10.1016/j.jcm.2010.06.004>
- de Sèze, M., Randriamnahisoa, T., Gaunelle, A., de Korvin, G., & Mazaux, J. M. (2013). Inter-observer reproducibility of back surface topography parameters allowing assessment of scoliotic thoracic gibbosity and comparison with two standard postures. *Annals of Physical and Rehabilitation Medicine*, 56(9-10), 599–612. <http://doi.org/10.1016/j.rehab.2013.10.003>
- Elgueta-Cancino, E., Schabrun, S., Danneels, L., & Hodges, P. (2014). A clinical test of lumbopelvic control: development and reliability of a clinical test of dissociation of lumbopelvic and thoracolumbar motion. *Manual Therapy*, 19(5), 418–424. <http://doi.org/10.1016/j.math.2014.03.009>
- Heiderscheit, B., & Boissonnault, W. (2008). Reliability of joint mobility and pain assessment of the thoracic spine and rib cage in asymptomatic individuals. *The Journal of Manual & Manipulative Therapy*, 16(4), 210–216.
- Johnson, K. D., Kim, K.-M., Yu, B.-K., Saliba, S. A., & Grindstaff, T. L. (2012). Reliability of thoracic spine rotation range-of-motion measurements in healthy adults. *Journal of Athletic Training*, 47(1), 52–60.
- Lamartina, C., & Berjano, P. (2014). Classification of sagittal imbalance based on spinal alignment and compensatory mechanisms. *European Spine Journal*, 23(6), 1177–1189. <http://doi.org/10.1007/s00586-014-3227-9>
- Lenke, L. G., Betz, R. R., Bridwell, K. H., Clements, D. H., Harms, J., Lowe, T. G., & Shufflebarger, H. L. (1998). Intraobserver and interobserver reliability of the classification of thoracic adolescent idiopathic scoliosis. *The Journal of Bone and Joint Surgery. American Volume*, 80(8), 1097–1106.
- Oppenheim, U., Kohen-Raz, R., Alex, D., Kohen-Raz, A., & Azarya, M. (1999). Postural characteristics of diabetic neuropathy. *Diabetes Care*, 22(2), 328–332.
- Paulet, T., & Fryer, G. (2009). International Journal of Osteopathic Medicine. *International Journal of Osteopathic Medicine*, 12(3), 92–96. <http://doi.org/10.1016/j.ijosm.2008.07.001>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
**Thoracic Region**

- Potter, L., McCarthy, C., & Oldham, J. (2006). Intraexaminer reliability of identifying a dysfunctional segment in the thoracic and lumbar spine. *Journal of Manipulative and Physiological Therapeutics*, 29(3), 203–207. <http://doi.org/10.1016/j.jmpt.2006.01.005>
- Redelmeier, D. A., Guyatt, G. H., & Goldstein, R. S. (1996). Assessing the minimal important difference in symptoms: a comparison of two techniques. *Journal of Clinical Epidemiology*, 49(11), 1215–1219.
- Shojaei, I., Arjmand, N., & Bazrgari, B. (2015). An optimization-based method for prediction of lumbar spine segmental kinematics from the measurements of thorax and pelvic kinematics. *International Journal for Numerical Methods in Biomedical Engineering*, 31(12). <http://doi.org/10.1002/cnm.2729>
- Stovall, B. A., & Kumar, S. (2010). Reliability of bony anatomic landmark asymmetry assessment in the lumbopelvic region: application to osteopathic medical education. *The Journal of the American Osteopathic Association*, 110(11), 667–674. <http://doi.org/10.1016/j.apmr.2010.12.042>
- van Middendorp, J. J., Patel, A. A., Schuetz, M., & Joaquim, A. F. (2013). The precision, accuracy and validity of detecting posterior ligamentous complex injuries of the thoracic and lumbar spine: a critical appraisal of the literature. *European Spine Journal*, 22(3), 461–474. <http://doi.org/10.1007/s00586-012-2602-7>
- Walker, B. F., Koppenhaver, S. L., Stomski, N. J., & Hebert, J. J. (2015). Interrater Reliability of Motion Palpation in the Thoracic Spine. *Evidence-Based Complementary and Alternative Medicine : eCAM*, 2015, 815407. <http://doi.org/10.1155/2015/815407>

## Upper Extremity Region

- Aartun, E., Aartun, E., Degerfalk, A., Degerfalk, A., Kentsdotter, L., Kentsdotter, L., & Hestbaek, L. (2014). Screening of the spine in adolescents: inter- and intra-rater reliability and measurement error of commonly used clinical tests. *BMC Musculoskeletal Disorders*, 15(1), 1–10. <http://doi.org/10.1186/1471-2474-15-37>
- Abouelela, A. A. K. H., & Zohiery, A. K. (2012). The triple compression stress test for diagnosis of tarsal tunnel syndrome. *Foot (Edinburgh, Scotland)*, 22(3), 146–149. <http://doi.org/10.1016/j.foot.2012.02.002>
- Ageberg, E., Bennell, K. L., Hunt, M. A., Simic, M., Roos, E. M., & Creaby, M. W. (2010). Validity and inter-rater reliability of medio-lateralknee motion observed during a single-limb minisquat. *BMC Musculoskeletal Disorders*, 11(1), 265. <http://doi.org/10.1186/1471-2474-11-265>
- AKGUL, O., Guldeste, Z., & Ozgocmen, S. (2014). The reliability of the clinical examination for detecting Baker's cyst in asymptomatic fossa. *International Journal of Rheumatic Diseases*, 17(2), 204–209. <http://doi.org/10.1111/1756-185X.12095>
- Alexiou, I., Germenis, A., Koutroumpas, A., Kontogianni, A., Theodoridou, K., & Sakkas, L. I. (2008). Anti-cyclic citrullinated peptide-2 (CCP2) autoantibodies and extra-articular manifestations in Greek patients with rheumatoid arthritis. *Clinical Rheumatology*, 27(4), 511–513. <http://doi.org/10.1007/s10067-007-0800-1>
- Allington, N. J., Leroy, N., & Doneux, C. (2002). Ankle joint range of motion measurements in spastic cerebral palsy children: intraobserver and interobserver reliability and reproducibility of goniometry and visual estimation. *Journal of Pediatric Orthopaedics. Part B / European Paediatric Orthopaedic Society, Pediatric Orthopaedic Society of North America*, 11(3), 236–239.
- Almoallim, H., Attar, S., Jannoudi, N., Al-Nakshabandi, N., Eldeek, B., Fathaddien, O., & Halabi, H. (2012). Sensitivity of standardised musculoskeletal examination of the hand and wrist joints in detecting arthritis in comparison to ultrasound findings in patients attending rheumatology clinics. *Clinical Rheumatology*. <http://doi.org/10.1007/s10067-012-2013-5>
- Alonso, A., Khoury, L., & Adams, R. (1998). Clinical tests for ankle syndesmosis injury: reliability and prediction of return to function. *The Journal of Orthopaedic and Sports Physical Therapy*, 27(4), 276–284. <http://doi.org/10.2519/jospt.1998.27.4.276>
- Altan, L., Ercan, I., & Konur, S. (2010). Reliability and validity of Turkish version of the patient rated tennis elbow evaluation. *Rheumatology International*, 30(8), 1049–1054. <http://doi.org/10.1007/s00296-009-1101-6>

- Amer, M., Hubert, G., Sullivan, S. J., Herbison, P., Franz, E. A., & Hammond-Tooke, G. D. (2012). Reliability and diagnostic characteristics of clinical tests of upper limb motor function. *Journal of Clinical Neuroscience : Official Journal of the Neurosurgical Society of Australasia*, 19(9), 1246–1251. <http://doi.org/10.1016/j.jocn.2011.12.007>
- Ames, R., Jarosz, B. S., Baglin, J., & Da Costa, C. (2015). Functional Knee Extension Test (FKET): Inter- rater palpatory agreement and visual analysis during a non-weight bearing functional assessment of the knee. *Journal of Bodywork and Movement Therapies*, 19(4), 656–665. <http://doi.org/10.1016/j.jbmt.2014.12.005>
- Andersson, J. K., Andersnord, D., Karlsson, J., & Fridén, J. (2015). Efficacy of Magnetic Resonance Imaging and Clinical Tests in Diagnostics of Wrist Ligament Injuries: A Systematic Review. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 31(10), 2014–20.e2. <http://doi.org/10.1016/j.arthro.2015.04.090>
- Ansari, N. N., Naghdi, S., Mashayekhi, M., Hasson, S., Fakhari, Z., & Jalaie, S. (2012). Intra-rater reliability of the Modified Modified Ashworth Scale (MMAS) in the assessment of upper-limb muscle spasticity. *NeuroRehabilitation*, 31(2), 215–222. <http://doi.org/10.3233/NRE-2012-0791>
- Armstrong, A. D., MacDermid, J. C., Chinchalkar, S., Stevens, R. S., & King, G. J. (1998). Reliability of range-of-motion measurement in the elbow and forearm. *Journal of Shoulder and Elbow Surgery /American Shoulder and Elbow Surgeons [Et Al]*, 7(6), 573–580.
- ATC, J. M. F. P. P. C., & Fedorczyk, J. M. (2012). Tendinopathies of the Elbow, Wrist, and Hand: Histopathology and Clinical Considerations. *Journal of Hand Therapy*, 25(2), 191–201. <http://doi.org/10.1016/j.jht.2011.12.001>
- Bagley, A. M., Molitor, F., Wagner, L. V., Tomhave, W., & James, M. A. (2006). The Unilateral Below Elbow Test: a function test for children with unilateral congenital below elbow deficiency. *Developmental Medicine and Child Neurology*, 48(7), 569–575. <http://doi.org/10.1017/S0012162206001204>
- Baldwin, C. E., Hons, C. E. B. B. P., PhD, J. D. P., & MD, A. D. B. (2013). Muscle strength assessment in critically ill patients with handheld dynamometry: An investigation of reliability, minimal detectable change, and timeto peak force generation. *Journal of Critical Care*, 28(1), 77–86. <http://doi.org/10.1016/j.jcrc.2012.03.001>
- Beekman, R., Schreuder, A. H. C. M. L., Rozeman, C. A. M., Koehler, P. J., & Uitdehaag, B. M. J. (2009). The diagnostic value of provocative clinical tests in ulnar neuropathy at the elbow is marginal. *Journal of Neurology, Neurosurgery & Psychiatry*, 80(12), 1369–1374. <http://doi.org/10.1136/jnnp.2009.180844>
- Bertilson, B. C., Grunnesjö, M., & Strender, L.-E. (2003). Reliability of Clinical Tests in the Assessment of Patients With Neck/Shoulder Problems—Impact of History. *Spine*, 28(19), 2222–2231. <http://doi.org/10.1097/01.BRS.0000089685.55629.2E>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine Upper Extremity Region

- Beumer, A., Swierstra, B. A., & Mulder, P. G. H. (2002). Clinical diagnosis of syndesmotic ankle instability: evaluation of stress tests behind the curtains. *Acta Orthopaedica Scandinavica*, 73(6), 667– 669. <http://doi.org/10.1080/000164702321039642>
- Bhardwaj, P., Venkatramani, H., Sabapathy, S. R., & Sabapathy, S. R. (2011). A modified towel test for assessment of elbow flexion in children less than 9 months old with brachial plexus birth palsy. *Journal of Hand Surgery (European Volume)*, 36(8), 707–708. <http://doi.org/10.1177/1753193411419787>
- Blonna, D., MD, D. B., C, P. C. Z. M. F., BSc, J. S. F., & C, S. W. O. P. M. F. (2012). Validation of a photography-based goniometry method for measuring joint range of motion. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 21(1), 29–35. <http://doi.org/10.1016/j.jse.2011.06.018>
- Blonna, D., Zarkadas, P. C., Fitzsimmons, J. S., & O'Driscoll, S. W. (2011). Accuracy and inter-observer reliability of visual estimation compared to clinical goniometry of the elbow. *Knee Surgery, Sports Traumatology, Arthroscopy : Official Journal of the ESSKA*. <http://doi.org/10.1007/s00167-011-1720-9>
- Boeree, N. R., & Ackroyd, C. E. (1991). Assessment of the menisci and cruciate ligaments: an audit of clinical practice. *Injury*, 22(4), 291–294.
- Boyle, K. L., Witt, P., & Riegger-Krugh, C. (2003). Intrarater and Interrater Reliability of the Beighton and Horan Joint Mobility Index. *Journal of Athletic Training*, 38(4), 281–285.
- Brandsma, J. W., Van Brakel, W. H., Anderson, A. M., Kortendijk, A. J., Gurung, K. S., & Sunwar, S. K. (1998). Intertester reliability of manual muscle strength testing in leprosy patients. *Leprosy Review*, 69(3), 257–266.
- Broekstra, D. C., Lanting, R., Werker, P. M. N., & van den Heuvel, E. R. (2015). Intra- and inter-observer agreement on diagnosis of Dupuytren disease, measurements of severity of contracture, and disease extent. *Manual Therapy*, 20(4), 580–586. <http://doi.org/10.1016/j.math.2015.01.010>
- Calfee, R. P., Calfee, R. P., Manske, P. R., Manske, P. R., Gelberman, R. H., Gelberman, R. H., et al. (2010). Clinical Assessment of the Ulnar Nerve at the Elbow: Reliability of Instability Testing and the Association of Hypermobility with Clinical Symptoms. *The Journal of Bone and Joint Surgery*, 92(17), 2801–2808. <http://doi.org/10.2106/JBJS.J.00097>
- Cejudo, A., Sainz de Baranda, P., Ayala, F., & Santonja, F. (2014). A simplified version of the weight-bearing ankle lunge test: description and test-retest reliability. *Manual Therapy*, 19(4), 355–359. <http://doi.org/10.1016/j.math.2014.03.008>
- Chang, C.-W., Wang, Y.-C., & Chu, C.-H. (2008). Increased Carrying Angle is a Risk Factor for Nontraumatic Ulnar Neuropathy at the Elbow. *Clinical Orthopaedics and Related Research*, 466(9), 2190–2195. <http://doi.org/10.1007/s11999-008-0308-2>

- Chatterton, B. D., Muller, S., Thomas, M. J., Menz, H. B., Rome, K., & Roddy, E. (2013). Inter and intra- rater repeatability of the scoring of foot pain drawings. *Journal of Foot and Ankle Research*, 6(1), 1–1. <http://doi.org/10.1186/1757-1146-6-44>
- Choa, R. M., & Giele, H. P. (2015). Inter- and intrarater reliability of osteoarthritis classification at the trapeziometacarpal joint. *The Journal of Hand Surgery*, 40(1), 23–26. <http://doi.org/10.1016/j.jhsa.2014.09.007>
- Cibere, J., Bellamy, N., Thorne, A., Esdaile, J. M., McGorm, K. J., Chalmers, A., et al. (2004). Reliability of the knee examination in osteoarthritis: Effect of standardization. *Arthritis and Rheumatism*, 50(2), 458–468. <http://doi.org/10.1002/art.20025>
- Cleffken, B., van Breukelen, G., van Mameren, H., Brink, P., & Damink, S. O. (2007). Test-retest reproducibility of elbow goniometric measurements in a rigid double-blinded protocol: Intervals for distinguishing between measurement error and clinical change. *Journal of Shoulder and Elbow Surgery*, 16(6), 788–794.e2. <http://doi.org/10.1016/j.jse.2007.02.134>
- Cook, J. L., Khan, K. M., Kiss, Z. S., Purdam, C. R., & Griffiths, L. (2001). Reproducibility and clinical utility of tendon palpation to detect patellar tendinopathy in young basketball players. Victorian Institute of Sport tendon study group. *British Journal of Sports Medicine*, 35(1), 65–69.
- Cooperman, J. M., Riddle, D. L., & Rothstein, J. M. (1990). Reliability and validity of judgments of the integrity of the anterior cruciate ligament of the knee using the Lachman's test. *Physical Therapy*, 70(4), 225–233.
- Cornwall, M. W., Fishco, W. D., McPoil, T. G., Lane, C. R., O'Donnell, D., & Hunt, L. (2004). Reliability and validity of clinically assessing first-ray mobility of the foot. *Journal of the American Podiatric Medical Association*, 94(5), 470–476.
- Currier, L. L., Froehlich, P. J., Carow, S. D., McAndrew, R. K., Cliborne, A. V., Boyles, R. E., et al. (2007). Development of a clinical prediction rule to identify patients with knee pain and clinical evidence of knee osteoarthritis who demonstrate a favorable short-term response to hip mobilization. *Physical Therapy*, 87(9), 1106–1119. <http://doi.org/10.2522/ptj.20060066>
- Cushnaghan, J., Cooper, C., Dieppe, P., Kirwan, J., McAlindon, T., & McCrae, F. (1990). Clinical assessment of osteoarthritis of the knee. *Annals of the Rheumatic Diseases*, 49(10), 768–770.
- Dawson, J., Doll, H., Boller, I., FITZPATRICK, R., Little, C., Rees, J., & Carr, A. (2012). Specificity and responsiveness of patient-reported and clinician-rated outcome measures in the context of elbow surgery, comparing patients with and without rheumatoid arthritis. *Orthopaedics & Traumatology, Surgery & Research : OTSR*, 98(6), 652–658. <http://doi.org/10.1016/j.otsr.2012.05.011>

- Dennis, R. J., Finch, C. F., Elliott, B. C., & Farhart, P. J. (2008). The reliability of musculoskeletal screening tests used in cricket. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 9(1), 25–33. <http://doi.org/10.1016/j.ptsp.2007.09.004>
- Derksen, R.-J., Bakker, F. C., Geervliet, P. C., de Lange de Klerk, E. S. M., Heilbron, E. A., Veenings, B., et al. (2005). Diagnostic accuracy and reproducibility in the interpretation of Ottawa ankle and foot rules by specialized emergency nurses. *The American Journal of Emergency Medicine*, 23(6), 725–729. <http://doi.org/10.1016/j.ajem.2005.02.054>
- Derwin, G. F., Stiell, I. G., Wells, G. A., Rody, K., & Grabowski, J. (2001). Physicians' accuracy and interrater reliability for the diagnosis of unstable meniscal tears in patients having osteoarthritis of the knee. *Canadian Journal of Surgery. Journal Canadien De Chirurgie*, 44(4), 267–274.
- Digiovanni, C. W., Holt, S., Czerniecki, J. M., Ledoux, W. R., & Sangeorzan, B. J. (2001). Can the presence of equinus contracture be established by physical exam alone? *Journal of Rehabilitation Research and Development*, 38(3), 335–340.
- Does Elbow Position Affect Strength and Reproducibility of Power Grip Measurements? (2005). Does Elbow Position Affect Strength and Reproducibility of Power Grip Measurements?, 1–5.
- Eathorne, S. W. (2005). The Wrist: Clinical Anatomy and Physical Examination—an Update. *Primary Care: Clinics in Office Practice*, 32(1), 17–33. <http://doi.org/10.1016/j.pop.2004.11.009>
- Elson, D. W., Jones, S., Caplan, N., Stewart, S., St Clair Gibson, A., Gibson, A. S. C., & Kader, D. F. (2011). The photographic knee pain map: Locating knee pain with an instrument developed for diagnostic, communication and research purposes. *The Knee*, 18(6), 417–423. <http://doi.org/10.1016/j.knee.2010.08.012>
- Eren, O. T. (2003). The accuracy of joint line tenderness by physical examination in the diagnosis of meniscal tears. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 19(8), 850– 854. [http://doi.org/10.1053/S0749-8063\(03\)00736-9](http://doi.org/10.1053/S0749-8063(03)00736-9)
- Erichsen, N., Lund, H., Ole Møller, J., Kaiser, T., Lykke Jensen, M., Märcher, I., et al. (2006). Inter-rater and intra-rater reliability of tests of translatory movements and range of movements in the subtalar and talocrural joints. *Advances in Physiotherapy*, 8(4), 161–167. <http://doi.org/10.1080/14038190601000066>
- Fredericson, M., & Yoon, K. (2006). Physical Examination and Patellofemoral Pain Syndrome. *American Journal of Physical Medicine & Rehabilitation / Association of Academic Physiatrists*, 85(3), 234–243. <http://doi.org/10.1097/01.phm.0000200390.67408.f0>

- Fritz, J. M., Delitto, A., Erhard, R. E., & Roman, M. (1998). An examination of the selective tissue tension scheme, with evidence for the concept of a capsular pattern of the knee. *Physical Therapy*, 78(10), 1046–56– discussion 1057–61.
- Gatt, A., & Chockalingam, N. (2012). Assessment of Ankle Joint Dorsiflexion: An Overview. *Revista Internacional De Ciencias Podológicas*, 6(1), 25–29. [http://doi.org/10.5209/rev\\_RICP.2012.v6.n1.37894](http://doi.org/10.5209/rev_RICP.2012.v6.n1.37894)
- Gelberman, R. H., Boone, S., Osei, D. A., Cherney, S., & Calfee, R. P. (2015). Trapeziometacarpal Arthritis: A Prospective Clinical Evaluation of the Thumb Adduction and Extension Provocative Tests. *The Journal of Hand Surgery*, 40(7), 1285–1291. <http://doi.org/10.1016/j.jhsa.2015.04.012>
- Gibbons, P., Dumper, C., & Gosling, C. (2000). Inter-examiner and intra-examiner agreement for assessing simulated leg length inequality using palpation and observation during a standing assessment. *Journal of Osteopathic Medicine*, 5(2), 53–58.
- Glasoe, W. M., Getsoian, S., Myers, M., Komnick, M., Kolkebeck, D., Oswald, W., & Liakos, P. (2005). Criterion-related validity of a clinical measure of dorsal first ray mobility. *The Journal of Orthopaedic and Sports Physical Therapy*, 35(9), 589–593. <http://doi.org/10.2519/jospt.2005.35.9.589>
- Grahame-Smith, D. (1995). Evidence based medicine: Socratic dissent. *BMJ (Clinical Research Ed)*, 310(6987), 1126–1127.
- H Charles Woodfield RPh, D. C., Woodfield, H. C., Gerstman, B. B., B Burt Gerstman DVM, M. P., Olaisen, R. H., Renate Henry Olaisen DC, M., et al. (2011). Interexaminer Reliability of Supine Leg Checks for Discriminating Leg-Length Inequality. *Journal of Manipulative and Physiological Therapeutics*, 34(4), 239–246. <http://doi.org/10.1016/j.jmpt.2011.04.009>
- Harris-Hayes, M., Steger-May, K., Koh, C., Royer, N. K., Graci, V., & Salsich, G. B. (2014). Classification of lower extremity movement patterns based on visual assessment: reliability and correlation with 2- dimensional video analysis. *Journal of Athletic Training*, 49(3), 304–310. <http://doi.org/10.4085/1062-6050-49.2.21>
- Hart, D. J., Spector, T. D., Brown, P., Wilson, P., Doyle, D. V., & Silman, A. J. (1991). Clinical signs of early osteoarthritis: reproducibility and relation to x ray changes in 541 women in the general population. *Annals of the Rheumatic Diseases*, 50(7), 467–470.
- Hauzeur, J. P., Mathy, L., & De Maertelaer, V. (1999). Comparison between clinical evaluation and ultrasonography in detecting hyarthrosis of the knee. *The Journal of Rheumatology*, 26(12), 2681– 2683.
- Hayes, K. W., & Petersen, C. M. (2001). Reliability of assessing end-feel and pain and resistance sequence in subjects with painful shoulders and knees. *The Journal of Orthopaedic and Sports Physical Therapy*, 31(8), 432–445.

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
**Upper Extremity Region**

- Hayes, K. W., Petersen, C., & Falconer, J. (1994a). An examination of Cyriax's passive motion tests with patients having osteoarthritis of the knee. *Physical Therapy*, 74(8), 697–707–discussion 707–9.
- Hayes, K. W., Petersen, C., & Falconer, J. (1994b). An examination of Cyriax's passive motion tests with patients having osteoarthritis of the knee. *Physical Therapy*, 74(8), 697–707–discussion 707–9.
- Hermans, G., Clerckx, B., & Vanhullebusch, T. (2012). Interobserver agreement of medical research council sum-score and handgrip strength in the intensive care unit. *Muscle & ....*
- Hirsch, C., Hirsch, M., John, M. T., & Bock, J. J. (2007). Reliability of the Beighton Hypermobility Index to determine the general joint laxity performed by dentists. *Journal of Orofacial Orthopedics = Fortschritte Der Kieferorthopädie : Organ/Official Journal Deutsche Gesellschaft Fur Kieferorthopädie*, 68(5), 342–352.  
<http://doi.org/10.1007/s00056-007-0708-z>
- Hsieh, C. Y., Hong, C. Z., Adams, A. H., Platt, K. J., Danielson, C. D., Hoehler, F. K., & Tobis, J. S. (2000). Interexaminer reliability of the palpation of trigger points in the trunk and lower limb muscles. *Archives of Physical Medicine and Rehabilitation*, 81(3), 258–264.
- Hurkmans, E. J., Hurkmans, E. J., van den Berg, M. H., van den Berg, M. H., Ronday, K. H., Ronday, K. H., et al. (2009). Maintenance of physical activity after Internet-based physical activity interventions in patients with rheumatoid arthritis. *Rheumatology*, 49(1), 167–172.  
<http://doi.org/10.1093/rheumatology/kep285>
- Hutchison, A.-M., Evans, R., Bodger, O., Pallister, I., Topliss, C., Williams, P., et al. (2013). Foot and Ankle Surgery. *Foot and Ankle Surgery*, 19(2), 112–117.  
<http://doi.org/10.1016/j.fas.2012.12.006>
- Jarvis, H. L., Nester, C. J., Jones, R. K., Williams, A., & Bowden, P. D. (2012). Inter-assessor reliability of practice based biomechanical assessment of the foot and ankle. *Journal of Foot and Ankle Research*, 5, 14. <http://doi.org/10.1186/1757-1146-5-14>
- Jensen, L. K., Mikkelsen, S., Loft, I. P., & Eenberg, W. (2000). Work-related knee disorders in floor layers and carpenters. *Journal of Occupational and Environmental Medicine*, 42(8), 835–842.
- Jepsen, J. R., Laursen, L. H., Hagert, C.-G., Kreiner, S., & Larsen, A. I. (2006). Diagnostic accuracy of the neurological upper limb examination I: inter-rater reproducibility of selected findings and patterns. *BMC Neurology*, 6, 8. <http://doi.org/10.1186/1471-2377-6-8>
- Jepsen, J., Laursen, L., Larsen, A., & Hagert, C.-G. (2004). Manual strength testing in 14 upper limb muscles: a study of inter-rater reliability. *Acta Orthopaedica Scandinavica*, 75(4), 442–448.

- Jie, K. E., van Dam, L. F., Verhagen, T. F., & Hammacher, E. R. (2014). Extension test and ossal point tenderness cannot accurately exclude significant injury in acute elbow trauma. *Annals of Emergency Medicine*, 64(1), 74–78. <http://doi.org/10.1016/j.annemergmed.2014.01.022>
- Jones, A., Hopkinson, N., Pattrick, M., Berman, P., & Doherty, M. (1992). Evaluation of a method for clinically assessing osteoarthritis of the knee. *Annals of the Rheumatic Diseases*, 51(2), 243–245.
- Junge, T., Balsnes, S. L., Runge, L., Juul-Kristensen, B., & Wedderkopp, N. (2012). Single leg mini squat: an inter-tester reproducibility study of children in the age of 9–10 and 12–14 years presented by various methods of kappa calculation. *BMC Musculoskeletal Disorders*, 13(1), 1–1. <http://doi.org/10.1186/1471-2474-13-203>
- Karim, Z., Wakefield, R. J., Quinn, M., Conaghan, P. G., Brown, A. K., Veale, D. J., et al. (2004). Validation and reproducibility of ultrasonography in the detection of synovitis in the knee: A comparison with arthroscopy and clinical examination. *Arthritis and Rheumatism*, 50(2), 387–394. <http://doi.org/10.1002/art.20054>
- Kaya, T., Karatepe, A. G., Goksel Karatepe, A., Gunaydin, R., Koc, A., & Altundal Ercan, U. (2011). Inter- rater reliability of the Modified Ashworth Scale and modified Modified Ashworth Scale in assessing poststroke elbow flexor spasticity. *International Journal of Rehabilitation Research*, 34(1), 59–64. <http://doi.org/10.1097/MRR.0b013e32833d6cdf>
- Kelly R Holt BSc Chiro, P., Chiro, D. G. R. B., Nicholas J Hoffmann BHLthSci, B. C., Chiro, B. I. B. B., Chiro, P. M. B. B., & Heidi Haavik Taylor BSc Chiro, P. (2009). Interexaminer Reliability of a Leg Length Analysis Procedure Among Novice and Experienced Practitioners. *Journal of Manipulative and Physiological Therapeutics*, 32(3), 216–222. <http://doi.org/10.1016/j.jmpt.2009.02.009>
- Khamwong, P., Nosaka, K., Pirunsan, U., & Paungmali, A. (2010a). Reliability of muscle function and sensory perception measurements of the wrist extensors. *Physiotherapy Theory and Practice*, 26(6), 408–415. <http://doi.org/10.3109/09593980903300470>
- Kim, J. P., & Park, M. J. (2008). Assessment of Distal Radioulnar Joint Instability After Distal Radius Fracture: Comparison of Computed Tomography and Clinical Examination Results. *The Journal of Hand Surgery*, 33(9), 1486–1492. <http://doi.org/10.1016/j.jhsa.2008.05.017>
- Kiter, E., & Bozkurt, M. (2005). The crossed-leg test for examination of ankle syndesmosis injuries. *Foot & Ankle International / American Orthopaedic Foot and Ankle Society [and] Swiss Foot and Ankle Society*, 26(2), 187–188.
- Kryger, A. I., Lassen, C. F., & Andersen, J. H. (2007a). The role of physical examinations in studies of musculoskeletal disorders of the elbow. *Occupational and Environmental Medicine*, 64(11), 776–781. <http://doi.org/10.1136/oem.2005.026260>

- Kwon, B. C., Kwon, B. C., Choi, S. J., Choi, S.-J., Song, S.-Y., Song, S. Y., et al. (2011). Modified Carpal Stretch Test as a Screening Test for Detection of Scapholunate Interosseous Ligament Injuries Associated with Distal Radial Fractures. *The Journal of Bone and Joint Surgery*, 93(9), 855–862. <http://doi.org/10.2106/JBJS.J.00361>
- LaStayo, P. C., & Wheeler, D. L. (1994). Reliability of passive wrist flexion and extension goniometric measurements: a multicenter study. *Physical Therapy*, 74(2), 162–74–discussion 174–6.
- Leard, J. S., EdD, J. S. L., PhD, B. A. C., Crane, B. A., PhD, K. A. B., & Ball, K. A. (2009). Intrarater and interrater reliability of 22 clinical measures associated with lower quarter malalignment. *Journal of Manipulative and Physiological Therapeutics*, 32(4), 270–276. <http://doi.org/10.1016/j.jmpt.2009.03.009>
- Lefevre-Colau, M.-M., Poiraudau, S., Oberlin, C., Demaille, S., Fermanian, J., Rannou, F., & Revel, M. (2003). Reliability, validity, and responsiveness of the modified kapandji index for assessment of functional mobility of the rheumatoid hand. *Archives of Physical Medicine and Rehabilitation*, 84(7), 1032–1038. [http://doi.org/10.1016/S0003-9993\(03\)00128-X](http://doi.org/10.1016/S0003-9993(03)00128-X)
- Lennox, I. A., Murali, S. R., & Porter, R. (1993). A study of the repeatability of the diagnosis of Dupuytren's contracture and its prevalence in the grampian region. *Journal of Hand Surgery (Edinburgh, Scotland)*, 18(2), 258–261.
- Lowe, B. D. (2004). Accuracy and validity of observational estimates of wrist and forearm posture. *Ergonomics*.
- Luukkainen, R., Sanila, M. T., Saltyshev, M., Huhtala, H., & Koski, J. M. (2004). Relationship between clinically detected joint swelling and effusion diagnosed by ultrasonography in elbow joints in patients with rheumatoid arthritis. *Clinical Rheumatology*, 24(3), 228–231. <http://doi.org/10.1007/s10067-004-1010-8>
- Maffulli, N., Kenward, M. G., Testa, V., Capasso, G., Regine, R., & King, J. B. (2003). Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clinical Journal of Sport Medicine*, 13(1), 11–15.
- Mallo, G., Stanat, S. J. C., & Gaffney, J. (2010). Use of the Gartland Classification System for Treatment of Pediatric Supracondylar Humerus Fractures. *Orthopedics*, 33(1), 19–22. <http://doi.org/10.3928/01477447-20091124-08>
- Maricar, N., Callaghan, M. J., Parkes, M. J., Felson, D. T., & O'Neill, T. W. (2016a). Clinical assessment of effusion in knee osteoarthritis-A systematic review. *Seminars in Arthritis and Rheumatism*, 45(5), 556– 563. <http://doi.org/10.1016/j.semarthrit.2015.10.004>
- Maricar, N., Callaghan, M. J., Parkes, M. J., Felson, D. T., & O'Neill, T. W. (2016b). Interobserver and Intraobserver Reliability of Clinical Assessments in Knee Osteoarthritis. *The Journal of Rheumatology*, 43(12), 2171–2178. <http://doi.org/10.3899/jrheum.150835>

- Marx, R. G., Bombardier, C., & Wright, J. G. (1999). What do we know about the reliability and validity of physical examination tests used to examine the upper extremity? *Yjhsu*, 24(1), 185–193. <http://doi.org/10.1053/jhsu.1999.jhsu24a0185>
- Marx, R. G., Hudak, P. L., Bombardier, C., Graham, B., Goldsmith, C., & Wright, J. G. (1998). The reliability of physical examination for carpal tunnel syndrome. *Journal of Hand Surgery (Edinburgh, Scotland)*, 23(4), 499–502.
- May, S., & Ross, J. (2009). The McKenzie classification system in the extremities: a reliability study using Mckenzie assessment forms and experienced clinicians. *Journal of Manipulative and Physiological Therapeutics*, 32(7), 556–563. <http://doi.org/10.1016/j.jmpt.2009.08.007>
- Menz, H. B., Tiedemann, A., Kwan, M. M.-S., Latt, M. D., Sherrington, C., & Lord, S. R. (2003). Reliability of clinical tests of foot and ankle characteristics in older people. *Journal of the American Podiatric Medical Association*, 93(5), 380–387.
- Mishra, S. (2006). A new test for demonstrating the action of flexor digitorum superficialis (FDS) tendon. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 59(12), 1342–1344. <http://doi.org/10.1016/j.bjps.2006.03.068>
- Mohapatra, D. P., & Mohapatra, D. P. (2011). Thumb Tip Flexion Test. *Yjhsu*, 36(11), 1882. <http://doi.org/10.1016/j.jhsa.2011.08.024>
- Mora-Relucio, R., Núñez-Nagy, S., Gallego-Izquierdo, T., Rus, A., Plaza-Manzano, G., Romero-Franco, N., et al. (2016). Experienced versus Inexperienced Interexaminer Reliability on Location and Classification of Myofascial Trigger Point Palpation to Diagnose Lateral Epicondylalgia: An Observational Cross-Sectional Study. *Evidence-Based Complementary and Alternative Medicine : eCAM*, 2016, 6059719. <http://doi.org/10.1155/2016/6059719>
- Morganti, C. M., McFarland, E. G., & Cosgarea, A. J. (2002). Saphenous neuritis: a poorly understood cause of medial knee pain. *The Journal of the American Academy of Orthopaedic Surgeons*, 10(2), 130–137.
- Morris-Chatta, R., Buchner, D. M., de Lateur, B. J., Cress, M. E., & Wagner, E. H. (1994). Isokinetic testing of ankle strength in older adults: assessment of inter-rater reliability and stability of strength over six months. *Archives of Physical Medicine and Rehabilitation*, 75(11), 1213–1216.
- Mulligan, E. P., Harwell, J. L., & Robertson, W. J. (2011). Reliability and diagnostic accuracy of the Lachman test performed in a prone position. *The Journal of Orthopaedic and Sports Physical Therapy*, 41(10), 749–757. <http://doi.org/10.2519/jospt.2011.3761>
- Myers, H. L., Thomas, E., Hay, E. M., & Dziedzic, K. S. (2011). Hand assessment in older adults with musculoskeletal hand problems: a reliability study. *BMC Musculoskeletal Disorders*, 12(1), 3. <http://doi.org/10.1186/1471-2474-12-3>

- Ng, G., & Fan, A. (2005). Does elbow position affect strength and reproducibility of power grip measurements? *Physiotherapy*, 1–5.
- Nguyen, H. T., Resnick, D. N., Caldwell, S. G., Elston, E. W., Bishop, B. B., Steinhouser, J. B., et al. (1999). Interexaminer reliability of activator methods' relative leg-length evaluation in the prone extended position. *Journal of Manipulative and Physiological Therapeutics*, 22(9), 565–569.
- Nussbaum, E. D., Hosea, T. M., Sieler, S. D., Incremona, B. R., & Kessler, D. E. (2001). Prospective evaluation of syndesmotic ankle sprains without diastasis. *The American Journal of Sports Medicine*, 29(1), 31–35.
- Ochi, K., MD, K. O., MD, Y. H., Horiuchi, Y., MD, A. T., Tanabe, A., et al. (2012). Shoulder internal rotation elbow flexion test for diagnosing cubital tunnel syndrome. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 21(6), 777–781. <http://doi.org/10.1016/j.jse.2011.10.015>
- Ocs, M. M. M. P. C. C., Merritt, M. M., Roddey, T. S., FAAOMPT, T. S. R. P. P. O., Costello, C., CHT, C. C. P. P., et al. (2010). Diagnostic Value of Clinical Grind Test for Carpometacarpal Osteoarthritis of the Thumb. *Journal of Hand Therapy*, 23(3), 261–268. <http://doi.org/10.1016/j.jht.2010.02.001>
- Palmer, K. T., Palmer, K., Walker-Bone, K., Linaker, C., Reading, I., Kellingray, S., et al. (2007). The Southampton Examination Schedule for the diagnosis of musculoskeletal disorders of the upper limb. *La Medicina Del Lavoro*, 98(2), 111–117.
- Peat, G., Wood, L., Wilkie, R., Thomas, E., & for the KNE-SCI Study Group. (2003). How reliable is structured clinical history-taking in older adults with knee problems? *Journal of Clinical Epidemiology*, 56(11), 1030–1037. [http://doi.org/10.1016/S0895-4356\(03\)00204-X](http://doi.org/10.1016/S0895-4356(03)00204-X)
- Peeler, J., & Anderson, J. E. (2008). Reliability of the Ely's test for assessing rectus femoris muscle flexibility and joint range of motion. *Journal of Orthopaedic Research : Official Publication of the Orthopaedic Research Society*, 26(6), 793–799. <http://doi.org/10.1002/jor.20556>
- Petersen, S. M., & Covill, L. G. (2010). Reliability of the radial and ulnar nerve biased upper extremity neural tissue provocation tests. *Physiotherapy Theory and Practice*, 26(7), 476–482. <http://doi.org/10.3109/09593981003607629>
- Polzer, H., Kanz, K.-G., Prall, W. C., Haasters, F., Ockert, B., Mutschler, W., & Grote, S. (2012). Diagnosis and treatment of acute ankle injuries: development of an evidence-based algorithm. *Orthopedic Reviews*, 4(1), e5. <http://doi.org/10.4081/or.2012.e5>
- Poulsen, D. R., & James, C. R. (2011). Concurrent validity and reliability of clinical evaluation of the single leg squat. *Physiotherapy Theory and Practice*, 27(8), 586–594. <http://doi.org/10.3109/09593985.2011.552539>

- Powden, C. J., Hoch, J. M., & Hoch, M. C. (2015). Reliability and minimal detectable change of the weight-bearing lunge test: A systematic review. *Manual Therapy*, 20(4), 524–532. <http://doi.org/10.1016/j.math.2015.01.004>
- Prosser, R., Harvey, L., LaStayo, P., Hargreaves, I., Scougall, P., & Herbert, R. D. (2011). Provocative wrist tests and MRI are of limited diagnostic value for suspected wrist ligament injuries: a cross-sectional study. *Journal of Physiotherapy*, 57(4), 247–253. [http://doi.org/10.1016/S1836-9553\(11\)70055-8](http://doi.org/10.1016/S1836-9553(11)70055-8)
- Proud, E. L., Miller, K. J., Bilney, B., Balachandran, S., McGinley, J. L., & Morris, M. E. (2015). Evaluation of measures of upper limb functioning and disability in people with Parkinson disease: a systematic review. *Archives of Physical Medicine and Rehabilitation*, 96(3), 540–551.e1. <http://doi.org/10.1016/j.apmr.2014.09.016>
- Rabin, A., Kozol, Z., Spitzer, E., & Finestone, A. (2014). Ankle dorsiflexion among healthy men with different qualities of lower extremity movement. *Journal of Athletic Training*, 49(5), 617–623. <http://doi.org/10.4085/1062-6050-49.3.14>
- Redler, L. H., Watling, J. P., Dennis, E. R., Swart, E., & Ahmad, C. S. (2016). Reliability of a field-based drop vertical jump screening test for ACL injury risk assessment. *The Physician and Sportsmedicine*, 44(1), 46–52. <http://doi.org/10.1080/00913847.2016.1131107>
- Remvig, L., Jensen, D. V., & Ward, R. C. (2007). Are diagnostic criteria for general joint hypermobility and benign joint hypermobility syndrome based on reproducible and valid tests? A review of the literature.
- Rosen, A. B., Ko, J., & Brown, C. N. (2015). Diagnostic accuracy of instrumented and manual talar tilt tests in chronic ankle instability populations. *Scandinavian Journal of Medicine & Science in Sports*, 25(2), e214–21. <http://doi.org/10.1111/sms.12288>
- Saban, B., & Masharawi, Y. (2016). Three single leg standing tests for clinical assessment of chronic plantar heel pain syndrome: static stance, half-squat and heel rise. *Physiotherapy*. <http://doi.org/10.1016/j.physio.2016.06.003>
- Sacco, I. C. N., Onodera, A. N., Butugan, M. K., Taddei, U. T., Mendes, Y. C., Galhardo, B., et al. (2010). Inter- and intra-tester reliability of clinical measurement to determine medio-lateral patellar position using a pachymeter or visual assessment. *The Knee*, 17(1), 92–95. <http://doi.org/10.1016/j.knee.2009.05.008>
- Salerno, D. F., Franzblau, A., Werner, R. A., Chung, K. C., Schultz, J. S., Becker, M. P., & Armstrong, T. J. (2000). Reliability of physical examination of the upper extremity among keyboard operators. *American Journal of Industrial Medicine*, 37(4), 423–430.
- Sanz, D. R., Lobo, C. C., López, D. L., Morales, C. R., Marín, C. S., & Corbalán, I. S. (2016). Interrater Reliability in the Clinical Evaluation of Myofascial Trigger Points in Three Ankle Muscles. *Journal of Manipulative and Physiological Therapeutics*, 39(9), 623–634. <http://doi.org/10.1016/j.jmpt.2016.09.002>

- Savva, C., Karagiannis, C., & Rushton, A. (2013). Test-retest reliability of grip strength measurement in full elbow extension to evaluate maximum grip strength. *Journal of Hand Surgery (European Volume)*, 38(2), 183–186. <http://doi.org/10.1177/1753193412449804>
- Schneider, M., Homonai, R., Moreland, B., & Delitto, A. (2007). Interexaminer Reliability of the Prone Leg Length Analysis Procedure. *Journal of Manipulative and Physiological Therapeutics*, 30(7), 514– 521. <http://doi.org/10.1016/j.jmpt.2007.07.001>
- Schwieterman, B., Haas, D., Columber, K., Knupp, D., & Cook, C. (2013). Diagnostic accuracy of physical examination tests of the ankle/foot complex: a systematic review. *International Journal of Sports Physical Therapy*, 8(4), 416–426.
- Sebastian, D., Chovvath, R., & Malladi, R. (2014). The sitting active and prone passive lag test: an inter- rater reliability study. *Journal of Bodywork and Movement Therapies*, 18(2), 204–209. <http://doi.org/10.1016/j.jbmt.2013.08.002>
- Senocak, O., Unsal, E., Akalin, E., & Ergör, G. (2003). Interobserver reliability of articular examination in juvenile idiopathic arthritis. *The Turkish Journal of Pediatrics*, 45(1), 29–32.
- Simonsen, O., Jensen, J., Mouritsen, P., & Lauritzen, J. (1984). The accuracy of clinical examination of injury of the knee joint. *Injury*, 16(2), 96–101.
- Sman, A. D., Hiller, C. E., & Refshauge, K. M. (2013). Diagnostic accuracy of clinical tests for diagnosis of ankle syndesmosis injury: a systematic review. *British Journal of Sports Medicine*, 47(10), 620–628. <http://doi.org/10.1136/bjsports-2012-091702>
- Smidt, N., van der Windt, D. A., & Assendelft, W. J. (2002). Interobserver reproducibility of the assessment of severity of complaints, grip strength, and pressure pain threshold in patients with lateral epicondylitis. *Archives of Physical ...*
- Smith, C. K., Bonauto, D. K., Silverstein, B. A., & Wilcox, D. (2010). Inter-Rater Reliability of Physical Examinations in a Prospective Study of Upper Extremity Musculoskeletal Disorders. *Journal of Occupational and Environmental Medicine*, 52(10), 1014–1018. <http://doi.org/10.1097/JOM.0b013e3181f4396b>
- Smith, T. O., Clark, A., Neda, S., Arendt, E. A., Post, W. R., Grelsamer, R. P., et al. (2012). The intra- and inter-observer reliability of the physical examination methods used to assess patients with patellofemoral joint instability. *The Knee*, 19(4), 404–410. <http://doi.org/10.1016/j.knee.2011.06.002>
- Snoeker, B. A. M., Lindeboom, R., Zwinderman, A. H., Vincken, P. W. J., Jansen, J. A., & Lucas, C. (2015). Detecting Meniscal Tears in Primary Care: Reproducibility and Accuracy of 2 Weight-Bearing Tests and 1 Non-Weight-Bearing Test. *The Journal of Orthopaedic and Sports Physical Therapy*, 45(9), 693–702. <http://doi.org/10.2519/jospt.2015.5712>
- Solomon, D. J., Simel, D. L., Bates, D. W., Katz, J. N., & Schaffer, J. L. (2001). Does This Patient Have a Torn Meniscus or Ligament of the Knee? *Jama*, 286(13), 1610–1620.

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
Upper Extremity Region

- Stiell, I. G., McKnight, R. D., Greenberg, G. H., Nair, R. C., McDowell, I., & Wallace, G. J. (1992). Interobserver agreement in the examination of acute ankle injury patients. *The American Journal of Emergency Medicine*, 10(1), 14–17.
- Stratford, P. W., Norman, G. R., & McIntosh, J. M. (1989). Generalizability of grip strength measurements in patients with tennis elbow. *Physical Therapy*, 69(4), 276–281.
- Sturgill, L. P. (2009). Interrater Reliability of a Clinical Scale to Assess Knee Joint Effusion. *The Journal of Orthopaedic and Sports Physical Therapy*. <http://doi.org/10.2519/jospt.2009.3143>
- Takeuchi, N., Kuwabara, T., & Usuda, S. (2009). Development and Evaluation of a New Measure for Muscle Tone of Ankle Plantar Flexors: the Ankle Plantar Flexors Tone Scale. *Archives of Physical Medicine and Rehabilitation*, 90(12), 2054–2061. <http://doi.org/10.1016/j.apmr.2009.08.141>
- Talebi, G. A., Oskouei, A. E., & Shakori, S. K. (2012). Reliability of upper limb tension test 1 in normal subjects and patients with carpal tunnel syndrome. *Journal of Back and Musculoskeletal Rehabilitation*, 25(3), 209–214. <http://doi.org/10.3233/BMR-2012-0330>
- Thaller, M., & Hughes, T. (2014). Inter-rater agreement of observable and elicitable neurological signs. *Clinical Medicine (London, England)*, 14(3), 264–267. <http://doi.org/10.7861/clinmedicine.14-3-264>
- The, B., PhD, B. T. M., Reininga, I. H. F., PhD, I. H. F. R., Moumni, El, M., Moumni MD, El, M., et al. (2013). Elbow-specific clinical rating systems: extent of established validity, reliability, and responsiveness. *Journal of Shoulder and Elbow Surgery / American Shoulder and Elbow Surgeons [Et Al]*, 22(10), 1380–1394. <http://doi.org/10.1016/j.jse.2013.04.013>
- Trojan, T. H. (2006). Single leg balance test to identify risk of ankle sprains \* Commentary 1 \* Commentary 2. *British Journal of Sports Medicine*, 40(7), 610–613. <http://doi.org/10.1136/bjism.2005.024356>
- Tucci, H. T., Martins, J., Sposito, G. de C., Camarini, P. M. F., & de Oliveira, A. S. (2014). Closed Kinetic Chain Upper Extremity Stability test (CKCUES test): a reliability study in persons with and without shoulder impingement syndrome. *BMC Musculoskeletal Disorders*, 15, 1. <http://doi.org/10.1186/1471-2474-15-1>
- Turchin, D. C., Beaton, D. E., & Richards, R. R. (1998). Validity of observer-based aggregate scoring systems as descriptors of elbow pain, function, and disability. *The Journal of Bone and Joint Surgery. American Volume*, 80(2), 154–162.
- Ulaşlı, A. M., Yaman, F., Dikici, Ö., Karaman, A., Kaçar, E., & Demirdal, Ü. S. (2014). Accuracy in detecting knee effusion with clinical examination and the effect of effusion, the patient's body mass index, and the clinician's experience. *Clinical Rheumatology*, 33(8), 1139–1143. <http://doi.org/10.1007/s10067-013-2356-6>

- Urban, V., KALBERER, F., Roos, M., & Dumont, C. E. (2002). [Reliability of active range-of-motion measurement of the rotation in the forearm: comparison of three measurement devices]. *Zeitschrift Für Orthopädie Und Ihre Grenzgebiete*, 140(1), 72–76.  
<http://doi.org/10.1055/s-2002-22094>
  
- van der Lee, J. H., de Groot, V., Beckerman, H., Wagenaar, R. C., Lankhorst, G. J., & Bouter, L. M. (2001). The intra- and interrater reliability of the action research arm test: A practical test of upper extremity function in patients with stroke. *Archives of Physical Medicine and Rehabilitation*, 82(1), 14–19. <http://doi.org/10.1053/apmr.2001.18668>
  
- van Dijk, C. N., Lim, L. S., Bossuyt, P. M., & Marti, R. K. (1996a). Physical examination is sufficient for the diagnosis of sprained ankles. *The Journal of Bone and Joint Surgery. British Volume*, 78(6), 958–962.
  
- van Dijk, C. N., Mol, B. W., Lim, L. S., Marti, R. K., & Bossuyt, P. M. (1996b). Diagnosis of ligament rupture of the ankle joint. Physical examination, arthrography, stress radiography and sonography compared in 160 patients after inversion trauma. *Acta Orthopaedica Scandinavica*, 67(6), 566–570. <http://doi.org/10.3109/17453679608997757>
  
- van Trijffel, E., van de Pol, R. J., Oostendorp, R. A., & Lucas, C. (2010). Inter-rater reliability for measurement of passive physiological movements in lower extremity joints is generally low: a systematic review. *Journal of Physiotherapy*, 56(4), 223–235.
  
- Vanpee, G., Hermans, G., Segers, J., & Gosselink, R. (2014). Assessment of Limb Muscle Strength in Critically Ill Patients. *Critical Care Medicine*, 42(3), 701–711.  
<http://doi.org/10.1097/CCM.0000000000000030>
  
- Wadey, V. M. R., Mohtadi, N. G. H., Bray, R. C., & Frank, C. B. (2007). Positive predictive value of maximal posterior joint-line tenderness in diagnosing meniscal pathology: a pilot study. *Canadian Journal of Surgery. Journal Canadien De Chirurgie*, 50(2), 96–100.
  
- Wadsworth, C. T., Krishnan, R., Sear, M., Harrold, J., & Nielsen, D. H. (1987). Intrarater reliability of manual muscle testing and hand-held dynametric muscle testing. *Physical Therapy*, 67(9), 1342–1347.
  
- Wai, E. K., Eugene K Wai MD, M. F., Kelly Howse BSc, M. D., Howse, K., Pollock, J. W., J Whitcomb Pollock MD, F., et al. (2009). The reliability of determining “leg dominant pain”. *The Spine Journal : Official Journal of the North American Spine Society*, 9(6), 447–453. <http://doi.org/10.1016/j.spinee.2008.11.009>
  
- Wainner, R. S., Fritz, J. M., Irrgang, J. J., Delitto, A., Allison, S., & Boninger, M. L. (2005). Development of a clinical prediction rule for the diagnosis of carpal tunnel syndrome. *Archives of Physical Medicine and Rehabilitation*, 86(4), 609–618.  
<http://doi.org/10.1016/j.apmr.2004.11.008>

- Walker-Bone, K. E., Palmer, K. T., Reading, I., & Cooper, C. (2003). Criteria for assessing pain and nonarticular soft-tissue rheumatic disorders of the neck and upper limb. *Seminars in Arthritis and Rheumatism*, 33(3), 168–184. [http://doi.org/10.1016/S0049-0172\(03\)00129-X](http://doi.org/10.1016/S0049-0172(03)00129-X)
- Walker-Bone, K., Byng, P., Linaker, C., Reading, I., Coggon, D., Palmer, K. T., & Cooper, C. (2002). Reliability of the Southampton examination schedule for the diagnosis of upper limb disorders in the general population. *Annals of the Rheumatic Diseases*, 61(12), 1103–1106.
- Walsh, J., & Hall, T. (2009). Reliability, validity and diagnostic accuracy of palpation of the sciatic, tibial and common peroneal nerves in the examination of low back related leg pain. *Manual Therapy*, 14(6), 623–629. <http://doi.org/10.1016/j.math.2008.12.007>
- Waters, P. M., Zurakowski, D., Patterson, P., Bae, D. S., & Nimec, D. (2004). Interobserver and intraobserver reliability of therapist-assisted videotaped evaluations of upper-limb hemiplegia. *The Journal of Hand Surgery*, 29(2), 328–334. <http://doi.org/10.1016/j.jhsa.2003.10.014>
- Watkins, M. A., Riddle, D. L., Lamb, R. L., & Personius, W. J. (1991). Reliability of goniometric measurements and visual estimates of knee range of motion obtained in a clinical setting. *Physical Therapy*, 71(2), 90–6– discussion 96–7.
- Watson, C. J., Leddy, H. M., Dynjan, T. D., & Parham, J. L. (2001). Reliability of the lateral pull test and tilt test to assess patellar alignment in subjects with symptomatic knees: student raters. *The Journal of Orthopaedic and Sports Physical Therapy*, 31(7), 368–374. <http://doi.org/10.2519/jospt.2001.31.7.368>
- Wood, L., Peat, G., Wilkie, R., Hay, E., Thomas, E., & Sim, J. (2006). A study of the noninstrumented physical examination of the knee found high observer variability. *Journal of Clinical Epidemiology*, 59(5), 512–520. <http://doi.org/10.1016/j.jclinepi.2005.11.004>
- Wrobel, J. S., & Armstrong, D. G. (2008). Reliability and validity of current physical examination techniques of the foot and ankle. *Journal of the American Podiatric Medical Association*, 98(3), 197– 206.
- Yasui, K., Mihata, T., Takeda, A., Watanabe, C., & Kinoshita, M. (2012). A new manual method for assessing elbow valgus laxity. *Sports Medicine, Arthroscopy, Rehabilitation, Therapy & Technology*, 4(1), 11. <http://doi.org/10.1186/1758-2555-4-11>

## Lower Extremity Region

- Abouelela, A. A. K. H., & Zohiery, A. K. (2012). The triple compression stress test for diagnosis of tarsal tunnel syndrome. *Foot (Edinburgh, Scotland)*, 22(3), 146–149. <http://doi.org/10.1016/j.foot.2012.02.002>
- Ageberg, E., Bennell, K. L., Hunt, M. A., Simic, M., Roos, E. M., & Creaby, M. W. (2010). Validity and inter-rater reliability of medio-lateral knee motion observed during a single-limb minisquat. *BMC Musculoskeletal Disorders*, 11(1), 265. <http://doi.org/10.1186/1471-2474-11-265>
- AKGUL, O., Guldeste, Z., & Ozgocmen, S. (2014). The reliability of the clinical examination for detecting Baker's cyst in asymptomatic fossa. *International Journal of Rheumatic Diseases*, 17(2), 204–209. <http://doi.org/10.1111/1756-185X.12095>
- Allington, N. J., Leroy, N., & Doneux, C. (2002). Ankle joint range of motion measurements in spastic cerebral palsy children: intraobserver and interobserver reliability and reproducibility of goniometry and visual estimation. *Journal of Pediatric Orthopaedics. Part B / European Paediatric Orthopaedic Society, Pediatric Orthopaedic Society of North America*, 11(3), 236–239.
- Alonso, A., Khoury, L., & Adams, R. (1998). Clinical tests for ankle syndesmosis injury: reliability and prediction of return to function. *The Journal of Orthopaedic and Sports Physical Therapy*, 27(4), 276–284. <http://doi.org/10.2519/jospt.1998.27.4.276>
- Ames, R., Jarosz, B. S., Baglin, J., & Da Costa, C. (2015). Functional Knee Extension Test (FKET): Inter-rater palpatory agreement and visual analysis during a non-weight bearing functional assessment of the knee. *Journal of Bodywork and Movement Therapies*, 19(4), 656–665. <http://doi.org/10.1016/j.jbmt.2014.12.005>
- Beumer, A., Swierstra, B. A., & Mulder, P. G. H. (2002). Clinical diagnosis of syndesmotic ankle instability: evaluation of stress tests behind the curtains. *Acta Orthopaedica Scandinavica*, 73(6), 667–669. <http://doi.org/10.1080/000164702321039642>
- Boeree, N. R., & Ackroyd, C. E. (1991). Assessment of the menisci and cruciate ligaments: an audit of clinical practice. *Injury*, 22(4), 291–294.
- Brandsma, J. W., Van Brakel, W. H., Anderson, A. M., Kortendijk, A. J., Gurung, K. S., & Sunwar, S. K. (1998). Intertester reliability of manual muscle strength testing in leprosy patients. *Leprosy Review*, 69(3), 257–266.
- Cejudo, A., Sainz de Baranda, P., Ayala, F., & Santonja, F. (2014). A simplified version of the weight-bearing ankle lunge test: description and test-retest reliability. *Manual Therapy*, 19(4), 355–359. <http://doi.org/10.1016/j.math.2014.03.008>
- Chang, C.-W., Wang, Y.-C., & Chu, C.-H. (2008). Increased Carrying Angle is a Risk Factor for Nontraumatic Ulnar Neuropathy at the Elbow. *Clinical Orthopaedics and Related Research*, 466(9), 2190–2195. <http://doi.org/10.1007/s11999-008-0308-2>

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/musculoskeletal medicine  
**Lower Extremity Region**

- Chatterton, B. D., Muller, S., Thomas, M. J., Menz, H. B., Rome, K., & Roddy, E. (2013). Inter and intra- rater repeatability of the scoring of foot pain drawings. *Journal of Foot and Ankle Research*, 6(1), 1–1. <http://doi.org/10.1186/1757-1146-6-44>
- Cibere, J., Bellamy, N., Thorne, A., Esdaile, J. M., McGorm, K. J., Chalmers, A., et al. (2004). Reliability of the knee examination in osteoarthritis: Effect of standardization. *Arthritis and Rheumatism*, 50(2), 458–468. <http://doi.org/10.1002/art.20025>
- Cook, J. L., Khan, K. M., Kiss, Z. S., Purdam, C. R., & Griffiths, L. (2001). Reproducibility and clinical utility of tendon palpation to detect patellar tendinopathy in young basketball players. Victorian Institute of Sport tendon study group. *British Journal of Sports Medicine*, 35(1), 65–69.
- Cooperman, J. M., Riddle, D. L., & Rothstein, J. M. (1990). Reliability and validity of judgments of the integrity of the anterior cruciate ligament of the knee using the Lachman's test. *Physical Therapy*, 70(4), 225–233.
- Cornwall, M. W., Fishco, W. D., McPoil, T. G., Lane, C. R., O'Donnell, D., & Hunt, L. (2004). Reliability and validity of clinically assessing first-ray mobility of the foot. *Journal of the American Podiatric Medical Association*, 94(5), 470–476.
- Currier, L. L., Froehlich, P. J., Carow, S. D., McAndrew, R. K., Cliborne, A. V., Boyles, R. E., et al. (2007). Development of a clinical prediction rule to identify patients with knee pain and clinical evidence of knee osteoarthritis who demonstrate a favorable short-term response to hip mobilization. *Physical Therapy*, 87(9), 1106–1119. <http://doi.org/10.2522/ptj.20060066>
- Cushnaghan, J., Cooper, C., Dieppe, P., Kirwan, J., McAlindon, T., & McCrae, F. (1990). Clinical assessment of osteoarthritis of the knee. *Annals of the Rheumatic Diseases*, 49(10), 768–770.
- Dennis, R. J., Finch, C. F., Elliott, B. C., & Farhart, P. J. (2008). The reliability of musculoskeletal screening tests used in cricket. *Physical Therapy in Sport : Official Journal of the Association of Chartered Physiotherapists in Sports Medicine*, 9(1), 25–33. <http://doi.org/10.1016/j.ptsp.2007.09.004>
- Derksen, R.-J., Bakker, F. C., Geervliet, P. C., de Lange de Klerk, E. S. M., Heilbron, E. A., Veenings, B., et al. (2005). Diagnostic accuracy and reproducibility in the interpretation of Ottawa ankle and foot rules
- by specialized emergency nurses. *The American Journal of Emergency Medicine*, 23(6), 725–729. <http://doi.org/10.1016/j.ajem.2005.02.054>
- Dervin, G. F., Stiell, I. G., Wells, G. A., Rody, K., & Grabowski, J. (2001). Physicians' accuracy and interrater reliability for the diagnosis of unstable meniscal tears in patients having osteoarthritis of the knee. *Canadian Journal of Surgery. Journal Canadien De Chirurgie*, 44(4), 267–274.

**Further Reading**, Reproducibility protocol for diagnostic procedures in manual/muskuloskeletal medicine  
**Lower Extremity Region**

- Digiovanni, C. W., Holt, S., Czerniecki, J. M., Ledoux, W. R., & Sangeorzan, B. J. (2001). Can the presence of equinus contracture be established by physical exam alone? *Journal of Rehabilitation Research and Development*, 38(3), 335–340.
- Elson, D. W., Jones, S., Caplan, N., Stewart, S., St Clair Gibson, A., Gibson, A. S. C., & Kader, D. F. (2011). The photographic knee pain map: Locating knee pain with an instrument developed for diagnostic, communication and research purposes. *The Knee*, 18(6), 417–423. <http://doi.org/10.1016/j.knee.2010.08.012>
- Eren, O. T. (2003). The accuracy of joint line tenderness by physical examination in the diagnosis of meniscal tears. *Arthroscopy : the Journal of Arthroscopic & Related Surgery : Official Publication of the Arthroscopy Association of North America and the International Arthroscopy Association*, 19(8), 850–854. [http://doi.org/10.1053/S0749-8063\(03\)00736-9](http://doi.org/10.1053/S0749-8063(03)00736-9)
- Erichsen, N., Lund, H., Ole Møller, J., Kaiser, T., Lykke Jensen, M., Mårcher, I., et al. (2006). Inter-rater and intra-rater reliability of tests of translatory movements and range of movements in the subtalar and talocrural joints. *Advances in Physiotherapy*, 8(4), 161–167. <http://doi.org/10.1080/14038190601000066>
- Fredericson, M., & Yoon, K. (2006). Physical Examination and Patellofemoral Pain Syndrome. *American Journal of Physical Medicine & Rehabilitation / Association of Academic Physiatrists*, 85(3), 234–243. <http://doi.org/10.1097/01.phm.0000200390.67408.f0>
- Fritz, J. M., Delitto, A., Erhard, R. E., & Roman, M. (1998). An examination of the selective tissue tension scheme, with evidence for the concept of a capsular pattern of the knee. *Physical Therapy*, 78(10), 1046–56– discussion 1057–61.
- Gatt, A., & Chockalingam, N. (2012). Assessment of Ankle Joint Dorsiflexion: An Overview. *Revista Internacional De Ciencias Podológicas*, 6(1), 25–29. [http://doi.org/10.5209/rev\\_RICP.2012.v6.n1.37894](http://doi.org/10.5209/rev_RICP.2012.v6.n1.37894)
- Gibbons, P., Dumper, C., & Gosling, C. (2000). Inter-examiner and intra-examiner agreement for assessing simulated leg length inequality using palpation and observation during a standing assessment. *Journal of Osteopathic Medicine*, 5(2), 53–58.
- H Charles Woodfield RPh, D. C., Woodfield, H. C., Gerstman, B. B., B Burt Gerstman DVM, M. P., Olaisen, R. H., Renate Henry Olaisen DC, M., et al. (2011). Interexaminer Reliability of Supine Leg Checks for Discriminating Leg-Length Inequality. *Journal of Manipulative and Physiological Therapeutics*, 34(4), 239–246. <http://doi.org/10.1016/j.jmpt.2011.04.009>
- Harris-Hayes, M., Steger-May, K., Koh, C., Royer, N. K., Graci, V., & Salsich, G. B. (2014). Classification of lower extremity movement patterns based on visual assessment: reliability and correlation with 2- dimensional video analysis. *Journal of Athletic Training*, 49(3), 304–310. <http://doi.org/10.4085/1062-6050-49.2.21>

- Hart, D. J., Spector, T. D., Brown, P., Wilson, P., Doyle, D. V., & Silman, A. J. (1991). Clinical signs of early osteoarthritis: reproducibility and relation to x ray changes in 541 women in the general population. *Annals of the Rheumatic Diseases*, 50(7), 467–470.
- Hauzeur, J. P., Mathy, L., & De Maertelaer, V. (1999). Comparison between clinical evaluation and ultrasonography in detecting hydrarthrosis of the knee. *The Journal of Rheumatology*, 26(12), 2681– 2683.
- Hayes, K. W., & Petersen, C. M. (2001). Reliability of assessing end-feel and pain and resistance sequence in subjects with painful shoulders and knees. *The Journal of Orthopaedic and Sports Physical Therapy*, 31(8), 432–445.
- Hayes, K. W., Petersen, C., & Falconer, J. (1994b). An examination of Cyriax's passive motion tests with patients having osteoarthritis of the knee. *Physical Therapy*, 74(8), 697–707–discussion 707–9.
- Hsieh, C. Y., Hong, C. Z., Adams, A. H., Platt, K. J., Danielson, C. D., Hoehler, F. K., & Tobis, J. S. (2000). Interexaminer reliability of the palpation of trigger points in the trunk and lower limb muscles. *Archives of Physical Medicine and Rehabilitation*, 81(3), 258–264.
- Hutchison, A.-M., Evans, R., Bodger, O., Pallister, I., Topliss, C., Williams, P., et al. (2013). Foot and Ankle Surgery. *Foot and Ankle Surgery*, 19(2), 112–117. <http://doi.org/10.1016/j.fas.2012.12.006>
- Jarvis, H. L., Nester, C. J., Jones, R. K., Williams, A., & Bowden, P. D. (2012). Inter-assessor reliability of practice based biomechanical assessment of the foot and ankle. *Journal of Foot and Ankle Research*, 5, 14. <http://doi.org/10.1186/1757-1146-5-14>
- Jensen, L. K., Mikkelsen, S., Loft, I. P., & Eenberg, W. (2000). Work-related knee disorders in floor layers and carpenters. *Journal of Occupational and Environmental Medicine*, 42(8), 835–842.
- Jepsen, J. R., Laursen, L. H., Hagert, C.-G., Kreiner, S., & Larsen, A. I. (2006). Diagnostic accuracy of the neurological upper limb examination I: inter-rater reproducibility of selected findings and patterns. *BMC Neurology*, 6, 8. <http://doi.org/10.1186/1471-2377-6-8>
- Jones, A., Hopkinson, N., Pattrick, M., Berman, P., & Doherty, M. (1992). Evaluation of a method for clinically assessing osteoarthritis of the knee. *Annals of the Rheumatic Diseases*, 51(2), 243–245.
- unge, T., Balsnes, S. L., Runge, L., Juul-Kristensen, B., & Wedderkopp, N. (2012). Single leg mini squat: an inter-tester reproducibility study of children in the age of 9–10 and 12–14 years presented by various methods of kappa calculation. *BMC Musculoskeletal Disorders*, 13(1), 1–1. <http://doi.org/10.1186/1471-2474-13-203>

- Karim, Z., Wakefield, R. J., Quinn, M., Conaghan, P. G., Brown, A. K., Veale, D. J., et al. (2004). Validation and reproducibility of ultrasonography in the detection of synovitis in the knee: A comparison with arthroscopy and clinical examination. *Arthritis and Rheumatism*, 50(2), 387–394. <http://doi.org/10.1002/art.20054>
- Kaya, T., Karatepe, A. G., Goksel Karatepe, A., Gunaydin, R., Koc, A., & Altundal Ercan, U. (2011). Inter-rater reliability of the Modified Ashworth Scale and modified Modified Ashworth Scale in assessing poststroke elbow flexor spasticity. *International Journal of Rehabilitation Research*, 34(1), 59–64. <http://doi.org/10.1097/MRR.0b013e32833d6cdf>
- Kelly R Holt BSc Chiro, P., Chiro, D. G. R. B., Nicholas J Hoffmann BHLthSci, B. C., Chiro, B. I. B. B., Chiro, P. M. B. B., & Heidi Haavik Taylor BSc Chiro, P. (2009). Interexaminer Reliability of a Leg Length Analysis Procedure Among Novice and Experienced Practitioners. *Journal of Manipulative and Physiological Therapeutics*, 32(3), 216–222. <http://doi.org/10.1016/j.jmpt.2009.02.009>
- Kiter, E., & Bozkurt, M. (2005). The crossed-leg test for examination of ankle syndesmosis injuries. *Foot & Ankle International / American Orthopaedic Foot and Ankle Society [and] Swiss Foot and Ankle Society*, 26(2), 187–188.
- Kwon, B. C., Kwon, B. C., Choi, S. J., Choi, S.-J., Song, S.-Y., Song, S. Y., et al. (2011). Modified Carpal Stretch Test as a Screening Test for Detection of Scapholunate Interosseous Ligament Injuries Associated with Distal Radial Fractures. *The Journal of Bone and Joint Surgery*, 93(9), 855–862. <http://doi.org/10.2106/JBJS.J.00361>
- Leard, J. S., EdD, J. S. L., PhD, B. A. C., Crane, B. A., PhD, K. A. B., & Ball, K. A. (2009). Intrarater and interrater reliability of 22 clinical measures associated with lower quarter malalignment. *Journal of Manipulative and Physiological Therapeutics*, 32(4), 270–276. <http://doi.org/10.1016/j.jmpt.2009.03.009>
- Lefevre-Colau, M.-M., Poiraudau, S., Oberlin, C., Demaille, S., Fermanian, J., Rannou, F., & Revel, M. (2003). Reliability, validity, and responsiveness of the modified kapandji index for assessment of functional mobility of the rheumatoid hand. *Archives of Physical Medicine and Rehabilitation*, 84(7), 1032–1038. [http://doi.org/10.1016/S0003-9993\(03\)00128-X](http://doi.org/10.1016/S0003-9993(03)00128-X)
- Maffulli, N., Kenward, M. G., Testa, V., Capasso, G., Regine, R., & King, J. B. (2003). Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clinical Journal of Sport Medicine*, 13(1), 11–15.
- Maricar, N., Callaghan, M. J., Parkes, M. J., Felson, D. T., & O'Neill, T. W. (2016a). Clinical assessment of effusion in knee osteoarthritis-A systematic review. *Seminars in Arthritis and Rheumatism*, 45(5), 556– 563. <http://doi.org/10.1016/j.semarthrit.2015.10.004>
- Maricar, N., Callaghan, M. J., Parkes, M. J., Felson, D. T., & O'Neill, T. W. (2016b). Interobserver and Intraobserver Reliability of Clinical Assessments in Knee Osteoarthritis. *The Journal of Rheumatology*, 43(12), 2171–2178. <http://doi.org/10.3899/jrheum.150835>

- May, S., & Ross, J. (2009). The McKenzie classification system in the extremities: a reliability study using McKenzie assessment forms and experienced clinicians. *Journal of Manipulative and Physiological Therapeutics*, 32(7), 556–563. <http://doi.org/10.1016/j.jmpt.2009.08.007>
- Menz, H. B., Tiedemann, A., Kwan, M. M.-S., Latt, M. D., Sherrington, C., & Lord, S. R. (2003). Reliability of clinical tests of foot and ankle characteristics in older people. *Journal of the American Podiatric Medical Association*, 93(5), 380–387.
- Mohapatra, D. P., & Mohapatra, D. P. (2011). Thumb Tip Flexion Test. *Yjhsu*, 36(11), 1882. <http://doi.org/10.1016/j.jhsa.2011.08.024>
- Morganti, C. M., McFarland, E. G., & Cosgarea, A. J. (2002). Saphenous neuritis: a poorly understood cause of medial knee pain. *The Journal of the American Academy of Orthopaedic Surgeons*, 10(2), 130–137.
- Morris-Chatta, R., Buchner, D. M., de Lateur, B. J., Cress, M. E., & Wagner, E. H. (1994). Isokinetic testing of ankle strength in older adults: assessment of inter-rater reliability and stability of strength over six months. *Archives of Physical Medicine and Rehabilitation*, 75(11), 1213–1216.
- Mulligan, E. P., Harwell, J. L., & Robertson, W. J. (2011). Reliability and diagnostic accuracy of the Lachman test performed in a prone position. *The Journal of Orthopaedic and Sports Physical Therapy*, 41(10), 749–757. <http://doi.org/10.2519/jospt.2011.3761>
- Nguyen, H. T., Resnick, D. N., Caldwell, S. G., Elston, E. W., Bishop, B. B., Steinhouser, J. B., et al. (1999). Interexaminer reliability of activator methods' relative leg-length evaluation in the prone extended position. *Journal of Manipulative and Physiological Therapeutics*, 22(9), 565–569.
- Nussbaum, E. D., Hosea, T. M., Sieler, S. D., Incremona, B. R., & Kessler, D. E. (2001). Prospective evaluation of syndesmotic ankle sprains without diastasis. *The American Journal of Sports Medicine*, 29(1), 31–35.
- Peat, G., Wood, L., Wilkie, R., Thomas, E., & for the KNE-SCI Study Group. (2003). How reliable is structured clinical history-taking in older adults with knee problems? *Journal of Clinical Epidemiology*, 56(11), 1030–1037. [http://doi.org/10.1016/S0895-4356\(03\)00204-X](http://doi.org/10.1016/S0895-4356(03)00204-X)
- Peeler, J., & Anderson, J. E. (2008). Reliability of the Ely's test for assessing rectus femoris muscle flexibility and joint range of motion. *Journal of Orthopaedic Research : Official Publication of the Orthopaedic Research Society*, 26(6), 793–799. <http://doi.org/10.1002/jor.20556>
- Polzer, H., Kanz, K.-G., Prall, W. C., Haasters, F., Ockert, B., Mutschler, W., & Grote, S. (2012). Diagnosis and treatment of acute ankle injuries: development of an evidence-based algorithm. *Orthopedic Reviews*, 4(1), e5. <http://doi.org/10.4081/or.2012.e5>

- Poulsen, D. R., & James, C. R. (2011). Concurrent validity and reliability of clinical evaluation of the single leg squat. *Physiotherapy Theory and Practice*, 27(8), 586–594. <http://doi.org/10.3109/09593985.2011.552539>
- Powden, C. J., Hoch, J. M., & Hoch, M. C. (2015). Reliability and minimal detectable change of the weight-bearing lunge test: A systematic review. *Manual Therapy*, 20(4), 524–532. <http://doi.org/10.1016/j.math.2015.01.004>
- Rabin, A., Kozol, Z., Spitzer, E., & Finestone, A. (2014). Ankle dorsiflexion among healthy men with different qualities of lower extremity movement. *Journal of Athletic Training*, 49(5), 617–623. <http://doi.org/10.4085/1062-6050-49.3.14>
- Redler, L. H., Watling, J. P., Dennis, E. R., Swart, E., & Ahmad, C. S. (2016). Reliability of a field-based drop vertical jump screening test for ACL injury risk assessment. *The Physician and Sportsmedicine*, 44(1), 46–52. <http://doi.org/10.1080/00913847.2016.1131107>
- Remvig, L., Jensen, D. V., & Ward, R. C. (2007). Are diagnostic criteria for general joint hypermobility and benign joint hypermobility syndrome based on reproducible and valid tests? A review of the literature.
- Rosen, A. B., Ko, J., & Brown, C. N. (2015). Diagnostic accuracy of instrumented and manual talar tilt tests in chronic ankle instability populations. *Scandinavian Journal of Medicine & Science in Sports*, 25(2), e214–21. <http://doi.org/10.1111/sms.12288>
- Saban, B., & Masharawi, Y. (2016). Three single leg standing tests for clinical assessment of chronic plantar heel pain syndrome: static stance, half-squat and heel rise. *Physiotherapy*. <http://doi.org/10.1016/j.physio.2016.06.003>
- Sacco, I. C. N., Onodera, A. N., Butugan, M. K., Taddei, U. T., Mendes, Y. C., Galhardo, B., et al. (2010). Inter- and intra-tester reliability of clinical measurement to determine medio-lateral patellar position using a pachymeter or visual assessment. *The Knee*, 17(1), 92–95. <http://doi.org/10.1016/j.knee.2009.05.008>
- Sanz, D. R., Lobo, C. C., López, D. L., Morales, C. R., Marín, C. S., & Corbalán, I. S. (2016). Interrater Reliability in the Clinical Evaluation of Myofascial Trigger Points in Three Ankle Muscles. *Journal of Manipulative and Physiological Therapeutics*, 39(9), 623–634. <http://doi.org/10.1016/j.jmpt.2016.09.002>
- Schneider, M., Homonai, R., Moreland, B., Delitto, A. (2007). Interexaminer Reliability of the Prone Leg Length Analysis Procedure. *Journal of Manipulative and Physiological Therapeutics*, 30(7), 514–521. <http://doi.org/10.1016/j.jmpt.2007.07.001>
- Schwieterman, B., Haas, D., Columer, K., Knupp, D., & Cook, C. (2013). Diagnostic accuracy of physical examination tests of the ankle/foot complex: a systematic review. *International Journal of Sports Physical Therapy*, 8(4), 416–426.

- Sebastian, D., Chovvath, R., & Malladi, R. (2014). The sitting active and prone passive lag test: an inter-rater reliability study. *Journal of Bodywork and Movement Therapies*, 18(2), 204–209. <http://doi.org/10.1016/j.jbmt.2013.08.002>
- Senocak, O., Unsal, E., Akalin, E., & Ergör, G. (2003). Interobserver reliability of articular examination in juvenile idiopathic arthritis. *The Turkish Journal of Pediatrics*, 45(1), 29–32.
- Simonsen, O., Jensen, J., Mouritsen, P., & Lauritzen, J. (1984). The accuracy of clinical examination of injury of the knee joint. *Injury*, 16(2), 96–101.
- Sman, A. D., Hiller, C. E., & Refshauge, K. M. (2013). Diagnostic accuracy of clinical tests for diagnosis of ankle syndesmosis injury: a systematic review. *British Journal of Sports Medicine*, 47(10), 620–628. <http://doi.org/10.1136/bjsports-2012-091702>
- Smith, T. O., Clark, A., Neda, S., Arendt, E. A., Post, W. R., Grelsamer, R. P., et al. (2012). The intra- and inter-observer reliability of the physical examination methods used to assess patients with patellofemoral joint instability. *The Knee*, 19(4), 404–410. <http://doi.org/10.1016/j.knee.2011.06.002>
- Snoeker, B. A. M., Lindeboom, R., Zwiderman, A. H., Vincken, P. W. J., Jansen, J. A., & Lucas, C. (2015). Detecting Meniscal Tears in Primary Care: Reproducibility and Accuracy of 2 Weight-Bearing Tests and 1 Non-Weight-Bearing Test. *The Journal of Orthopaedic and Sports Physical Therapy*, 45(9), 693–702. <http://doi.org/10.2519/jospt.2015.5712>
- Solomon, D. J., Simel, D. L., Bates, D. W., Katz, J. N., & Schaffer, J. L. (2001). Does This Patient Have a Torn Meniscus or Ligament of the Knee? *Jama*, 286(13), 1610–1620.
- Stiell, I. G., McKnight, R. D., Greenberg, G. H., Nair, R. C., McDowell, I., & Wallace, G. J. (1992). Interobserver agreement in the examination of acute ankle injury patients. *The American Journal of Emergency Medicine*, 10(1), 14–17.
- Sturgill, L. P. (2009). Interrater Reliability of a Clinical Scale to Assess Knee Joint Effusion. *The Journal of Orthopaedic and Sports Physical Therapy*. <http://doi.org/10.2519/jospt.2009.3143>
- Takeuchi, N., Kuwabara, T., & Usuda, S. (2009). Development and Evaluation of a New Measure for Muscle Tone of Ankle Plantar Flexors: the Ankle Plantar Flexors Tone Scale. *Archives of Physical Medicine and Rehabilitation*, 90(12), 2054–2061. <http://doi.org/10.1016/j.apmr.2009.08.141>
- Thaller, M., & Hughes, T. (2014). Inter-rater agreement of observable and elicitable neurological signs. *Clinical Medicine (London, England)*, 14(3), 264–267. <http://doi.org/10.7861/clinmedicine.14-3-264>
- Trojjan, T. H. (2006). Single leg balance test to identify risk of ankle sprains \* Commentary 1 \* Commentary 2. *British Journal of Sports Medicine*, 40(7), 610–613. <http://doi.org/10.1136/bjsm.2005.024356>

- Ulaşlı, A. M., Yaman, F., Dikici, Ö., Karaman, A., Kaçar, E., & Demirdal, Ü. S. (2014). Accuracy in detecting knee effusion with clinical examination and the effect of effusion, the patient's body mass index, and the clinician's experience. *Clinical Rheumatology*, 33(8), 1139–1143. <http://doi.org/10.1007/s10067-013-2356-6>
- van Dijk, C. N., Lim, L. S., Bossuyt, P. M., & Marti, R. K. (1996a). Physical examination is sufficient for the diagnosis of sprained ankles. *The Journal of Bone and Joint Surgery. British Volume*, 78(6), 958–962.
- van Dijk, C. N., Mol, B. W., Lim, L. S., Marti, R. K., & Bossuyt, P. M. (1996b). Diagnosis of ligament rupture of the ankle joint. Physical examination, arthrography, stress radiography and sonography compared in 160 patients after inversion trauma. *Acta Orthopaedica Scandinavica*, 67(6), 566–570. <http://doi.org/10.3109/17453679608997757>
- Vanpee, G., Hermans, G., Segers, J., & Gosselink, R. (2014). Assessment of Limb Muscle Strength in Critically Ill Patients. *Critical Care Medicine*, 42(3), 701–711. <http://doi.org/10.1097/CCM.0000000000000030>
- Wadey, V. M. R., Mohtadi, N. G. H., Bray, R. C., & Frank, C. B. (2007). Positive predictive value of maximal posterior joint-line tenderness in diagnosing meniscal pathology: a pilot study. *Canadian Journal of Surgery. Journal Canadien De Chirurgie*, 50(2), 96–100.
- Wai, E. K., Eugene K Wai MD, M. F., Kelly Howse BSc, M. D., Howse, K., Pollock, J. W., J Whitcomb Pollock MD, F., et al. (2009). The reliability of determining "leg dominant pain". *The Spine Journal : Official Journal of the North American Spine Society*, 9(6), 447–453. <http://doi.org/10.1016/j.spinee.2008.11.009>
- Walsh, J., & Hall, T. (2009). Reliability, validity and diagnostic accuracy of palpation of the sciatic, tibial and common peroneal nerves in the examination of low back related leg pain. *Manual Therapy*, 14(6), 623–629. <http://doi.org/10.1016/j.math.2008.12.007>
- Watkins, M. A., Riddle, D. L., Lamb, R. L., & Personius, W. J. (1991). Reliability of goniometric measurements and visual estimates of knee range of motion obtained in a clinical setting. *Physical Therapy*, 71(2), 90–96— discussion 96–7.
- Watson, C. J., Leddy, H. M., Dynjan, T. D., & Parham, J. L. (2001). Reliability of the lateral pull test and tilt test to assess patellar alignment in subjects with symptomatic knees: student raters. *The Journal of Orthopaedic and Sports Physical Therapy*, 31(7), 368–374. <http://doi.org/10.2519/jospt.2001.31.7.368>
- Wood, L., Peat, G., Wilkie, R., Hay, E., Thomas, E., & Sim, J. (2006). A study of the noninstrumented physical examination of the knee found high observer variability. *Journal of Clinical Epidemiology*, 59(5), 512–520. <http://doi.org/10.1016/j.jclinepi.2005.11.004>

- Wrobel, J. S., & Armstrong, D. G. (2008). Reliability and validity of current physical examination techniques of the foot and ankle. *Journal of the American Podiatric Medical Association*, 98(3), 197– 206.