

Outpatient Diuretic Intensification: a Simple Prognostic Marker in Cardiac Transthyretin Amyloidosis

Clinical Research in Cardiology

- Online Resource 1 -

Richard J. Nies¹, Svenja Ney¹, Jasper F. Nies², Katharina Seuthe¹, Lukas Klösches¹, Monique Brüwer¹, Stephan Nienaber¹, Sascha Macherey-Meyer¹, Matthieu Schäfer¹, Roman Pfister¹

¹ University of Cologne, Faculty of Medicine and University Hospital Cologne, Clinic III for Internal Medicine, Kerpener Str. 62, 50937 Cologne

² Department of Nephrology, University of Cologne, Kerpener Str. 62, D-50937 Cologne, Germany

Corresponding author:

Dr. med. Richard Nies

University of Cologne, Faculty of Medicine and University Hospital Cologne

Clinic III for Internal Medicine

Kerpener Straße 62


D-50937 Köln, Germany

Phone: +49 221 47876653

Fax: +49 221 47832343

E-mail: richard.nies@uk-koeln.de

Online Resource 1 Flowchart regarding patient selection



AA = amyloid A; AL = amyloid light-chain; ATTR-CM = transthyretin amyloidosis cardiomyopathy; EMB = endomyocardial biopsy.