

Supplementary table Means and standard deviations of the QUIP-RS scales for the whole sample and patients with and without at least subsyndromal disorders.

	<i>Whole sample</i>				<i>Negatives</i>			<i>Positives</i>		
	<i>n</i>	Range	<i>M</i>	<i>SD</i>	<i>n</i>	<i>M₀</i>	<i>SD₀</i>	<i>n</i>	<i>M₁</i>	<i>SD₁</i>
Gambling	153	0-16	0.79	2.21	147	0.59	1.33	6	7.83	5.85
Sex	152	0-12	2.97	3.40	126	2.21	2.77	26	6.69	3.76
Buying	153	0-12	2.05	2.75	141	1.65	2.32	12	6.75	3.13
Eating	147	0-16	2.95	3.47	127	2.24	2.87	20	7.45	3.65
Punding	151	0-16	2.71	3.55	137	2.36	3.33	14	6.14	4.00
Hobbyism	149	0-16	3.61	3.84	133	3.23	3.66	16	6.81	3.87
DDS	153	0-16	2.72	4.06	146	2.50	3.88	6	8.33	4.97
Sum score	138	0-96	17.25	17.10	81	12.21	15.06	57	24.42	17.39

n: subjects with no missing values for the respective scale; *M*, *SD*: mean and standard deviation of the whole sample; *M₀*, *SD₀*: mean and standard deviation for patients without the particular disorder; *M₁*, *SD₁* mean and standard deviation for patients with the particular disorder.