

Article Title: Separation of Stroke from Vestibular Neuritis using the Video Head Impulse Test: Machine Learning Models versus Expert Clinicians

Authors: Chao Wang, Jeevan Sreerama, Benjamin Nham, Nicole Reid, Nese Ozalp, James O. Thomas, Cecilia Cappelen-Smith, Zeljka Calic, Andrew P. Bradshaw, Sally M. Rosengren, Deborah A. Black, Glden Akdal, G. Michael Halmagyi, Mukesh Prasad, Gnana K. Bharathy, Miriam S. Welgampola

Journal: Journal of Neurology

Corresponding Author: Miriam S. Welgampola; Central Clinical School, University of Sydney, Australia; miriam@icn.usyd.edu.au

Supplemental Table 2: Patient Characteristics

Characteristic	First Cohort (Training Set) (n = 252)		Second Cohort (Test Set) (n = 49)		Statistical Testing
	Vestibular Neuritis	Posterior Circulation Stroke	Vestibular Neuritis	Posterior Circulation Stroke	
n (%)	149 (59)	103 (41)	33 (67)	16 (33)	N/A
Sex, female, n (%)	57 (38)	30 (29)	11 (33)	6 (38)	$\chi^2(1) = 0.28, p = 0.60^a$ $\chi^2(1) = 0.46, p = 0.50^b$
Age, years, mean (SD)	56.2 (17.1)	66.7 (11.2)	61.3 (13.1)	68.1 (11.8)	$U = 2145, p = 0.25^a$ $t(117) = 0.45, p = 0.66^b$

^a Comparison between vestibular neuritis patients of first cohort and second cohort

^b Comparison between posterior circulation stroke patients of first cohort and second cohort