

Subcutaneous Natalizumab Administration in Relapsing-Remitting Multiple Sclerosis: Results of EASIER 2 Study

Massimo Filippi¹, Luigi ME Grimaldi^{2,3}, Vincenzo Brescia Morra⁴, Antonella Conte^{5,6}, Cinzia Cordioli⁷, Rocco Totaro⁸, Giacomo Lus⁹, Augusto Rini¹⁰, Fabiana Marinelli¹¹, Paola Valentino¹², Paola Perini¹³, Girolama Alessandra Marfia¹⁴, Mariarosaria Valente¹⁵, Simona Malucchi¹⁶, Chiara Zanetta¹, Lorenzo Pradelli¹⁷, Daria Perini¹⁸, Laura Santoni¹⁸, on behalf of the EASIER 2 study working Group

¹Neurology Unit, Neurorehabilitation Unit, Neurophysiology Service, and Neuroimaging Research Unit, “Vita e Salute” University and IRCCS San Raffaele Scientific Institute, Milan, Italy; ²Neurology Unit, Multiple Sclerosis Center, Fondazione Istituto G. Giglio, Cefalù, Italy; ³UniCamillus–Saint Camillus International University of Health Sciences, Rome, Italy; ⁴Multiple Sclerosis Clinical Care and Research Center, Federico II University - Department of Neuroscience (NSRO), Naples, Italy; ⁵Department of Human Neurosciences, Sapienza, University of Rome, Rome, Italy; ⁶IRCCS Neuromed, Pozzilli, IS, Italy; ⁷Multiple Sclerosis Center, ASST Spedali Civili di Brescia, Montichiari Hospital (Brescia), Italy; ⁸Demyelinating Disease Center, Department of Neurology, San Salvatore Hospital, L’Aquila, Italy; ⁹Multiple Sclerosis Center, UOC II Neurology, Department of Advanced Medical and Surgical Sciences; University of Campania “L. Vanvitelli”, Naples, Italy; ¹⁰Multiple Sclerosis Center, Division of Neurology, A. Perrino Hospital, Brindisi, Italy; ¹¹MS Center - Neurology unit, F. Spaziani Hospital, Frosinone (FR), Italy; ¹²Department of Neurology, Magna Graecia University of Catanzaro, Catanzaro, Italy; ¹³Multiple Sclerosis Centre, University Hospital of Padua, Padua, Italy; ¹⁴Multiple Sclerosis Clinical and Research Unit, Department of Systems Medicine, Tor Vergata University, Rome, Italy; ¹⁵Clinical Neurology, Department of Medicine (DMED) University of Udine, Udine, Italy; ¹⁶SCDO Neurologia - CReSM, AOU San Luigi Gonzaga, Orbassano Torino; ¹⁷AdRes, Torin, Italy; ¹⁸Biogen Italia, Milan, Italy

Corresponding author: Massimo Filippi; filippi.massimo@hsr.it

Average day	N (%)	Mean (SD)	Median (range)
Full cohort			
• Without natalizumab administration	222 (92%)	74.8 (19.8)	80 (5–100)
• With IV administration	213 (88%)	56.9 (24.8)	60 (0–100)
• With SC administration	213 (88%)	70.4 (22.0)	75 (0–100)
Patients with EDSS 0–4			
• Without natalizumab administration	136 (56%)	79.7 (17.3)	80 (5–100)
• With IV administration	131 (54%)	58.8 (22.5)	60 (0–100)
• With SC administration	132 (55%)	70.4 (20.9)	75 (0–100)
Patients with EDSS ≥4			
• Without natalizumab administration	40 (17%)	60.6 (18.8)	60 (10–100)
• With IV administration	38 (16%)	50.9 (25.4)	50 (0–100)
• With SC administration	40 (17%)	61.9 (23.2)	60 (0–100)

Online Resource 4. Quality of life assessment of a day based on the type of administration (scale 0–100)

EDSS Expanded Disability Status Scale, IV intravenous, SC subcutaneous, SD standard deviation