

Subcutaneous Natalizumab Administration in Relapsing-Remitting Multiple Sclerosis: Results of EASIER 2 Study

Massimo Filippi¹, Luigi ME Grimaldi^{2,3}, Vincenzo Brescia Morra⁴, Antonella Conte^{5,6}, Cinzia Cordioli⁷, Rocco Totaro⁸, Giacomo Lus⁹, Augusto Rini¹⁰, Fabiana Marinelli¹¹, Paola Valentino¹², Paola Perini¹³, Girolama Alessandra Marfia¹⁴, Mariarosaria Valente¹⁵, Simona Malucchi¹⁶, Chiara Zanetta¹, Lorenzo Pradelli¹⁷, Daria Perini¹⁸, Laura Santoni¹⁸, on behalf of the EASIER 2 study working Group

¹Neurology Unit, Neurorehabilitation Unit, Neurophysiology Service, and Neuroimaging Research Unit, “Vita e Salute” University and IRCCS San Raffaele Scientific Institute, Milan, Italy; ²Neurology Unit, Multiple Sclerosis Center, Fondazione Istituto G. Giglio, Cefalù, Italy; ³UniCamillus–Saint Camillus International University of Health Sciences, Rome, Italy; ⁴Multiple Sclerosis Clinical Care and Research Center, Federico II University - Department of Neuroscience (NSRO), Naples, Italy; ⁵Department of Human Neurosciences, Sapienza, University of Rome, Rome, Italy; ⁶IRCCS Neuromed, Pozzilli, IS, Italy; ⁷Multiple Sclerosis Center, ASST Spedali Civili di Brescia, Montichiari Hospital (Brescia), Italy; ⁸Demyelinating Disease Center, Department of Neurology, San Salvatore Hospital, L’Aquila, Italy; ⁹Multiple Sclerosis Center, UOC II Neurology, Department of Advanced Medical and Surgical Sciences; University of Campania “L. Vanvitelli”, Naples, Italy; ¹⁰Multiple Sclerosis Center, Division of Neurology, A. Perrino Hospital, Brindisi, Italy; ¹¹MS Center - Neurology unit, F. Spaziani Hospital, Frosinone (FR), Italy; ¹²Department of Neurology, Magna Graecia University of Catanzaro, Catanzaro, Italy; ¹³Multiple Sclerosis Centre, University Hospital of Padua, Padua, Italy; ¹⁴Multiple Sclerosis Clinical and Research Unit, Department of Systems Medicine, Tor Vergata University, Rome, Italy; ¹⁵Clinical Neurology, Department of Medicine (DMED) University of Udine, Udine, Italy; ¹⁶SCDO Neurologia - CReSM, AOU San Luigi Gonzaga, Orbassano Torino; ¹⁷AdRes, Torin, Italy; ¹⁸Biogen Italia, Milan, Italy

Corresponding author: Massimo Filippi; filippi.massimo@hsr.it

Non-healthcare costs	Mean (SD)	Median (range)
Transport cost per patient (€)	31.98 (44.71)	19.15 (0.32–363.23)
• Private vehicle (€)	34.39 (46.09)	20.18 (0.50–363.23)
• Public transport (€)	7.25 (8.23)	6.44 (0.32–31.23)
• Taxi (€)	13.20 (13.30)	13.20 (6.20–20.20)
Cost for formal assistance (€)	1.24 (7.04)	0.00 (0.00–57.28)
• In patients who benefit from it only (€)	33.41 (16.79)	35.80 (7.16–57.28)

Online Resource 6. Non-healthcare costs borne by the patient

SD standard deviation