

Immune function differs among tropical environments but is not downregulated during reproduction in three year-round breeding equatorial lark populations

Submitted to *Oecologia*

Henry K. Ndithia^{1, 2,*}, Kevin D. Matson³, Muchane Muchai^{1, 4}, B. Irene Tieleman²

¹Ornithology Section, Department of Zoology, National Museums of Kenya, P.O. Box 40658 – 00100 GPO, Nairobi, Kenya

²Groningen Institute for Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands;

³Resource Ecology Group, Department of Environmental Sciences, Wageningen University, Droevendaalsesteegh 3a, 6708 PB Wageningen, The Netherlands

⁴Department of Clinical Studies (Wildlife and Conservation), College of Agriculture and Veterinary Sciences, University of Nairobi. Box 30197-00100, Nairobi, Kenya

*Corresponding author:

Email: hndithia@gmail.com

CFSK	0.09 (2.77)
------	-------------

b) T_{max}

	INCKE	CFKE	INCKE	CFKE
CFKE	<0.01 (5.05)			
NBKE	<0.01 (8.57)	0.02 (3.25)		
INCSK			<0.01 (7.64)	
INCNK			<0.01 (7.30)	
CFSK				<0.01 (4.79)
