




Fig. S2: Spatial distribution of zircon grains with heavily pitted surfaces (class 4 of Gärtner et al., 2013) and their possible correlation to glacial deposits and landforms.

Compiled from:

- Andrews, G.D., McGrady, A.T., Brown, S.R., Maynard, S.M., 2019. First description of subglacial megalineations from the late Paleozoic ice age in Southern Africa. *PLoS ONE* 14(1): e0210673.
- Bechstädt, T., Jäger, H., Spence, G., Werner, G., 2009. Late Cryogenian (Neoproterozoic) glacial and post-glacial successions at the southern margin of the Congo Craton, northern Namibia: facies, palaeogeography and hydrocarbon perspective. *Geological Society, London, Special Publications* 326, 255-287.
- Bechstädt, T., Jäger, H., Rittersbacher, A., Schweisfurth, B., Spence, G., Werner, G., Boni, M., 2018. The Cryogenian Ghaub Formation of Namibia - New insights into Neoproterozoic glaciations. *Earth Science Reviews* 177, 678-714.
- Geiger, M., 1999. An Explanation of the Geological Map 1:10 000 of the Namibian borderland along the Orange River at Zwartbas - Warmbad District - Karas Region - Namibia. *Diplomkartierung, Würzburg University*, 80pp.
- Halverson, G.P., Hoffman, P.F., Schrag, D.P., 2002. A major perturbation of carbon cycle before Ghaub glaciation (Neoproterozoic) in Namibia: Prelude to snowball Earth? *Geochemistry Geophysics Geosystems* 3(6), doi: 10.1029/2001GC000244.
- Hoffman, P.F., 2011. Strange bedfellows: glacial diamictite and cap carbonate from the Marinoan (635 Ma) glaciation in Namibia. *Sedimentology* 58, 57-119.
- Hofmann, M., Linnemann, U., Hoffmann, K.-H., Gerdes, A., Eckelmann, K., Gärtner, A., 2013. The Namuskluft and Dreigratberg sections in southern Namibia (Kalahari Craton, Gariep Belt): a geological history of Neoproterozoic rifting and recycling of cratonic crust during the dispersal of Rodinia until the amalgamation of Gondwana. *International Journal of Earth Sciences* 103, 1187-1202.
- Hofmann, M., Linnemann, U., Hoffmann, K.-H., Germs, G., Gerdes, A., Marko, L., Eckelmann, K., Gärtner, A., Krause, R., 2015. The four Neoproterozoic glaciations of southern Namibia and their detrital zircon record: The fingerprints of four crustal growth events during two supercontinent cycles. *Precambrian Research* 259, 176-188.
- MacDonald, F.A., Strauss, J.V., Rose, C.V., Dudás, F.O., Schrag, D., 2010. Stratigraphy of the Port Nolloth Group of Namibia and South Africa and implications for the age of Neoproterozoic iron formations. *American Journal of Science* 310, 862-888.
- Zieger, J., Rothe, J., Hofmann, M., Gärtner, A., Linnemann, U., 2019. The Permian-Carboniferous Dwyka Group of the Aranos Basin (Namibia) – How detrital zircons help understanding sedimentary recycling during a major glaciation. *International Journal of African Earth Sciences* 158, 103555.
- Zimmermann, U., Tait, J., Crowley, Q.G., Pashley, V., Straathof, G., 2011. The Witputs diamictite in southern Namibia and associated rocks: constraints for a global glaciation? *International Journal of Earth Sciences* 100, 511-526.