

Online resource 1: Reference list of included studies

Title: Clinical course of pain and disability following primary lumbar discectomy: systematic review and meta-analysis

Journal: European Spine Journal

Authors: Rushton A, Heneghan NR, Heymans MW, Staal JB, Goodwin PC

Corresponding author: Dr Alison Rushton a.b.rushton@bham.ac.uk

- Abou-Zeid A PJ, Gnanalingham K. Day case lumbar discectomy—Viable option in the UK? British journal of neurosurgery 2014;**28**(3):320-23.
- Alò KM, Wright, R.E., Sutcliffe, J. and Brandt, S.A. Percutaneous lumbar discectomy: clinical response in an initial cohort of fifty consecutive patients with chronic radicular pain. Pain Practice 2004;**4**(1):19-29
- Alò KM, Wright, R.E., Sutcliffe, J. and Brandt, S.A. Percutaneous lumbar discectomy: One-year follow-up in an initial cohort of fifty consecutive patients with chronic radicular pain. . Pain Practice 2005;**5**(2):116-24
- Amoretti N, Huwart L, Marcy PY, Foti P, Hauger O, Boileau P. CT- and fluoroscopy-guided percutaneous discectomy for lumbar radiculopathy related to disc herniation: a comparative prospective study comparing lateral to medial herniated discs. Skeletal Radiol 2013;**42**(1):49-53 doi: 10.1007/s00256-012-1422-5[published Online First: Epub Date]].
- Andrade P, Hoogland G, Teernstra OP, et al. Elevated levels of tumor necrosis factor-alpha and TNFR1 in recurrent herniated lumbar discs correlate with chronicity of postoperative sciatic pain. Spine J 2016;**16**(2):243-51 doi: 10.1016/j.spinee.2015.10.038[published Online First: Epub Date]].
- Arpino L, Iavarone A, Parlato C, Moraci A. Prognostic role of depression after lumbar disc surgery. Neurol Sci 2004;**25**(3):145-7 doi: 10.1007/s10072-004-0248-x[published Online First: Epub Date]].
- Asch HL, Lewis, P.J., Moreland, D.B., Egnatchik, J.G., Young, J.Y., Clabeaux, D.E. and Hyland, A.H., . Prospective multiple outcomes study of outpatient lumbar microdiscectomy: Should 75 to 80% success rates be the norm? Journal of Neurosurgery 2002;**96**(1 SUPPL):34-44
- Atlas SJ, Deyo RA, Keller RB, et al. The Maine Lumbar Spine Study, Part II. 1-year outcomes of surgical and nonsurgical management of sciatica. Spine (Phila Pa 1976) 1996;**21**(15):1777-86
- Atlas SJ, Keller RB, Chang Y, Deyo RA, Singer DE. Surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: five-year outcomes from the Maine Lumbar Spine Study. Spine (Phila Pa 1976) 2001;**26**(10):1179-87
- Atlas SJ, Keller RB, Wu YA, Deyo RA, Singer DE. Long-term outcomes of surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: 10 year results from the maine lumbar spine study. Spine (Phila Pa 1976) 2005;**30**(8):927-35
- Casal-Moro R, Castro-Menendez M, Hernandez-Blanco M, Bravo-Ricoy JA, Jorge-Barreiro FJ. Long-term outcome after microendoscopic discectomy for lumbar disk herniation: a prospective clinical study with a 5-year follow-up. Neurosurgery 2011;**68**(6):1568-75; discussion 75 doi: 10.1227/NEU.0b013e31820cd16a[published Online First: Epub Date]].
- Chaichana KL, Mukherjee D, Adogwa O, Cheng JS, McGirt MJ. Correlation of preoperative depression and somatic perception scales with postoperative disability and quality of life after lumbar discectomy. J Neurosurg Spine 2011;**14**(2):261-7 doi: 10.3171/2010.10.SPINE10190[published Online First: Epub Date]].

- Chen HT, Tsai CH, Chao SC, et al. Endoscopic discectomy of L5-S1 disc herniation via an interlaminar approach: Prospective controlled study under local and general anesthesia. *Surg Neurol Int* 2011;**2**:93 doi: 10.4103/2152-7806.82570[published Online First: Epub Date]].
- Chin KR, Tomlinson DT, Auerbach JD, Shatsky JB, Deirmengian CA. Success of lumbar microdiscectomy in patients with modic changes and low-back pain: a prospective pilot study. *J Spinal Disord Tech* 2008;**21**(2):139-44 doi: 10.1097/BSD.0b013e318093e5dc[published Online First: Epub Date]].
- Choi G, Modi HN, Prada N, et al. Clinical results of XMR-assisted percutaneous transforaminal endoscopic lumbar discectomy. *J Orthop Surg Res* 2013;**8**(1):14 doi: 10.1186/1749-799X-8-14[published Online First: Epub Date]].
- D'Angelo C, Mirijello A, Ferrulli A, et al. Role of trait anxiety in persistent radicular pain after surgery for lumbar disc herniation: a 1-year longitudinal study. *Neurosurgery* 2010;**67**(2):265-71 doi: 10.1227/01.NEU.0000371971.51755.1C[published Online First: Epub Date]].
- Dedering A. Lumbar muscle fatigue and subjective health measurements in patients with lumbar disc herniation 2 years after surgery. *Eur Spine J* 2012;**21**(4):646-54 doi: 10.1007/s00586-011-2109-7[published Online First: Epub Date]].
- Dedering A, Harms-Ringdahl K, Nemeth G. Back extensor muscle fatigue in patients with lumbar disc herniation. Pre-operative and post-operative analysis of electromyography, endurance time and subjective factors. *Eur Spine J* 2006;**15**(5):559-69 doi: 10.1007/s00586-005-0934-2[published Online First: Epub Date]].
- den Boer JJ, Oostendorp RA, Beems T, Munneke M, Evers AW. Continued disability and pain after lumbar disc surgery: the role of cognitive-behavioral factors. *Pain* 2006;**123**(1-2):45-52 doi: 10.1016/j.pain.2006.02.008[published Online First: Epub Date]].
- den Boer JJ, Oostendorp RA, Beems T, Munneke M, Evers AW. Reduced work capacity after lumbar disc surgery: the role of cognitive-behavioral and work-related risk factors. *Pain* 2006;**126**(1-3):72-8 doi: 10.1016/j.pain.2006.06.010[published Online First: Epub Date]].
- Dewing CB, Provencher MT, Riffenburgh RH, Kerr S, Manos RE. The outcomes of lumbar microdiscectomy in a young, active population: correlation by herniation type and level. *Spine (Phila Pa 1976)* 2008;**33**(1):33-8 doi: 10.1097/BRS.0b013e31815e3a42[published Online First: Epub Date]].
- Dubourg G, Rozenberg S, Fautrel B, et al. A pilot study on the recovery from paresis after lumbar disc herniation. *Spine (Phila Pa 1976)* 2002;**27**(13):1426-31; discussion 31
- Eguchi Y, Oikawa Y, Suzuki M, et al. Diffusion tensor imaging of radiculopathy in patients with lumbar disc herniation: preliminary results. *Bone Joint J* 2016;**98-B**(3):387-94 doi: 10.1302/0301-620X.98B3.36036[published Online First: Epub Date]].
- Elkan P, Sten-Linder M, Hedlund R, Willers U, Ponzer S, Gerdhem P. Markers of inflammation and fibrinolysis in relation to outcome after surgery for lumbar disc herniation. A prospective study on 177 patients. *Eur Spine J* 2016;**25**(1):186-91 doi: 10.1007/s00586-015-3998-7[published Online First: Epub Date]].
- Eicker SO, Rhee S, Steiger HJ, Herdmann J, Floeth FW. Transtubular microsurgical approach to treating extraforaminal lumbar disc herniations. *Neurosurg Focus* 2013;**35**(2):E1 doi: 10.3171/2013.4.FOCUS13126[published Online First: Epub Date]].
- Fakouri B, Stovell MG, Allom R. A Comparative Cohort Study of Lumbar Microdiscectomy in Obese and Nonobese Patients. *J Spinal Disord Tech* 2015;**28**(6):E352-7 doi: 10.1097/BSD.0b013e318290bf4a[published Online First: Epub Date]].
- Gadjradj PS, van Tulder MW, Dirven CM, Peul WC, Harhangi BS. Clinical outcomes after percutaneous transforaminal endoscopic discectomy for lumbar disc herniation: a prospective case series. *Neurosurg Focus* 2016;**40**(2):E3 doi: 10.3171/2015.10.FOCUS15484[published Online First: Epub Date]].

- Graver V, Haaland AK, Magnaes B, Loeb M. Seven-year clinical follow-up after lumbar disc surgery: results and predictors of outcome. *Br J Neurosurg* 1999;**13**(2):178-84
- Guilfoyle MR, Ganesan D, Seeley H, Laing RJ. Prospective study of outcomes in lumbar discectomy. *Br J Neurosurg* 2007;**21**(4):389-95 doi: 10.1080/02688690701477310[published Online First: Epub Date]].
- HÄKKINEN A, Ylinen, J., Kautiainen, H., Airaksinen, O., Herno, A. and Kiviranta, I. Does the outcome 2 months after lumbar disc surgery predict the outcome 12 months later? Disability and rehabilitation 2003;**25**(17):968-72
- Häkkinen A, Kautiainen, H., Sintonen, H. and Ylinen, J. Health related quality of life after lumbar disc surgery: a prospective study of 145 patients. . *Disability and rehabilitation* 2005;**27**(3):94-100
- Hussein M. Minimal Incision, Multifidus-sparing Microendoscopic Discectomy Versus Conventional Microdiscectomy for Highly Migrated Intracanal Lumbar Disk Herniations. *J Am Acad Orthop Surg* 2016;**24**(11):805-13 doi: 10.5435/JAAOS-D-15-00588[published Online First: Epub Date]].
- Kahanovitz N, Viola K, Goldstein T, Dawson E. A multicenter analysis of percutaneous discectomy. *Spine (Phila Pa 1976)* 1990;**15**(7):713-5
- Keller RB, Atlas, S.J., Soule, D.N., Singer, D.E. and Deyo, R.A. Relationship between rates and outcomes of operative treatment for lumbar disc herniation and spinal stenosis. *JBJS* 1999;**81**(6):752-62
- Kerr D, Zhao, W. and Lurie, J.D. What are long-term predictors of outcomes for lumbar disc herniation? A randomized and observational study. . *Clinical Orthopaedics and Related Research®* 2015;**473**(6):1920-30
- Lurie JD, Tosteson, T.D., Tosteson, A.N., Zhao, W., Morgan, T.S., Abdu, W.A., Herkowitz, H. and Weinstein, J.N. Surgical versus non-operative treatment for lumbar disc herniation: eight-year results for the Spine Patient Outcomes Research Trial (SPORT). . *Spine* 2014;**39**(1):3
- Kim HS, Ju CI, Kim SW, Kim JG. Endoscopic transforaminal suprapedicular approach in high grade inferior migrated lumbar disc herniation. *J Korean Neurosurg Soc* 2009;**45**(2):67-73 doi: 10.3340/jkns.2009.45.2.67[published Online First: Epub Date]].
- Kleinstueck FS, Fekete, T., Jeszenszky, D., Mannion, A.F., Grob, D., Lattig, F., Mutter, U. and Porchet, F. The outcome of decompression surgery for lumbar herniated disc is influenced by the level of concomitant preoperative low back pain. . *European Spine Journal* 2011;**20**(7):1166-73
- Koen EA, Antonarakos PD, Katranitsa LT, et al. Lumbar Discectomy: Generic Outcome Measures for Specific Outcome Prediction: A Short-term, Observational, Pilot Study. *Clin Spine Surg* 2017;**30**(2):E132-E37 doi: 10.1097/BSD.0000000000000414[published Online First: Epub Date]].
- Kong W, Liao W, Ao J, Cao G, Qin J, Cai Y. The Strategy and Early Clinical Outcome of Percutaneous Full-Endoscopic Interlaminar or Extraforaminal Approach for Treatment of Lumbar Disc Herniation. *Biomed Res Int* 2016;**2016**:4702946 doi: 10.1155/2016/4702946[published Online First: Epub Date]].
- Kuiper JJ, Verbeek, J.H., Frings-Dresen, M.H., Everts, V., van Acker, R.E., Kuiper, S. and Straub, J.P. Exploration of the use of biomarkers to monitor recovery after surgery for lumbar disc herniation: a prospective cohort study. . *Clinical Spine Surgery* 2002;**15**(5):398-403
- Lagerbäck T, Elkan, P., Möller, H., Grauers, A., Diarbakerli, E. and Gerdhem, P. An observational study on the outcome after surgery for lumbar disc herniation in adolescents compared with adults based on the Swedish Spine Register. *The Spine Journal* 2015;**15**(6):1241-47
- Lebow RL, Adogwa O, Parker SL, Sharma A, Cheng J, McGirt MJ. Asymptomatic same-site recurrent disc herniation after lumbar discectomy: results of a prospective longitudinal study with 2-year serial imaging. *Spine (Phila Pa 1976)* 2011;**36**(25):2147-51 doi: 10.1097/BRS.0b013e3182054595[published Online First: Epub Date]].
- Lebow R, Parker SL, Adogwa O, et al. Microdiscectomy improves pain-associated depression, somatic anxiety, and mental well-being in patients with herniated lumbar disc. *Neurosurgery*

- 2012;**70**(2):306-11; discussion 11 doi: 10.1227/NEU.0b013e3182302ec3.[published Online First: Epub Date]].
- Leinonen V, Kankaanpää M, Luukkainen M, et al. Lumbar paraspinal muscle function, perception of lumbar position, and postural control in disc herniation-related back pain. *Spine (Phila Pa 1976)* 2003;**28**(8):842-8
- Li ZZ, Hou, S.X., Shang, W.L., Song, K.R. and Zhao, H.L. The strategy and early clinical outcome of full-endoscopic L5/S1 discectomy through interlaminar approach. *Clinical Neurology and Neurosurgery* 2015;**133**:40-45
- Li ZZ, Hou SX, Shang WL, Song KR, Zhao HL. Modified Percutaneous Lumbar Foraminoplasty and Percutaneous Endoscopic Lumbar Discectomy: Instrument Design, Technique Notes, and 5 Years Follow-up. *Pain Physician* 2017;**20**(1):E85-E98
- Liao Z, Chen W, Wang CH. Transforaminal percutaneous endoscopic surgery for far lateral lumbar intervertebral disk herniation. *Orthopedics* 2014;**37**(8):e717-27 doi: 10.3928/01477447-20140728-58[published Online First: Epub Date]].
- Lima-Ramírez PG B-RD, Viera-Ordóñez JY, Ruíz-Gurría JA, Castillo-Vergara ID, Hernández Carbajal B, León-López DF. PERCUTANEOUS DISCECTOMY: A CURRENT TREATMENT FOR LUMBAR DISC HERNIATION. *Coluna/Columna* 2016;**15**(2):127-30
- Mannion AF, Dvorak, J., Mùntener, M. and Grob, D. A prospective study of the interrelationship between subjective and objective measures of disability before and 2 months after lumbar decompression surgery for disc herniation. *European Spine Journal* 2005;**14**(5):454-65
- McCormick ZL SC, Kotcharian A, Chhatre A, Bender III FJ, Salam A, Menkin S, Kennedy DJ, Plastaras C. Percutaneous lumbar disc decompression using the dekompressor: a prospective long-term outcome study. *Pain Medicine* 2016;**17**(6):1023-30
- Millisdotter M, Stromqvist B, Jonsson B. Proximal neuromuscular impairment in lumbar disc herniation: a prospective controlled study. *Spine (Phila Pa 1976)* 2003;**28**(12):1281-9 doi: 10.1097/01.BRS.0000066458.43883.17[published Online First: Epub Date]].
- Ng LC, Sell P. Predictive value of the duration of sciatica for lumbar discectomy. A prospective cohort study. *J Bone Joint Surg Br* 2004;**86**(4):546-9
- Nie H ZJ, Song Y, Chen G, Wang X, Li Z, Jiang H, Kong Q. Percutaneous Endoscopic Lumbar Discectomy for L5–S1 Disc Herniation Via an Interlaminar Approach Versus a Transforaminal Approach A Prospective Randomized Controlled Study With 2-Year follow Up. *Spine* 2016;**41**(19B):B30–B37
- Ohtori S, Yamashita M, Yamauchi K, et al. Low back pain after lumbar discectomy in patients showing endplate modic type 1 change. *Spine (Phila Pa 1976)* 2010;**35**(13):E596-600 doi: 10.1097/BRS.0b013e3181cd2cb8[published Online First: Epub Date]].
- Okoro TaS, P. A short report comparing outcomes between L4/L5 and L5/S1 single-level discectomy surgery. *Clinical Spine Surgery* 2010;**23**(1):40-42
- Parker SL, Mendenhall SK, Godil SS, et al. Incidence of Low Back Pain After Lumbar Discectomy for Herniated Disc and Its Effect on Patient-reported Outcomes. *Clin Orthop Relat Res* 2015;**473**(6):1988-99 doi: 10.1007/s11999-015-4193-1[published Online First: Epub Date]].
- Pearson A, Lurie, J., Tosteson, T., Zhao, W., Abdu, W., Mirza, S. and Weinstein, J. Who should have Surgery for an Intervertebral Disc Herniation?: Comparative Effectiveness Evidence from SPORT. *Spine* 2012;**37**(2):140-49
- Pearson A, Lurie, J., Tosteson, T., Zhao, W., Abdu, W. and Weinstein, J. Who should have surgery for spinal stenosis?: Treatment effect predictors in SPORT. *Spine* 2012;**37**(21):1791
- Peng CWB, Yeo, W. and Tan, S.B. Percutaneous endoscopic discectomy: clinical results and how it affects the quality of life. *Clinical Spine Surgery* 2010;**23**(6):425-30.
- Porchet F, Bartanusz, V., Kleinstueck, F.S., Lattig, F., Jeszenszky, D., Grob, D. and Mannion, A.F. Microdiscectomy compared with standard discectomy: an old problem revisited with new

- outcome measures within the framework of a spine surgical registry. . *European spine journal* 2009;**18**(3):360-66
- Ramberg N, Sahlstrand T. Early course and long-term follow-up after automated percutaneous lumbar discectomy. *J Spinal Disord* 2001;**14**(6):511-6; discussion 16-7
- Righesso O, Falavigna, A. and Avanzi, O. Correlation between persistent neurological impairment and clinical outcome after microdiscectomy for treatment of lumbar disc herniation. . *Neurosurgery* 2012;**70**(2):390-97
- Ruetten S, Komp M, Merk H, Godolias G. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine (Phila Pa 1976)* 2008;**33**(9):931-9 doi: 10.1097/BRS.0b013e31816c8af7[published Online First: Epub Date]].
- Sahlstrand T, Lonntoft M. A prospective study of preoperative and postoperative sequential magnetic resonance imaging and early clinical outcome in automated percutaneous lumbar discectomy. *J Spinal Disord* 1999;**12**(5):368-74
- Sasani M, Ozer, A.F., Oktenoglu, T., Canbulat, N. and Sarioglu, A.C. Percutaneous endoscopic discectomy for far lateral lumbar disc herniations: prospective study and outcome of 66 patients. *min-Minimally Invasive. Neurosurgery* 2007;**50**(02):91-97
- Schick U, Elhabony R. Prospective comparative study of lumbar sequestrectomy and microdiscectomy. *Minim Invasive Neurosurg* 2009;**52**(4):180-5 doi: 10.1055/s-0029-1237379[published Online First: Epub Date]].
- Schizas C, Tsiridis E, Saksena J. Microendoscopic discectomy compared with standard microsurgical discectomy for treatment of uncontained or large contained disc herniations. *Neurosurgery* 2005;**57**(4 Suppl):357-60; discussion 57-60
- Sharma MK, Chichanovskaia, L.V. and Shlemski, V.A. A comprehensive study of early outcome (at the time of discharge from the hospital) after lumbar discectomy for degenerative spine disease. . *NN Burdenko Journal of Neurosurgery* 2013;**77**(2):30-4
- Silverplats K, Lind B, Zoega B, et al. Clinical factors of importance for outcome after lumbar disc herniation surgery: long-term follow-up. *Eur Spine J* 2010;**19**(9):1459-67 doi: 10.1007/s00586-010-1433-7[published Online First: Epub Date]].
- Solberg T, Johnsen, L.G., Nygaard, Ø.P. and Grotle, M.. . Can we define success criteria for lumbar disc surgery? Estimates for a substantial amount of improvement in core outcome measures. . *Acta orthopaedica* 2013;**84**(2):196-201
- Solberg TK, Nygaard, Ø.P., Sjaavik, K., Hofoss, D. and Ingebrigtsen, T. The risk of “getting worse” after lumbar microdiscectomy. . *European Spine Journal* 2005;**14**(1):49-54
- Sørli A, Moholdt, V., Kvistad, K.A., Nygaard, Ø.P., Ingebrigtsen, T., Iversen, T., Kloster, R. and Solberg, T.K. Modic type I changes and recovery of back pain after lumbar microdiscectomy. . *European Spine Journal* 2012;**21**(11):2252-58
- Sun Z, Li W, Chen Z, et al. Decompression alone versus decompression with instrumented fusion for young patients with single-level lumbar disc herniation: a short-term prospective comparative study. *Chin Med J (Engl)* 2014;**127**(11):2037-42
- Thaler M, Lechner, R., Foedinger, B., Haid, C., Kavakebi, P., Galiano, K. and Obwegeser, A. Driving reaction time before and after surgery for lumbar disc herniation in patients with radiculopathy. . *European Spine Journal* 2012;**21**(11):2259-64
- Than KD CJ, Resnick DK, Shaffrey CI, Ghogawala Z, Mummaneni PV. How to predict return to work after lumbar discectomy: answers from the NeuroPoint-SD registry. . *Journal of Neurosurgery: Spine* 2016;**25**(2):181-86
- Toyone T, Tanaka T, Kato D, Kaneyama R. Low-back pain following surgery for lumbar disc herniation. A prospective study. *J Bone Joint Surg Am* 2004;**86-A**(5):893-6

- Tschugg A, Lener S, Hartmann S, et al. Preoperative sport improves the outcome of lumbar disc surgery: a prospective monocentric cohort study. *Neurosurg Rev* 2017;**40**(4):597-604 doi: 10.1007/s10143-017-0811-6[published Online First: Epub Date]].
- Tullberg T, Grane P, Isacson J. Gadolinium-enhanced magnetic resonance imaging of 36 patients one year after lumbar disc resection. *Spine (Phila Pa 1976)* 1994;**19**(2):176-82
- Veresciagina K, Spakauskas B, Ambrozaitis KV. Clinical outcomes of patients with lumbar disc herniation, selected for one-level open-discectomy and microdiscectomy. *Eur Spine J* 2010;**19**(9):1450-8 doi: 10.1007/s00586-010-1431-9[published Online First: Epub Date]].
- Wallbom AS, Geisser, M.E., Koch, J., Haig, A.J., Guido, C. and Hoff, J.T. Lumbar flexion and dynamic EMG among persons with single level disk herniation pre-and postsurgery with radicular low-back pain. . *American journal of physical medicine & rehabilitation* 2009;**88**(4):302-07
- Weinstein JN, Lurie, J.D., Tosteson, T.D., Tosteson, A.N., Blood, E., Abdu, W.A., Herkowitz, H., Hilibrand, A., Albert, T. and Fischgrund, J. Surgical versus non-operative treatment for lumbar disc herniation: four-year results for the Spine Patient Outcomes Research Trial (SPORT). *Spine* 2008;**33**(25):2789
- Xu H, Liu, X., Liu, G., Zhao, J., Fu, Q. and Xu, B. Learning curve of full-endoscopic technique through interlaminar approach for L5/S1 disk herniations. . *Cell biochemistry and biophysics* 2014;**70**(2):1069-74
- Yadav YR, Parihar V, Namdev H, Agarwal M, Bhatele PR. Endoscopic interlaminar management of lumbar disc disease. *J Neurol Surg A Cent Eur Neurosurg* 2013;**74**(2):77-81 doi: 10.1055/s-0032-1333127[published Online First: Epub Date]].
- Yeom KS, Choi YS. Full endoscopic contralateral transforaminal discectomy for distally migrated lumbar disc herniation. *J Orthop Sci* 2011;**16**(3):263-9 doi: 10.1007/s00776-011-0048-0[published Online First: Epub Date]].
- Ying J, Huang K, Zhu M, et al. The Effect and Feasibility Study of Transforaminal Percutaneous Endoscopic Lumbar Discectomy Via Superior Border of Inferior Pedicle Approach for Down-Migrated Intracanal Disc Herniations. *Medicine (Baltimore)* 2016;**95**(8):e2899 doi: 10.1097/MD.0000000000002899[published Online First: Epub Date]].
- Yoon SM, Ahn, S.S., Kim, K.H., Kim, Y.D., Cho, J.H. and Kim, D.H. Comparative study of the outcomes of percutaneous endoscopic lumbar discectomy and microscopic lumbar discectomy using the tubular retractor system based on the VAS, ODI, and SF-36. . *Korean Journal of Spine* 2012;**9**(3):215