

ONLINE RESOURCE 2: references of excluded studies (n=31)

Title: Clinical course of pain and disability following primary lumbar discectomy: systematic review and meta-analysis

Journal: European Spine Journal

Authors: Rushton A, Heneghan NR, Heymans MW, Staal JB, Goodwin PC

Corresponding author: Dr Alison Rushton a.b.rushton@bham.ac.uk

Authors responded – no raw data available

Lonne G, Solberg TK, Sjaavik K, Nygaard OP. Recovery of muscle strength after microdiscectomy for lumbar disc herniation: a prospective cohort study with 1-year follow-up. Eur Spine J 2012;**21**(4):655-9 doi: 10.1007/s00586-011-2122-x[published Online First: Epub Date]].

Unable to contact authors

Amoretti N, David P, Grimaud A, et al. Clinical follow-up of 50 patients treated by percutaneous lumbar discectomy. Clin Imaging 2006;**30**(4):242-4 doi: 10.1016/j.clinimag.2006.01.019[published Online First: Epub Date]].

Asher R MA, Weiner J, Fessler RG. The relationship between preoperative general mental health and postoperative quality of life in minimally invasive lumbar spine surgery. Neurosurgery 2015;**76**:672-79

Azimi P BE, Montazeri A. Predictive Score Card in Lumbar Disc Herniation: Is It Reflective of Patient Surgical Success after Discectomy? Plos One 2016;**11**(4)

Belykh E, Giers MB, Preul MC, Theodore N, Byvaltsev V. Prospective Comparison of Microsurgical, Tubular-Based Endoscopic, and Endoscopically Assisted Discectomies: Clinical Effectiveness and Complications in Railway Workers. World Neurosurg 2016;**90**:273-80 doi: 10.1016/j.wneu.2016.02.047[published Online First: Epub Date]].

El-Badry A BT. Results of Lumbar Discectomy in Patients with Recurrent Disc Herniation Compared to Patients with Primary Disc Herniation. Egypt J Neurol Psychiat Neurosurg 2012;**49**:335-40

Eyenga VC, Ngowe NN, Minkande JZ, Ngah JE. Kinetics of regression of sciatica and pain in the low back after lumbar macrodiscectomy in human immunodeficiency virus carriers. Spine (Phila Pa 1976) 2008;**33**(13):E411-3 doi: 10.1097/BRS.0b013e318175c32b[published Online First: Epub Date]].

Fisher C, Noonan V, Bishop P, et al. Outcome evaluation of the operative management of lumbar disc herniation causing sciatica. J Neurosurg 2004;**100**(4 Suppl Spine):317-24

Gautschi OP SN, Joswigc H, Corniolaa MV, Schallera K, Hildebrandtc G, Stienena MN. Influence of age on pain intensity, functional impairment and health-related quality of life before and after surgery for lumbar degenerative disc disease. Clinical Neurology and Neurosurgery 2016;**150**:33–39

- Gautschi OP, Joswig H, Corniola MV, et al. Pre- and postoperative correlation of patient-reported outcome measures with standardized Timed Up and Go (TUG) test results in lumbar degenerative disc disease. *Acta Neurochir (Wien)* 2016;**158**(10):1875-81 doi: 10.1007/s00701-016-2899-9[published Online First: Epub Date]].
- Hegarty D, Shorten G. Multivariate prognostic modeling of persistent pain following lumbar discectomy. *Pain Physician* 2012;**15**(5):421-34
- Hickey OT, Burke SM, Hafeez P, et al. Determinants of outcome for patients undergoing lumbar discectomy: a pilot study. *Eur J Anaesthesiol* 2010;**27**(8):696-701 doi: 10.1097/EJA.0b013e32833b46dc[published Online First: Epub Date]].
- Kahanovitz N, Viola K, Goldstein T, Dawson E. A multicenter analysis of percutaneous discectomy. *Spine (Phila Pa 1976)* 1990;**15**(7):713-5
- Kapetanakis S GG, Thomaidis T, Charitoudis G, Kazakos K. Comparison of quality of life between men and women who underwent Transforaminal Percutaneous Endoscopic Discectomy for lumbar disc herniation. *Endoscopic Minimally Invasive Surgery* 2017;**11**(4):218 - 28
- Khan Z SS, Ali M. Surgical outcome after microdiscectomy for lumbar disc herniation. *J. Med. Sci. (Peshawar)* 2013;**21**(2):74-76
- K K. Closed Drainage versus Non-Drainage for Single-Level Lumbar Disc Surgery: Relationship between Epidural Hematoma and Fibrosis. *Asian Spine J* 2016;**10**(6):1072-78
- Lundin A MA, Nilsson O. A stiff and straight back preoperatively is associated with a good outcome 2 years after lumbar disc surgery A prospective study of 80 patients. *Acta Orthopaedica* 2009;**80**(5):573-78
- Pneumatics SG CA, Hipp J, Chatziioannou SN. Prediction of successful discectomy using MRI quantitation of dural sac and herniated disc dimensions. *Int J Clin Pract* 2010;**64**(1):13-18
- Quon JA, Sobolev BG, Levy AR, et al. The effect of waiting time on pain intensity after elective surgical lumbar discectomy. *Spine J* 2013;**13**(12):1736-48 doi: 10.1016/j.spinee.2013.05.038[published Online First: Epub Date]].
- Rothoerl DR WC, Holzschuh M, Schlaier J. . Are there differences in the symptoms, signs and outcome after lumbar disc surgery in the elderly compared with younger patients? *British journal of neurosurgery* 1998;**12**(3):250-53
- Shah M KM, Safi T, Aman A, Ahmad A, Shah MA, Ahmed F. Effectiveness of disc excision in the treatment of herniated lumbar intervertebral disc. *J. Med. Sci. (Peshawar)* 2015;**23**(3):153-57
- Sørensen LV MO, Skovlund O. . A prospective study of the importance of psychological and social factors for the outcome after surgery in patients with slipped lumbar disk operated upon for the first time. *Acta neurochirurgica* 1987;**88**(3-4):119-25
- LV. S. Preoperative psychological testing with the MMPI at first operation for prolapsed lumbar disc. Five-year follow up. . *Danish medical bulletin* 1992;**39**(2):186-90
- Tafazal SI, Sell PJ. Outcome scores in spinal surgery quantified: excellent, good, fair and poor in terms of patient-completed tools. *Eur Spine J* 2006;**15**(11):1653-60 doi: 10.1007/s00586-005-0028-1[published Online First: Epub Date]].
- Thomas KC, Fisher CG, Boyd M, Bishop P, Wing P, Dvorak MF. Outcome evaluation of surgical and nonsurgical management of lumbar disc protrusion causing radiculopathy. *Spine (Phila Pa 1976)* 2007;**32**(13):1414-22 doi: 10.1097/BRS.0b013e318060a5d1[published Online First: Epub Date]].
- Tschugg A LS, Hartmann S, Neururer S, Wildauer M, Thomé C, Löscher WN. . Improvement of sensory function after sequestrectomy for lumbar disc herniation: a prospective clinical study using quantitative sensory testing. . *European Spine Journal* 2016;**25**(11):3543-9
- Wang B, Lu G, Patel AA, Ren P, Cheng I. An evaluation of the learning curve for a complex surgical technique: the full endoscopic interlaminar approach for lumbar disc herniations. *Spine J* 2011;**11**(2):122-30 doi: 10.1016/j.spinee.2010.12.006[published Online First: Epub Date]].

Weir B. Prospective study of 100 lumbosacral discectomies. *Journal of Neurosurgery* 1979;**50**:283-89

Whitmore RG, Curran JN, Ali ZS, et al. Predictive value of 3-month lumbar discectomy outcomes in the NeuroPoint-SD Registry. *J Neurosurg Spine* 2015;**23**(4):459-66 doi: 10.3171/2015.1.SPINE14890[published Online First: Epub Date] |.

Woertgen C RR, Holzschuh M, Breme K, Brawanski A. . Are prognostic factors still what they are expected to be after long-term follow-up? *Journal of spinal disorders* 1998;**11**(5):395-9

Woertgen C, Rothoerl RD, Breme K, Altmeyen J, Holzschuh M, Brawanski A. Variability of outcome after lumbar disc surgery. *Spine (Phila Pa 1976)* 1999;**24**(8):807-11