

Genome size variation in *Deschampsia cespitosa* sensu lato (Poaceae) in Eurasia

Plant Systematics and Evolution

Josef Greimler^{1*}, Eva M. Temsch¹, Zhiqing Xue¹, Hanna Weiss-Schneeweiss¹, Polina Volkova², Markus Peintinger³, Pawel Wasowicz⁴, Hui Shang⁵, Ivan Schanzer⁶, Jorge O. Chiapella⁷

¹ Department für Botanik und Biodiversitätsforschung, Universität Wien, Rennweg 14, 1030 Wien, Austria

² Papanin Institute for Biology of Inland Waters RAS, Borok, Yaroslavl Region, 152742, Russia.

³ WSL Swiss Federal Research Institute, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

⁴ Icelandic Institute of Natural History (Náttúrufræðistofnun Íslands), Borgir við Nordurslóð, 600 Akureyri, Iceland


⁵ Shanghai Botanical Garden, 997 Longwu Rd, Xuhui District, Shanghai, China

⁶ Tsitsin Main Botanical Garden of Russian Academy of Sciences, Botanicheskaya Ul. 4, 127276 Moscow, Russia.

⁷ Instituto de Investigaciones en Biodiversidad y Medioambiente (INIBIOMA-CONICET-Universidad Nacional del Comahue), Quintral 1250, R8400FRF Bariloche, Río Negro, Argentina.

* Author for correspondence: Josef Greimler, josef.greimler@univie.ac.at

Online Resource 4. Regional variation in monoploid genome size of *Deschampsia cespitosa*



Variation in monoploid genome size (Cx, pg) from western Europe to eastern Asia along longitude classes in steps of 10 degrees. Missing classes because of missing samples from some regions.