

Article title: Spatial genetic structure in the Madeiran endemic *Dactylorhiza foliosa* (Orchidaceae)

Journal name: Plant Systematics and Evolution

Author name: Mikael Hedrén

Affiliation: Dept of Biology, Lund University, Sweden

e-mail: mikael.hedren [at] biol.lu.se

Online resource 1 Sampling locations of *Dactylorhiza foliosa*

Locality	Latitude	Longitude	approximate elevation (m asl)	sample size	Distance (m) between adjacent samples (mean, standard deviation, range)
Ribeiro Frio	32°44'05''N	16°53'04''W	900 – 950	31	26.5 ± 35.5; 1 – 160
Encumeada North	32°45'18''N	17°00'53''W	950 – 1000	40	12.8 ± 16.9; 1 – 71
Encumeada West	32°44'46''N	17°02'27''W	1190	12	66.0 ± 166.8; 2 – 564
Encumeada West Central	32°44'59''N	17°01'52''W	1120	3	42.0 ± 49.5; 7 – 77
Portela	32°44'31''N	16°51'13''W	850	25	151.2 ± 256.5; 2 – 1051
Rabaçal	32°45'42''N	17°08'05''W	970 – 990	7	106.7 ± 120.4; 2 – 489
Pico das Pedras	32°46'32''N	16°53'35''W	850	16	68.0 ± 143.7; 2 – 540
Queimados	32°46'56''N	16°55'36''W	850 – 900	39	91.2 ± 127.7; 1 – 495