

ONLINE RESOURCE 1 - Supplementary information

Title article: Awake craniotomy does not lead to increased psychological complaints

Journal: Acta neurochirurgica

Authors: I.M.C. Huenges Wajer^{1,2}, PhD – J. Kal^{1,2}, MSc – P.A.J.T. Robe¹, MD, PhD – M. J. E. van Zandvoort^{1,2}, PhD – C. Ruis^{1,2}, PhD

¹ *Department of Neurology and Neurosurgery, Rudolf Magnus Institute for Neuroscience, University Medical Center Utrecht, Utrecht, The Netherlands*

² *Experimental Psychology, Helmholtz Institute, Utrecht University, Utrecht, The Netherlands*

Corresponding author: I.M.C. Huenges Wajer (I.M.C.HuengesWajer@umcutrecht.nl)

Table 1: Kendall's tau correlation between sex, age and complaints of anxiety, depression and PTSD

	Anxiety						Depression						PTSD			
	T0		T1		T2		T0		T1		T2		T1		T2	
	r_τ	p	r_τ	p	r_τ	p	r_τ	p	r_τ	p	r_τ	p	r_τ	p	r_τ	p
Sex	.321	.012*	.210	.092	.194	.109	.327	.011*	.185	.141	.124	.306	.131	.292	.048	.689
Age	-.144	.180	-.066	.530	.007	.948	-.008	.939	.091	.388	.197	.053	.016	.876	-.054	.591

T0 = pre-operative; T1 = 4 weeks after surgery; T2 = 3 months after surgery

** $p < .05$*