

**Soft tissue changes in upper incisors from tooth movements using a new measurement method
based on digital scans superimposition in adult patients.**

Luis Sastre-Buades; Verónica García-Sanz; Beatriz Tarazona-Álvarez; Natalia Zamora-Martínez;
Sara Camañes-Gonzalvo; José María Montiel-Company; Vanessa Paredes-Gallardo; Carlos Bellot-
Arcís.

Online Resource

Clinical Oral Investigations

Correspondence to: Verónica García-Sanz, Orthodontics Teaching Unit, Department of Stomatology
(University of Valencia), Gascó Oliag, 1. 46010. Valencia, Spain.

e-mail: veronica.garcia-sanz@uv.es

Fig. S1 A. Pre-treatment sagittal section (2.1); B. Post-treatment sagittal section (2.1). C.
Superposition of the pre- and post-treatment sagittal slices of the 2.1




Fig. S2 A. Protrusive/retrusive movement measurement (mm); B. Retroclination/proclination measurement ($^{\circ}$); C. Tip movement measurement ($^{\circ}$); D. Rotation movement measurement ($^{\circ}$); E. Intrusion/extrusion movement measurement (mm).




Fig. S3 Correlation between free gingival gain/loss (+/-) and proclination movement.




Figure S4. Correlation between free gingival gain/loss (+/-) and retroclination movement.

