

Clinical score for early diagnosis of myotonic dystrophy type 2

Neurological Sciences

Vukan Ivanovic¹, Stojan Peric^{1*}, Jovan Pesovic², Radoje Tubic³, Ivo Bozovic¹, Ivana Petrovic⁴,
Dusanka Savic-Pavicevic², Giovanni Meola⁵, Vidosava Rakocevic-Stojanovic¹

¹University of Belgrade – Faculty of Medicine, University Clinical Center of Serbia – Neurology Clinic, Belgrade, Serbia

²University of Belgrade – Faculty of Biology, Center for Human Molecular Genetics, Belgrade, Serbia

³Institute of Oncology and Radiology of Serbia, Belgrade, Serbia

⁴University of Belgrade – Faculty of Medicine, University Clinical Center of Serbia – Endocrinology Clinic, Belgrade, Serbia


⁵ Department of Neurorehabilitation Sciences - Casa di Cura del Policlinico, Department of Biomedical Sciences for Health, University of Milan, Italy

* Corresponding Author:

Stojan Peric, MD, PhD, neurologist

Email. stojanperic@gmail.com

Online Resource 1. ROC curve analysis of DM2-EDS in our cohort of patients



DM2-EDS – DM2 Early Diagnosis Score; ROC – Receiver operating characteristic