

Online Resource 2 Stool culture

Group D n = 28		Group N n = 11	
Bacterium	Number (%)	Bacterium	Number (%)
<i>Enterococcus</i>	15 (53.4%)	<i>Enterococcus</i>	6 (54.5%)
<i>Escherichia coli</i>	9 (32.1%)	<i>Escherichia coli</i>	4 (36.4%)
<i>Klebsiella pneumoniae</i>	7 (25.0%)	<i>Culture negative</i>	2 (18.2%)
<i>Candida</i>	5 (17.9%)	<i>Klebsiella pneumoniae</i>	1 (9.1%)
<i>Pseudomonas</i>	5 (17.9%)	<i>Candida</i>	1 (9.1%)
<i>Citrobacter</i>	2 (7.1%)	<i>Citrobacter</i>	1 (9.1%)
<i>Klebsiella oxytoca</i>	2 (7.1%)	<i>Serratia</i>	1 (9.1%)
<i>Enterobacter cloacae</i>	2 (7.1%)	<i>Aeromonas</i>	1 (9.1%)
<i>Clostridium difficile</i>	2 (7.1%)		
<i>Stenotrophomonas maltophilia</i>	2 (7.1%)		
<i>Lactobacillus</i>	1 (3.6%)		
<i>Proteus</i>	1 (3.6%)		
<i>Serratia</i>	1 (3.6%)		
<i>Streptococcus</i>	1 (3.6%)		
<i>Acinetobacter</i>	1 (3.6%)		
<i>Raoultella ornithinolytica</i>	1 (3.6%)		
<i>Morganella morganii</i>	1 (3.6%)		

Clinical impact of diarrhea during enteral feeding after esophagectomy

Ryoma Haneda, MD¹, Yoshihiro Hiramatsu, MD, Ph.D^{1,2}, Sanshiro Kawata, MD, Ph.D¹,
Wataru Soneda, MD¹, Eisuke Booka, MD, Ph.D¹, Tomohiro Murakami, MD, Ph.D¹,
Tomohiro Matsumoto, MD, Ph.D¹, Yoshifumi Morita, MD, Ph.D¹, Hirotoshi Kikuchi,
MD, Ph.D¹, and Hiroya Takeuchi, MD, Ph.D¹

1. Department of Surgery, Hamamatsu University School of Medicine, Hamamatsu,
Shizuoka, Japan

2. Department of Perioperative Functioning Care and Support, Hamamatsu University
School of Medicine, Hamamatsu, Shizuoka, Japan

Corresponding author: Yoshihiro Hiramatsu, MD, Ph.D.

Department of Perioperative Functioning Care and Support, Hamamatsu University
School of Medicine

1-20-1 Handayama, Higashi-ku, Hamamatsu, Shizuoka 431-3192, Japan

E-mail: hiramatu@hama-med.ac.jp

Phone: +81-53-435-2427; Fax: +81-53-435-2423