

Online Resource 4 Independent relevant factors of postoperative high NLR

	Univariate analysis			Multivariate analysis		
	OR	P	95% CI	OR	P	95% CI
Age, ≥ 70	1.111	0.745	0.588—2.101	1.090	0.800	0.559—2.126
Transthoracic approach, Thoracotomy	0.864	0.703	0.409—1.828	0.820	0.626	0.368—1.825
Abdominal approach, Laparotomy	2.314	0.022	1.126—4.753	2.303	0.029	1.090—4.864
Postoperative infectious complications, +	1.250	0.504	0.649—2.409	1.004	0.991	0.498—2.025
Diarrhea during EF	2.446	0.015	1.193—5.015	2.577	0.014	1.213—5.472

NLR, neutrophil-to-lymphocyte ratio; *OR*, odds ratio; *CI*, confidence interval; *EF*, enteral feeding

Clinical impact of diarrhea during enteral feeding after esophagectomy

Ryoma Haneda, MD¹, Yoshihiro Hiramatsu, MD, Ph.D^{1,2}, Sanshiro Kawata, MD, Ph.D¹,
Wataru Soneda, MD¹, Eisuke Booka, MD, Ph.D¹, Tomohiro Murakami, MD, Ph.D¹,
Tomohiro Matsumoto, MD, Ph.D¹, Yoshifumi Morita, MD, Ph.D¹, Hirotoshi Kikuchi,
MD, Ph.D¹, and Hiroya Takeuchi, MD, Ph.D¹

1. Department of Surgery, Hamamatsu University School of Medicine, Hamamatsu,
Shizuoka, Japan

2. Department of Perioperative Functioning Care and Support, Hamamatsu University
School of Medicine, Hamamatsu, Shizuoka, Japan

Corresponding author: Yoshihiro Hiramatsu, MD, Ph.D.

Department of Perioperative Functioning Care and Support, Hamamatsu University
School of Medicine

1-20-1 Handayama, Higashi-ku, Hamamatsu, Shizuoka 431-3192, Japan

E-mail: hiramatu@hama-med.ac.jp

Phone: +81-53-435-2427; Fax: +81-53-435-2423