




Online Resource 5. Kaplan–Meier analysis based on the postoperative nutritional and immunological status. **a.** Comparison of overall survival between low- and high-CRP

groups. **b.** Comparison of recurrence-free survival between low- and high-CRP groups. **c.** Comparison of overall survival between low- and high-PNI groups. **d.** Comparison of recurrence-free survival between low- and high-PNI groups. **e.** Comparison of overall survival between low- and high-NLR groups. **f.** Comparison of recurrence-free survival between low- and high-NLR groups.

Clinical impact of diarrhea during enteral feeding after esophagectomy

Ryoma Haneda, MD¹, Yoshihiro Hiramatsu, MD, Ph.D^{1,2}, Sanshiro Kawata, MD, Ph.D¹, Wataru Soneda, MD¹, Eisuke Booka, MD, Ph.D¹, Tomohiro Murakami, MD, Ph.D¹, Tomohiro Matsumoto, MD, Ph.D¹, Yoshifumi Morita, MD, Ph.D¹, Hirotoshi Kikuchi, MD, Ph.D¹, and Hiroya Takeuchi, MD, Ph.D¹

1. Department of Surgery, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan
2. Department of Perioperative Functioning Care and Support, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan

Corresponding author: Yoshihiro Hiramatsu, MD, Ph.D.

Department of Perioperative Functioning Care and Support, Hamamatsu University School of Medicine

1-20-1 Handayama, Higashi-ku, Hamamatsu, Shizuoka 431-3192, Japan

E-mail: hiramatu@hama-med.ac.jp

Phone: +81-53-435-2427; Fax: +81-53-435-2423