

Electronic supplementary material

Efficacy and safety of esaxerenone (CS-3150) in Japanese patients with type 2 diabetes and macroalbuminuria: a multicenter, single-arm, open-label phase III study

Clinical and Experimental Nephrology

Sadayoshi Ito, Naoki Kashihara, Kenichi Shikata, Masaomi Nangaku, Takashi Wada,
Yasuyuki Okuda, Tomoko Sawanobori

Corresponding author:

Sadayoshi Ito, MD, PhD

Division of Nephrology, Endocrinology and Vascular Medicine, Department of Medicine,
Tohoku University School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8575,
Japan

E-mail: db554@med.tohoku.ac.jp

Online Resource 3 Study schedule of assessments

Assessments	Run-in period			Treatment period													Follow-up period
Timing of visit (weeks)	Start	2	3 ^a	Start	2	4	6	8	12	16	20	24	28 (EOT)	2 weeks after dose increase	After visits where serum K ⁺ ≥5.5 mEq/L	Treatment discontinuation	4 weeks after EOT
Urine sampling UACR	X	X	X ^a			X		X	X	X	X	X	X			X	X
Blood sampling																	
Hematology		X				X			X				X			X	
Biochemistry		X				X			X				X			X	
Potassium		X			X	X	X	X	X	X	X	X	X	X	X ^b	X	X
Creatinine		X			X	X	X	X	X	X	X	X	X	X	X ^b	X	X
Vital signs																	
BP, pulse	X	X			X	X	X	X	X	X	X	X	X	X		X	X
Adverse events				X	X	X	X	X	X	X	X	X	X	X	X	X	X

^aAssessed at week 3 if urinary albumin-to-creatinine ratio criteria were not met at week 2.

^bSerum potassium and creatinine measurement only.

BP blood pressure; *EOT* end of treatment; *UACR* urinary albumin-to-creatinine ratio.

