

Supplementary Information

Online Resource TEAEs occurring in $\geq 5\%$ of the total patients (SAS)

Adverse event by SOC	<50 kg group (N=22)	≥ 50 kg group (N=5)	Total (N=27)
Any TEAEs	19 (86.4%)	5 (100.0%)	24 (88.9%)
Gastrointestinal disorders			
Nasopharyngitis	9 (40.9%)	2 (40.0)	11 (40.7%)
Constipation	3 (13.6%)	1 (20.0)	4 (14.8%)
Influenza	3 (13.6%)	1 (20.0)	4 (14.8%)
Gastroenteritis	4 (18.2%)	0	4 (14.8%)
Epistaxis	2 (9.1%)	1 (20.0%)	3 (11.1%)
Abdominal pain	3 (13.6%)	0	3 (11.1%)
Pyrexia	3 (13.6%)	0	3 (11.1%)
Otitis media	3 (13.6%)	0	3 (11.1%)
Dizziness	3 (13.6%)	0	3 (11.1%)
Stomatitis	2 (9.1%)	0	2 (7.4%)
Sinusitis	2 (9.1%)	0	2 (7.4%)
Upper respiratory tract infection	2 (9.1%)	0	2 (7.4%)
Contusion	2 (9.1%)	0	2 (7.4%)
Ligament sprain	2 (9.1%)	0	2 (7.4%)
Serum creatinine increased	2 (9.1%)	0	2 (7.4%)
Renal impairment	1 (4.5%)	1 (20.0%)	2 (7.4%)
Asthma	2 (9.1%)	0	2 (7.4%)
Cough	2 (9.1%)	0	2 (7.4%)
Dermatitis atopic	2 (9.1%)	0	2 (7.4%)

Values are n (%).

SAS safety analysis set, **SOC** System Organ Class, **TEAE** treatment-emergent adverse event