

		baseline			6W			12W			6M			1Y			2Y		
		mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE
UNa (mEq/gCr)	Standard (31)	154.0	93.5	16.8	137.4	62.4	11.2	157.0	85.7	15.4	146.1	76.3	13.7	82.3 (4)†	44.2	22.1	207.3 (3)†	69.8	40.3
	<i>p</i> value*, vs. baseline	-			0.448			0.411			0.158			0.0077*			0.922		
	SB (31)	153.6	79.1	14.2	161.7	81.3	14.6	193.5	123.6	22.2	177.7	106.3	19.1	112.4 (7)†	49.7	18.8	132.5 (5)†	943.8	422.1
	<i>p</i> value*, vs. baseline	-			0.808			0.58			0.76			0.042			0.196		
	PCSC (32)	131.7	107.1	17.5	134.4	72.1	13.2	157.4	72.6	13.7	164.5	93.7	17.4	146.1 (10)†	61.3	19.4	177.5 (9)†	117.3	39.1
	<i>p</i> value*, vs. baseline	-			0.881			0.163			0.139			0.506			0.305		
	<i>p</i> value among 3 groups**	-			0.319			0.338			0.382			0.908			0.718		
	<i>p</i> value by ANOVA*	0.0115*																	
UK (mEq/gCr)		baseline			6W			12W			6M			1Y			2Y		
		mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE
	Standard (31)	53.7	31.8	5.71	51.0	25.1	4.51	53.1	24.4	4.39	51.1	23.1	4.14	40.6 (4)†	11.6	5.79	49.6 (3)†	17.5	10.1
	<i>p</i> value*, vs. baseline	-			0.202			0.303			0.014			0.531			0.318		
	SB (31)	45.7	18.7	3.36	46.6	21.8	3.92	49.0	26.5	4.76	48.3	29.0	5.20	25.2 (7)†	15.4	5.82	48.0 (5)†	10.1	4.48
	<i>p</i> value*, vs. baseline	-			0.530			0.663			0.170			0.142			0.548		
	PCSC (32)	52.8	28.1	4.97	56.1	26.6	4.71	59.1	38.9	6.87	63.4	41.6	7.35	45.0 (10)†	13.7	4.33	58.4 (9)†	1.94	6.48
<i>p</i> value*, vs. baseline	-			0.198			0.284			0.170			0.142			0.548			
	<i>p</i> value among 3 groups**	-			0.314			0.372			0.117			0.124			0.06		
	<i>p</i> value by ANOVA*	0.1627																	
UNa/K ratio		baseline			6W			12W			6M			1Y			2Y		
		mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE	mean	SD	SE
	Standard (31)	3.17	1.89	0.34	2.90	1.22	0.22	3.16	1.69	0.21	2.95	1.17	0.21	2.05 (4)†	1.10	0.55	2.21 (3)†	0.19	0.11
	<i>p</i> value*, vs. baseline	-			0.740			0.272			0.366			0.066			0.057		
	SB (31)	3.46	1.74	0.313	3.72	1.89	0.34	4.44	2.56	0.46	4.09	2.51	0.45	3.53 (7)†	0.77	0.29	2.86 (5)†	1.32	0.59
	<i>p</i> value*, vs. baseline	-			0.265			0.297			0.374			0.102			0.738		
	PCSC (32)	2.57	1.47	0.26	2.37	0.90	0.16	3.06	1.75	0.31	2.7	1.41	0.25	2.95 (10)†	1.87	0.59	3.10 (9)†	1.80	0.6
<i>p</i> value*, vs. baseline	-			0.562			0.135			0.531			0.054			0.794			
	<i>p</i> value among 3 groups**	-			0.003**			0.031			0.035			0.630			0.003*		
	<i>p</i> value by ANOVA*	0.0951																	

Supplemental Table 1. Additional data of urinary excretion of Na, and K, and urinary Na/K ratio. Values were described mean, standard deviation (SD) and standard error (SE). The significance of comparison vs. baseline was defined as **p*<0.05, and the significance of comparison among the three groups and ANOVA were ***p*<0.0167.

(N) †; Number of patients at 1Y or 2Y for the long-term study.