

Supporting material for: Vindstad, O.P.L., Schott, T., Hagen, S.B., Jepsen, J.U., Kapari, L. & Ims, R.A. "How rapidly do invasive birch forest geometrids recruit larval parasitoids? – Insights from comparison with a sympatric native geometrid" (Biological invasions). Corresponding author: Ole Petter Laksforsmo Vindstad, Department of Arctic and Marine Biology, University of Tromsø, Norway, e-mail: ole.p.vindstad@uit.no

Appendix 1. Outdated parasitoid species names originally given in the cited papers and corresponding currently valid names.

Parasitoid family	Outdated species name	Valid species name
Ichneumonidae	<i>Lissonota femorata</i> (Hassell 1966)	<i>Lissonota biguttata</i>
	<i>Eulimneria rufifemur</i> (Wylie 1960; Herting 1965)	<i>Sinophorus turionum</i>
	<i>Sinophorus rufifemur</i> (Sechser 1970)	
Braconidae	<i>Apanteles jucundus</i> (Wylie 1960; Herting 1965; Sechser 1970)	<i>Cotesia Jucunda</i>
	<i>Rogas testaceus</i> (Wylie 1960; Herting 1965; Sechser 1970)	<i>Rogas luteus</i>
Tachinidae	<i>Ernestia nielsenii</i> (Hassell 1966; Herting 1965)	<i>Panzeria laevigata</i>
	<i>Phryxe longicauda</i> (Wylie 1960; Herting 1965; Sechser 1970)	<i>Phryxe magnicornis</i>