

Characterization of the Functional Dynamics in the Neonatal Brain during REM and NREM Sleep States by means of Microstate Analysis

Journal: Brain Topography

Mohammad Khazaei^{1*}, Khadijeh Raeisi^{1*}, Pierpaolo Croce¹, Gabriella Tamburro^{1,2}, Anton Tokariev^{3,4}, Sampsa Vanhatalo^{3,4}, Filippo Zappasodi^{1,5}, Silvia Comani^{1,2}

¹ Department of Neuroscience, Imaging and Clinical Sciences, University “Gabriele d’Annunzio” of Chieti–Pescara, Chieti, Italy

² Behavioral Imaging and Neural Dynamics Center, University “Gabriele d’Annunzio” of Chieti–Pescara, Chieti, Italy

³ BABA center, Pediatric Research Center, Department of Clinical Neurophysiology, Children’s Hospital, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

⁴ Neuroscience center, Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland

⁵ Institute for Advanced Biomedical Technologies, University “Gabriele d’Annunzio” of Chieti–Pescara, Chieti, Italy

**Authors contributed equally to this work.*

Corresponding Author’s Email: filippo.zappasodi@unich.it

Fig. S1 Mean across subjects and standard deviation of the Power Spectral Density (PSD) in Active Sleep (AS, blu) and Quiet Sleep (QS, red) states. For spectral estimation, a Welch method was used with a frequency resolution of 0.1 Hz.

