

Supplementary Table 1: Trends in EAC and ESCC incidence and mortality by race/ethnicity in the United States

	Trend 1		Trend 2		Trend 3		Trend 4		Average APC
	Years	APC	Years	APC	Years	APC	Years	APC	Entire study period‡
EAC									
Incidence									
Overall	1992-2004	3.15*	2004-2016	0.00					1.6*
NHW	1992-1999	6.14*	1999-2016	0.97*					2.4*
Hispanic	1992-2016	1.04							1.0
NHB	1992-2016	1.80*							1.8*
NHAPI	1992-2016	1.62*							1.6*
NHAI/AN	1992-2016	2.36*							2.4*
Mortality									
Overall	1993-1995	17.77*	1995-1999	4.65*	1999-2012	1.23*	2012-2016	-1.67	2.6*
NHW	1993-1997	11.97*	1997-2010	2.01*	2010-2016	-0.34			3.1*
Hispanic	1993-2016	0.85							0.9*
NHB	1993-2016	1.60*							1.6*
NHAPI	1993-2016	0.65							0.7
NHAI/AN	1993-2016	2.26							2.3
ESCC									
Incidence									
Overall	1992-2016	-3.18*							-3.2*
NHW	1992-2016	-2.57*							-2.6*
Hispanic	1992-2016	-3.11*							-3.1*
NHB	1992-2016	-5.27*							-5.3*
NHAPI	1992-2016	-2.57*							-2.6*
NHAI/AN	1992-2016	-1.60							-1.6
Mortality									
Overall IBM	1993-1996	4.47	1996-2016	-3.08*					-2.1*
NHW IBM	1993-1995	14.18	1995-2016	-2.53*					-1.2
Hispanic IBM	1993-2016	-2.78*							-2.8*
NHB IBM	1993-2016	-4.53*							-4.5*
NHAPI IBM	1993-2016	-2.38*							-2.4*
NHAI/AN IBM	1993-2016	-1.79							-1.8

Average incidence rates are age-adjusted to the 2000 US standard population.

APC, annual percent change; EAC, esophageal adenocarcinoma; ESCC, esophageal squamous cell carcinoma; IR, incidence rate; NHW, non-Hispanic White; NHB, non-Hispanic Black; NHAPI, non-Hispanic Asian/Pacific Islander; NHAI/AN, non-Hispanic American Indian/Alaskan Native.

*The APC is significantly different from 0.0 (P < .05).

‡The entire study period for incidence rate is from 1992-2016 while the entire study period for incidence-based mortality is from 1993-2016.

Supplementary Table 2: Esophageal adenocarcinoma annual mortality rates (95% Cis) in the United States by race/ethnicity

Year of Death	Overall	NHW	Hispanic	NHB	NHAPI	NHAI/AN
1993-2016	2.5 (2.5-2.5)	3.2 (3.1-3.2)	1.5 (1.4-1.6)	0.8 (0.7-0.9)	0.6 (0.5-0.6)	1.8 (1.4-2.2)
1993	1.4 (1.24-1.56)	1.64 (1.45-1.85)	1.06 (0.61-1.68)	0.62 (0.32-1.09)	0.11 (0.01-0.38)	0.8 (0.02-4.05)
1994	1.8 (1.63-1.99)	2.2 (1.98-2.43)	0.83 (0.49-1.32)	0.55 (0.27-1)	0.34 (0.14-0.71)	1.79 (0.35-5.34)
1995	1.99 (1.81-2.18)	2.3 (2.08-2.54)	1.76 (1.18-2.51)	0.9 (0.51-1.45)	0.74 (0.4-1.26)	1.73 (0.2-5.68)
1996	2.07 (1.89-2.27)	2.52 (2.29-2.77)	1.34 (0.87-1.95)	0.48 (0.22-0.92)	0.46 (0.22-0.84)	1.74 (0.2-5.68)
1997	2.19 (2-2.39)	2.73 (2.49-2.98)	1.07 (0.64-1.67)	0.81 (0.46-1.31)	0.32 (0.13-0.67)	0.49 (0.01-2.94)
1998	2.29 (2.1-2.49)	2.8 (2.56-3.06)	1.56 (1.04-2.22)	0.69 (0.37-1.16)	0.55 (0.28-0.96)	1.41 (0.16-4.79)
1999	2.47 (2.27-2.67)	3.07 (2.82-3.34)	1.66 (1.14-2.33)	0.61 (0.31-1.06)	0.58 (0.31-0.97)	0.44 (0.01-2.72)
2000	2.46 (2.27-2.66)	3.13 (2.87-3.4)	1.97 (1.37-2.71)	0.68 (0.37-1.14)	0.33 (0.14-0.65)	0.39 (0.01-2.51)
2001	2.47 (2.28-2.67)	3.06 (2.81-3.33)	1.53 (1.05-2.14)	0.91 (0.55-1.42)	0.57 (0.32-0.94)	2.08 (0.51-5.42)
2002	2.45 (2.26-2.65)	3 (2.76-3.27)	2.03 (1.49-2.69)	0.69 (0.38-1.14)	0.69 (0.41-1.08)	1.66 (0.2-5.25)
2003	2.45 (2.26-2.64)	3.2 (2.94-3.47)	1.17 (0.78-1.67)	0.56 (0.29-0.98)	0.25 (0.1-0.52)	2.34 (0.58-5.93)
2004	2.55 (2.36-2.75)	3.27 (3.01-3.54)	1.35 (0.95-1.86)	0.61 (0.33-1.03)	0.65 (0.39-1.02)	1.51 (0.28-4.42)
2005	2.7 (2.51-2.91)	3.45 (3.19-3.73)	1.71 (1.24-2.28)	0.62 (0.34-1.04)	0.65 (0.39-1.01)	1.95 (0.61-4.7)
2006	2.64 (2.45-2.84)	3.39 (3.13-3.66)	1.5 (1.08-2.03)	0.67 (0.37-1.11)	0.88 (0.57-1.27)	2.32 (0.67-5.55)
2007	2.61 (2.42-2.81)	3.37 (3.11-3.65)	1.26 (0.89-1.72)	1.01 (0.64-1.52)	0.69 (0.44-1.04)	0.98 (0.2-3.11)
2008	2.74 (2.55-2.94)	3.59 (3.32-3.87)	1.57 (1.15-2.08)	0.88 (0.54-1.35)	0.46 (0.26-0.74)	1.48 (0.39-3.94)
2009	2.66 (2.48-2.86)	3.45 (3.19-3.73)	1.29 (0.95-1.71)	0.73 (0.43-1.17)	0.75 (0.49-1.1)	3.94 (1.77-7.47)
2010	2.82 (2.63-3.02)	3.75 (3.49-4.04)	1.27 (0.92-1.69)	1.03 (0.68-1.5)	0.54 (0.33-0.83)	2.9 (0.99-6.29)
2011	2.69 (2.51-2.89)	3.58 (3.32-3.85)	1.59 (1.21-2.04)	0.61 (0.36-0.97)	0.62 (0.39-0.92)	1.33 (0.43-3.35)
2012	2.9 (2.71-3.1)	3.78 (3.52-4.06)	1.84 (1.43-2.33)	0.83 (0.52-1.25)	0.67 (0.44-0.97)	1.91 (0.56-4.58)
2013	2.74 (2.56-2.93)	3.61 (3.35-3.88)	1.85 (1.44-2.34)	0.86 (0.54-1.28)	0.65 (0.43-0.95)	1.62 (0.48-3.95)
2014	2.76 (2.58-2.95)	3.69 (3.43-3.96)	1.64 (1.27-2.08)	1.03 (0.69-1.49)	0.54 (0.34-0.81)	2.09 (0.71-4.67)
2015	2.67 (2.5-2.86)	3.57 (3.32-3.84)	1.55 (1.19-1.97)	1.02 (0.7-1.45)	0.47 (0.3-0.71)	2.32 (0.9-4.85)
2016	2.64 (2.47-2.82)	3.58 (3.34-3.85)	1.68 (1.31-2.12)	0.74 (0.46-1.13)	0.47 (0.29-0.71)	1.79 (0.61-4.04)

95% CI, 95% confidence interval; NHW, Non-Hispanic White; NHB, Non-Hispanic Black; NHAPI, Non-Hispanic Asian and Pacific Islander; NHAI/AN, Non-Hispanic American Indian/Alaska Native.

Rates are age-adjusted to the 2000 US standard population (19 age groups-census p25-1130) standard.

Supplementary Table 3: Esophageal squamous cell carcinoma age-adjusted annual mortality rates (95% CIs) in the United States by race/ethnicity

Year of Death	Overall	NHW	Hispanic	NHB	NHAPI	NHAI/AN
1993-2016	2.1 (2.1-2.1)	1.7 (1.6-1.7)	1.5 (1.4-1.6)	5.9 (5.7-6.1)	2.2 (2.1-2.3)	2.5 (2.1-3.0)
1993	2.34 (2.15-2.55)	1.66 (1.47-1.87)	1.79 (1.2-2.55)	9.4 (8.04-10.92)	1.9 (1.32-2.64)	2.75 (0.69-7.08)
1994	2.74 (2.52-2.96)	1.95 (1.74-2.17)	2.41 (1.74-3.24)	10.14 (8.72-11.71)	2.88 (2.13-3.81)	2.03 (0.36-5.99)
1995	2.68 (2.47-2.9)	2.16 (1.95-2.39)	1.99 (1.36-2.78)	8.07 (6.84-9.45)	2.53 (1.85-3.37)	2.12 (0.57-5.57)
1996	2.75 (2.54-2.97)	2.1 (1.89-2.33)	1.63 (1.1-2.31)	10.19 (8.8-11.73)	2.13 (1.54-2.87)	1.3 (0.15-4.54)
1997	2.66 (2.45-2.87)	2.01 (1.8-2.23)	2.1 (1.49-2.86)	9.03 (7.74-10.47)	2.62 (1.95-3.42)	2.1 (0.4-5.97)
1998	2.85 (2.64-3.07)	2.2 (1.99-2.43)	1.67 (1.15-2.34)	8.24 (7.03-9.59)	3.55 (2.8-4.43)	7.36 (3.6-13.07)
1999	2.62 (2.42-2.83)	1.99 (1.79-2.21)	1.51 (1.01-2.15)	8.56 (7.31-9.95)	3.31 (2.59-4.14)	2.19 (0.57-5.61)
2000	2.32 (2.13-2.52)	1.77 (1.58-1.97)	1.85 (1.31-2.52)	7.01 (5.89-8.27)	2.78 (2.15-3.52)	1.78 (0.34-5.1)
2001	2.41 (2.22-2.61)	1.88 (1.68-2.09)	2.17 (1.6-2.87)	6.79 (5.72-8.01)	2.49 (1.92-3.17)	2.4 (0.76-5.64)
2002	2.32 (2.13-2.51)	1.89 (1.7-2.11)	1.71 (1.21-2.34)	6.44 (5.4-7.61)	2.34 (1.79-3)	4.88 (1.88-9.84)
2003	2.23 (2.06-2.42)	1.74 (1.56-1.95)	1.48 (1.04-2.03)	6.71 (5.66-7.9)	2.43 (1.88-3.08)	3.43 (1.29-7.19)
2004	2.15 (1.97-2.33)	1.71 (1.53-1.92)	1.48 (1.02-2.05)	6.76 (5.71-7.94)	1.86 (1.39-2.42)	3.11 (1.04-6.88)
2005	2.18 (2-2.36)	1.77 (1.58-1.97)	1.79 (1.31-2.37)	5.48 (4.55-6.55)	2.41 (1.89-3.04)	2.58 (0.92-5.69)
2006	2.1 (1.93-2.28)	1.79 (1.6-2)	1.53 (1.09-2.06)	5.03 (4.14-6.05)	2.11 (1.63-2.69)	4.4 (2-8.28)
2007	1.87 (1.71-2.03)	1.41 (1.25-1.59)	1.35 (0.96-1.83)	5.35 (4.45-6.38)	2.31 (1.81-2.91)	2.13 (0.52-5.42)
2008	1.95 (1.79-2.12)	1.62 (1.44-1.81)	1.25 (0.87-1.73)	5.32 (4.42-6.33)	1.99 (1.54-2.53)	2.86 (0.97-6.3)
2009	1.85 (1.69-2.01)	1.5 (1.33-1.68)	1.42 (1.03-1.9)	4.86 (4.02-5.82)	1.88 (1.45-2.39)	1.95 (0.48-4.96)
2010	1.77 (1.62-1.94)	1.35 (1.19-1.53)	1.5 (1.1-1.98)	4.93 (4.08-5.89)	2.15 (1.7-2.68)	1.05 (0.13-3.51)
2011	1.81 (1.66-1.97)	1.56 (1.39-1.75)	1.34 (0.97-1.79)	4.18 (3.43-5.05)	1.82 (1.42-2.3)	1.63 (0.47-4.04)
2012	1.62 (1.48-1.77)	1.48 (1.31-1.66)	1.01 (0.7-1.4)	3.72 (3.01-4.55)	1.47 (1.11-1.9)	1.45 (0.36-3.83)
2013	1.63 (1.49-1.78)	1.3 (1.15-1.47)	1.02 (0.71-1.4)	4.59 (3.82-5.47)	1.73 (1.35-2.18)	3.03 (1.23-6.08)
2014	1.47 (1.34-1.61)	1.18 (1.04-1.34)	1.23 (0.91-1.62)	3.21 (2.58-3.93)	1.97 (1.58-2.44)	0.26 (0.01-1.73)
2015	1.61 (1.48-1.76)	1.39 (1.24-1.56)	0.94 (0.67-1.28)	3.53 (2.86-4.3)	1.85 (1.47-2.29)	3.32 (1.45-6.38)
2016	1.64 (1.51-1.78)	1.34 (1.19-1.51)	1.35 (1.02-1.76)	4.09 (3.4-4.89)	1.64 (1.29-2.05)	2.2 (0.91-4.51)
95% CI, 95% confidence interval; NHW, Non-Hispanic White; NHB, Non-Hispanic Black; NHAPI, Non-Hispanic Asian and Pacific Islander; NHAI/AN, Non-Hispanic American Indian/Alaska Native.						
Rates are age-adjusted to the 2000 US standard population (19 age groups-census p25-1130) standard.						

Supplementary Table 4: Esophageal adenocarcinoma annual proportion of unstaged and late stage disease at diagnosis in the United States by race/ethnicity

	Overall		NHW		Hispanic		NHB		NHAPI	
Year of diagnosis	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)
1992-2015	10.8 (0.2)	65.5 (0.3)	10.4 (0.2)	65.3 (0.4)	13.6 (1.0)	66.6 (1.3)	15.6 (1.6)	65.3 (2.1)	12.6 (1.5)	69.2 (2.1)
1992	22.9 (1.9)	55.2 (2.3)	21.7 (2)	55.5 (2.4)	30.4 (9.6)	60.9 (10.2)	46.2 (13.8)	46.2 (13.8)	30 (14.5)	50 (15.8)
1993	17.9 (1.7)	57.9 (2.2)	18.3 (1.9)	58.7 (2.4)	4.3 (4.3)	52.2 (10.4)	23.1 (11.7)	69.2 (12.8)	12.5 (11.7)	25 (15.3)
1994	17.3 (1.7)	54.1 (2.2)	17.4 (1.8)	53.7 (2.4)	19.4 (6.6)	52.8 (8.3)	18.2 (11.6)	72.7 (13.4)	6.7 (6.4)	60 (12.6)
1995	17.5 (1.7)	58.8 (2.2)	18.5 (1.8)	57.7 (2.3)	15.4 (7.1)	57.7 (9.7)	0 (0)	88.9 (10.5)	0 (0)	88.9 (10.5)
1996	17 (1.5)	56.8 (1.9)	16.8 (1.6)	56.1 (2.1)	10.8 (5.1)	64.9 (7.8)	33.3 (10.3)	52.4 (10.9)	13.3 (8.8)	73.3 (11.4)
1997	17.6 (1.5)	54.1 (2)	17.4 (1.6)	53.6 (2.1)	21.1 (6.6)	57.9 (8)	23.1 (11.7)	46.2 (13.8)	15.4 (10)	69.2 (12.8)
1998	18.7 (1.5)	58.3 (1.9)	18.2 (1.6)	58.4 (2)	20 (6.8)	54.3 (8.4)	33.3 (15.7)	55.6 (16.6)	22.2 (13.9)	66.7 (15.7)
1999	13.5 (1.3)	61.6 (1.8)	11.7 (1.3)	62.7 (1.9)	32.6 (7.1)	46.5 (7.6)	27.8 (10.6)	61.1 (11.5)	23.1 (11.7)	53.8 (13.8)
2000	12.1 (1.2)	59.3 (1.8)	10.9 (1.2)	60.2 (1.9)	21.6 (6.8)	54.1 (8.2)	30 (10.2)	40 (11)	17.6 (9.2)	58.8 (11.9)
2001	12 (1.2)	61.8 (1.8)	11.6 (1.3)	61 (1.9)	17 (5.5)	70.2 (6.7)	11.1 (7.4)	72.2 (10.6)	20 (10.3)	73.3 (11.4)
2002	9.7 (1.1)	66.9 (1.7)	8.8 (1.1)	68.3 (1.8)	21.4 (6.3)	57.1 (7.6)	16.7 (8.8)	55.6 (11.7)	6.7 (6.4)	53.3 (12.9)
2003	12.2 (1.2)	61.7 (1.8)	11.2 (1.2)	61.5 (1.9)	21.1 (6.6)	57.9 (8)	26.7 (11.4)	66.7 (12.2)	13.3 (8.8)	80 (10.3)
2004	10.7 (1)	63.9 (1.6)	10.8 (1.1)	63.2 (1.7)	9.7 (3.8)	72.6 (5.7)	0 (0)	72.2 (10.6)	19.4 (7.1)	58.1 (8.9)
2005	9.9 (1)	66.9 (1.6)	9.4 (1.1)	67.7 (1.7)	12.9 (4.3)	62.9 (6.1)	33.3 (10.3)	57.1 (10.8)	0 (0)	64.5 (8.6)
2006	7.5 (0.9)	69.9 (1.6)	7.3 (1)	69.2 (1.7)	12.8 (5.4)	76.9 (6.7)	12.5 (8.3)	62.5 (12.1)	0 (0)	87.5 (6.8)
2007	6.8 (0.8)	66.5 (1.6)	7 (0.9)	64.9 (1.7)	5.7 (3.2)	77.4 (5.7)	3.8 (3.8)	80.8 (7.7)	5.6 (5.4)	83.3 (8.8)
2008	7.4 (0.8)	69.5 (1.5)	6.8 (0.9)	70.3 (1.6)	8.6 (3.7)	67.2 (6.2)	11.1 (6)	51.9 (9.6)	17.9 (7.2)	67.9 (8.8)
2009	8.2 (0.9)	68.4 (1.5)	8 (0.9)	67.8 (1.6)	9.1 (3.3)	72.7 (5.1)	6.3 (4.3)	78.1 (7.3)	13 (7)	73.9 (9.2)
2010	8.6 (0.9)	68.4 (1.5)	8 (1)	68.3 (1.6)	11.3 (4)	75.8 (5.4)	12.5 (6.8)	66.7 (9.6)	17.9 (7.2)	60.7 (9.2)
2011	8 (0.8)	71.2 (1.4)	7.7 (0.9)	70.6 (1.5)	11.6 (3.5)	72.1 (4.8)	9.5 (6.4)	76.2 (9.3)	4 (3.9)	88 (6.5)
2012	8.1 (0.8)	70.7 (1.4)	7.8 (0.9)	70.2 (1.5)	10.1 (3)	73.7 (4.4)	6.1 (4.2)	78.8 (7.1)	15.6 (6.4)	62.5 (8.6)
2013	5.9 (0.7)	70.6 (1.4)	5.4 (0.7)	70.5 (1.5)	13.9 (4.1)	65.3 (5.6)	2.7 (2.7)	73 (7.3)	9.5 (6.4)	85.7 (7.6)
2014	8.9 (0.9)	71.8 (1.4)	8 (0.9)	72.7 (1.5)	11 (3.5)	67.1 (5.2)	20.6 (6.9)	55.9 (8.5)	14.7 (6.1)	76.5 (7.3)
2015	6.4 (0.7)	73.7 (1.3)	5.4 (0.7)	74.5 (1.4)	9.3 (2.8)	72.2 (4.3)	11.5 (6.3)	69.2 (9.1)	14.3 (5.9)	68.6 (7.8)

NHW, Non-Hispanic White; NHB, Non-Hispanic Black; NHAPI, Non-Hispanic Asian and Pacific Islander.

SE, standard error

Counts by year were too sparse for Non-Hispanic American Indian/Alaska Native.

Supplementary Table 5: Esophageal squamous cell carcinoma annual proportion of late and unstaged disease at diagnosis in the United States by race/ethnicity

	Overall		NHW		Hispanic		NHB		NHAPI	
Year of diagnosis	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)	Proportion Unstaged (SE)	Proportion Late Stage (SE)
1992-2015	16.1 (0.3)	60.1 (0.4)	16.4 (0.4)	58.4 (0.5)	19.2 (1.2)	59.7 (1.5)	15.2 (0.6)	61.5 (0.8)	14.3 (0.8)	64.8 (1.1)
1992	23.1 (1.5)	47.7 (1.8)	25.6 (2.1)	45.1 (2.4)	20 (5.4)	52.7 (6.7)	19.3 (2.6)	49.8 (3.3)	22 (6.5)	56.1 (7.8)
1993	22.3 (1.5)	47 (1.8)	24.5 (2.1)	46.1 (2.4)	18.4 (5.5)	49 (7.1)	20.5 (2.7)	48.4 (3.4)	18.2 (4.7)	47 (6.1)
1994	22 (1.6)	50.4 (1.9)	23.4 (2.1)	50.5 (2.5)	11.9 (5)	59.5 (7.6)	21.7 (2.9)	46 (3.5)	20.8 (5.6)	58.5 (6.8)
1995	21.7 (1.5)	45.9 (1.8)	20.8 (2)	46 (2.4)	26.1 (6.5)	39.1 (7.2)	20.4 (2.9)	48.5 (3.6)	27.9 (5.7)	42.6 (6.3)
1996	21.3 (1.5)	50.6 (1.9)	22.7 (2.1)	49.4 (2.5)	20 (6.3)	57.5 (7.8)	20.3 (2.7)	52.1 (3.4)	18.4 (4.4)	46.1 (5.7)
1997	21.7 (1.6)	49.6 (1.9)	22.7 (2.1)	46.1 (2.5)	24.3 (7.1)	56.8 (8.1)	21.9 (3.1)	52.8 (3.7)	15.5 (4.3)	54.9 (5.9)
1998	19.2 (1.5)	55.3 (1.9)	22.1 (2.1)	51.2 (2.5)	17.6 (6.5)	61.8 (8.3)	13.9 (2.7)	60.8 (3.8)	17 (4)	63.6 (5.1)
1999	17.9 (1.5)	54.7 (1.9)	19.4 (2)	51.1 (2.6)	20.8 (5.6)	50.9 (6.9)	17.1 (3.1)	59.9 (4)	9.2 (3.3)	65.8 (5.4)
2000	17.1 (1.5)	54.6 (2)	15.8 (2)	50 (2.7)	23.4 (6.2)	57.4 (7.2)	18.2 (3.1)	59.7 (4)	16.7 (4.4)	62.5 (5.7)
2001	13.4 (1.3)	61.8 (1.9)	13.4 (1.7)	61.6 (2.5)	28.3 (6.6)	50 (7.4)	12.8 (2.7)	59.7 (4)	6.8 (2.9)	73 (5.2)
2002	14.4 (1.4)	64.6 (1.9)	14.3 (1.9)	59.9 (2.7)	21.1 (6.6)	63.2 (7.8)	14 (2.8)	70 (3.7)	11.5 (3.6)	74.4 (4.9)
2003	15.2 (1.5)	62 (2)	14.2 (1.9)	60.7 (2.7)	18.2 (5.8)	68.2 (7)	16.7 (2.9)	63.7 (3.7)	15.5 (4.8)	56.9 (6.5)
2004	13 (1.3)	60.3 (1.9)	11.3 (1.6)	62.2 (2.5)	19.6 (5.6)	51 (7)	14.4 (2.9)	56.2 (4.1)	15.7 (3.9)	60.7 (5.2)
2005	12.2 (1.4)	65.2 (2)	12 (1.8)	65.6 (2.7)	11.6 (4.9)	67.4 (7.1)	15.1 (3.2)	61.9 (4.3)	9.7 (3.5)	66.7 (5.6)
2006	12.8 (1.4)	64.5 (2)	12.3 (1.8)	64.2 (2.7)	23.3 (6.4)	60.5 (7.5)	10.2 (2.6)	65 (4.1)	13.3 (3.7)	66.3 (5.2)
2007	12.7 (1.4)	65.1 (2)	14.2 (2)	63.2 (2.7)	18.8 (5.6)	58.3 (7.1)	8.3 (2.4)	72.2 (3.9)	11 (3.7)	65.8 (5.6)
2008	13.9 (1.5)	67.8 (2)	14 (2)	68.6 (2.7)	17.1 (5.9)	53.7 (7.8)	10.7 (2.8)	71.9 (4.1)	15.1 (3.9)	67.4 (5.1)
2009	13.7 (1.4)	66.7 (1.9)	11.7 (1.7)	65.1 (2.6)	26.5 (6.3)	63.3 (6.9)	15.4 (3.2)	73.1 (3.9)	10.3 (3.7)	66.2 (5.7)
2010	11.7 (1.3)	69.7 (1.9)	9.9 (1.7)	66.8 (2.7)	22.4 (5.1)	59.7 (6)	8.9 (2.6)	75.8 (3.8)	13.5 (3.5)	78.1 (4.2)
2011	12.3 (1.4)	69.5 (2)	13.5 (1.9)	66.1 (2.6)	14 (5.3)	67.4 (7.1)	7.3 (2.5)	80.7 (3.8)	13.9 (4.1)	69.4 (5.4)
2012	9.4 (1.3)	73.5 (1.9)	8.3 (1.6)	75.2 (2.5)	14.3 (5)	75.5 (6.1)	8.5 (2.7)	67.9 (4.5)	11.1 (3.7)	73.6 (5.2)
2013	9.7 (1.2)	73.6 (1.8)	9.6 (1.7)	73.2 (2.5)	8.8 (3.7)	80.7 (5.2)	9.5 (2.6)	72.2 (4)	10.2 (3.2)	72.7 (4.7)
2014	13 (1.4)	69.5 (2)	14.6 (2.1)	66.8 (2.7)	14 (4.6)	64.9 (6.3)	10 (2.9)	76.4 (4.1)	12.2 (3.6)	73.2 (4.9)
2015	12.2 (1.3)	72.2 (1.8)	11.7 (1.8)	69.8 (2.5)	20.4 (5.5)	63 (6.6)	6.5 (2.4)	84.3 (3.5)	16 (3.7)	72 (4.5)

NHW, Non-Hispanic White; NHB, Non-Hispanic Black; NHAPI, Non-Hispanic Asian and Pacific Islander; NHAI/AN, Non-Hispanic American Indian/Alaska Native.

SE, standard error

Counts by year were too sparse for Non-Hispanic American Indian/Alaska Native.