

Online Resource – Conservation Genetics

Genetic structure of coastal and inland populations of *Spergularia media* (L.) C. Presl (Caryophyllaceae) in Central Europe

Kathleen Prinz^{1*}, Kurt Weising², and Isabell Hensen³

¹Forest Genetics and Forest Tree Breeding, Georg-August-University Göttingen, Buisen-Institute, Buisenweg 2, D-37077 Göttingen, Germany


²Plant Molecular Systematics, University of Kassel, Heinrich-Plett-Straße 40, D-34109 Kassel, Germany

³Institute of Biology/Geobotany and Botanical Garden, Martin-Luther-University Halle-Wittenberg, Am Kirchtor 1, D-06108 Halle/Saale, Germany


***Corresponding author:** kathleen.prinz@forst.uni-goettingen.de

telephone: 0049-551-3914281

fax number: 0049-551-398367



ESM 1 NeighborNet based on Nei-Li distances (1979) among 179 individual plants of *S. media*. The numbers at the ends of the branches refer to the populational assignment (1 to 18) of the respective individuals (see Table 1).



ESM 2 Individual-level results of STRUCTURE analyses on based on 179 individual plants of *S. media* for $K = 5$ (upper panel) and $K = 7$ (lower panel). Each of the coloured vertical bars represents a single individual. The vertical axis refers to the Bayesian admixture coefficients. The numbers below the diagrams denote the population assignment of the 179 individuals (population 1 to 18; see Table 1).