

Conservation Genetics

Conservation Genomics Reveals Multiple Evolutionary Units Within Bell's Vireo (*Vireo bellii*)

Luke B. Klicka^{1,2}, Barbara E. Kus³, Pascal O. Title⁴, and Kevin J. Burns¹

1. Department of Biology, San Diego State University, 5500 Campanile Drive, San Diego, CA, 92182

2. Current address: University of Kansas Biodiversity Institute, 1345 Jayhawk Blvd, Lawrence, KS 66049

3. United States Geological Survey, 4165 Spruance Rd, San Diego, CA 92101

4. Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109

Corresponding author: Luke B. Klicka, Klicka@ku.edu, telephone: 763-222-8373

Table S2 AMOVA results for both ND2 and SNP datasets partitioned by sampling populations

Dataset	Source of Variation	Percent of Variation	Φ -statistics	P-value
SNPs	East vs. West	6.46	0.065	0.016
	Among populations within East or West	4.63	0.049	< 0.00001
	Within populations	88.91	0.11	<0.00001
ND2	East vs. West	90.00	0.914	< 0.00001
	Among populations within East or West	1.43	0.143	0.0039
	Within populations	8.57	0.900	0.0105