

Title: Grass-skipper (Hesperiinae) trends in midwestern USA grasslands during 1988-2013

Journal: Journal of Insect Conservation

Ann B. Swengel* and Scott R. Swengel

No affiliation; 909 Birch Street, Baraboo, WI 53913 USA

* corresponding author (email: swengel@naba.org; tel. +608-356-9543)

Online Resource Table 3 4th of July Counts in the northern tallgrass prairie region analyzed in this study. N years = number of years the Count has been reported. Co = County; NC = Nature Center; SP = State Park

State, Count	latitude	longitude	first year	last year	N years	Party-hours
Iowa						
Waubonsie	40.7662	95.7401	1986	2012	27	249.80
Illinois						
Beardstown	39.9484	90.4278	1995	2012	16	88.75
McDonough Co	40.3683	90.7719	1977	2011	26	120.28
Minnesota						
Bluestem	46.8552	96.4734	1993	2012	10	68.17
Central Polk Co	47.595	96.3658	1994	2012	19	167.43
Ramsey Co	45.0903	93.1667	1993	2012	18	79.64
Nebraska						
Indian Cave SP	40.3151	95.6149	1986	2012	27	207.08
South Dakota						
Strandburg	45.0410	96.7607	1992	2012	21	85.0533
Yankton	42.9959	97.3702	1993	2012	20	71.60
Wisconsin						
Baraboo	43.4891	89.7387	1986	2012	27	205.637
Madison	43.0899	89.3838	1991	2012	22	339.9
Mud Lake	43.4118	89.3044	1989	2012	24	188.60
Riveredge NC	43.3959	87.9793	1987	2012	19	270.5
Trempealeau	44.0701	91.4112	1989	2012	23	264.9
Wehr NC	42.9304	87.9791	1988	2012	25	158.44