

The graph displays the performance of four gene identification methods across six datasets. The y-axis represents the number of genes identified, ranging from 0 to 10,000. The x-axis lists the datasets: bee, dorsata, tetragonula, cerana, florea, and notbee. The legend indicates: wbnc (blue), wboc (orange), wobnc (green), and woboc (red).

Dataset	wbnc	wboc	wobnc	woboc
bee	~1,500	~3,500	~1,500	~1,500
dorsata	~4,500	~7,500	~5,500	~9,500
tetragonula	~7,500	~7,500	~9,500	~9,500
cerana	~3,500	~1,000	~5,500	~3,500
florea	~8,500	~8,500	~9,500	~9,500
notbee	~9,500	~9,000	~9,500	~9,500

IMAGE BASED SOUTH ASIAN BEE SPECIES IDENTIFICATION: A MACHINE LEARNING APPROACH
Srinivas C L¹(srinivas.cl@niser.ac.in), Sneering M¹(sneeringm@niser.ac.in), Ronita Mukherjee²(ronita@niser.ac.in),
Soubadra M Dey³(soubadra@niser.org), Ritik Deb⁴(ritikdeb@niser.ac.in),
Subhanika Mishra¹(mishra@niser.ac.in)
¹National Institute of Science Education and Research, Bhubaneswar
²Ashoka Trust for Research in Ecology and the Environment, Bangalore
Journal of Insect Conservation