


Manometric determination of the oxygen diffusion coefficients in



J. Kargin^{a(*)}, H.E Sanchez Cornejo^b, K. Zhangozin^c, T. Zhanabergenov^c, Zh. Kassymkhanov^a, Jiwon Seo^d, S.N. Holmes^e, C.H.W. Barnes^f, J. Albino Aguiar^g, L. De Los Santos Valladares^{f,g(*)}

^a *L.N. Gumilyov Eurasian National University, NPJSC, Astana, Kazakhstan*

^b *Laboratorio de Ceramicos y Nanomateriales, Facultad de Ciencias Físicas, Universidad Nacional Mayor de San Marcos, Ap. Postal 14-0149, Lima, Peru*

^c *TSK-Vostok, Astana, Kazakhstan.*

^d *College of Science and Technology, Yonsei University, 1 Yonseidae-gil, Wonju, Gangwon-do, 26493, South Korea.*

^e *Department of Electronic and Electrical Engineering, University College London, Torrington Place, London WC1E 7JE, UK.*

^f *Cavendish Laboratory, Department of Physics, University of Cambridge, J.J. Thomson Ave., Cambridge CB3 0HE, UK.*

^g *Programa de Pós-Graduação em Ciências de Materiais, Centro de Ciências Exatas e da Natureza, Universidade Federal de Pernambuco, 50670-901 Recife-PE, Brazil.*

(*) Corresponding authors emails: kjb_orken@mail.ru (J. Kargin) and ld301@cam.ac.uk (L. De Los Santos Valladares)

Supporting Material




Figure S1 -Typical view of the vacuum degassing curve under continuous pumping conditions.