

Multi-scale landscape composition and local factors jointly shape the occurrence patterns of declining peatland dragonflies

Giacomo Assandri¹, Felix Puff^{2,3}, Leonardo Siddi¹, Matteo Anderle³, Chiara Fedrigotti⁴

¹ Dipartimento di Scienze e Innovazione Tecnologica, Università del Piemonte Orientale “Amedeo Avogadro”, 15121 Alessandria (AL), Italy.

²Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030 Vienna, Austria

³ Institute for Alpine Environment, Eurac Research, Drususallee/Viale Druso 1, I-39100, Bolzano/Bozen, Italy

⁴ Museum and Research Collection Office, MUSE-Science Museum Trento, Corso del Lavoro e della Scienza, 3 - 38122 Trento, Italy.

Supplementary informations



Figure S1. Example of peatland habitats occurring in the study area. (a) Wooded bogs [Paluda La Lot], (b) bogs, in the foreground is visible a clear example of *Rhynchosporion* depression on peat substrate habitat, often occurring intermixed with *Sphagnum* hummocks at bogs [Palù Tremole]; (c) poor/intermediate fens [Lago delle Buse], and (d) rich fens [Torbiera delle Grave].