

Electronic Supplementary Material 3 to

Can methyl jasmonate treatment of conifer seedlings be used as a tool to stop height growth in nursery forest trees?

Published in New Forests

Frauke Fedderwitz^{1*}, Niklas Björklund¹, Ronny Anngren^{2^}, Anders Lindström³, Göran Nordlander¹

¹Department of Ecology, Swedish University of Agricultural Sciences (SLU), Box 7044, SE-75007 Uppsala, Sweden

²Nässja Tree Nursery, Bergvik Skog Plantor AB, Berreksvägen 45, SE-81020 Österfärnebo, Sweden

³School of Industrial Technology and Business Studies, Dalarna University, SE-79188 Falun, Sweden

*Current address: Crop Research Centre, Teagasc, Oak Park, Ireland; Corresponding author: email: frauke.fedderwitz@teagasc.ie

^Current address: Rural Economy and Agricultural Society Halland, Lilla Böslid 146, SE-30596 Eldsberga, Sweden

Online Resource 3 Exact p-values for freezing tolerance at –10°C and –25°C (one-sample Wilcoxon rank sum tests against a cut-off value of 4; all data from batch B seedlings). Comparisons indicated with “–” were not carried out

Treatment	September		October		November
	-10	-25	-10	-25	-25
RLN	0.031	0.031	0.031	0.031	0.031
0 mM MeJA	0.313	1	0.031	0.031	0.031
5 mM MeJA	-	-	0.156	1	0.031
10 mM MeJA	1	1	0.5	1	0.031
10 mM MeJA no LN	1	1	-	-	-
15 mM MeJA	-	-	1	1	0.094