

Expression Patterns of Carbonic Anhydrase IX and Immune Response Biomarkers in Pediatric Gliomas and Glioneuronal Tumors

Journal of Neuro-Oncology

Monika Kapitančukė¹, Donatas Petroška², Agnė Petrošiūtė³, Asta Lučiūnaitė³, Švitrigailė Grincevičienė³, Daumantas Matulis³, Saulius Ročka¹, Ramūnas Raugalas¹, Dainius Rinkevičius¹, Irena Adomaitienė⁴, Gabrielė Tarutytė⁵, Jelena Rascon¹

¹ Institute of Clinical Medicine, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

² National Center of Pathology, Affiliate of Vilnius University Hospital Santaros Klinikos, Vilnius, Lithuania

³ Life Sciences Center, Vilnius University, Vilnius, Lithuania

⁴ Center of Radiology, Vilnius University Hospital Santaros Klinikos, Vilnius, Lithuania

⁵ Department of Research and Innovation, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

Corresponding author:

Monika Kapitančukė


Center of Pediatric Oncology and Hematology

Vilnius University Hospital Santaros Klinikos

Santariskiu 7, LT-08406

Vilnius, Lithuania

E-mail: monika.kapitancuke@santa.lt



SI2. CAIX H-score staining intensity categories

(A) Negative IHC reaction. (B) Weak staining intensity. (C) Moderate staining intensity. (D) Strong staining intensity. No cases with strong staining intensity were observed in the pediatric glioma and GNT cohort. An example of strong CAIX expression is shown from a

renal cell carcinoma specimen used as a positive control. All images were acquired at 400× magnification.