

International Journal of Clinical Pharmacy

Online Resource 2

to

Effect of CYP2D6 pharmacogenetic phenotype and phenoconversion on serum concentrations of antidepressants and antipsychotics - a retrospective cohort study

Maike Scherf-Clavel, PhD¹, Amelie Frantz², Andreas Eckert, MD², Heike Weber, PhD^{1,2}, Stefan Unterecker, MD¹, Jürgen Deckert, MD¹, Andreas Reif, MD², Martina Hahn, PhD^{2,3}

¹ Department of Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University Hospital of Würzburg, 97080 Würzburg, Germany

² Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt, 60528 Frankfurt, Germany

³ Department of mental health, varisano Hospital Frankfurt Hoechst, Germany

Corresponding author:

Dr. rer. nat. Maike Scherf-Clavel
Department of Psychiatry, Psychosomatics and Psychotherapy
University Hospital of Würzburg
Margarete-Höppel-Platz 1
97080 Würzburg, Germany
Tel.: +49/931/201 77546
Fax: +49/931/201 77262
E-Mail: Scherf_M@ukw.de

Table 2 Drugs with the propensity to cause phenoconversion due to inhibitory or inducing effects on CYP2D6 according to the Flockhart table [The Flockhart Cytochrome P450 Drug-Drug Interaction Table. Updated 2021. <https://drug-interactions.medicine.iu.edu/>; cited: January, 12, 2022].

CYP2D6	
Inhibitors	Inducers
STRONG	Dexamethasone
Bupropion	Oritavancin
Cinacalcet	Rifampin
Fluoxetine	
Paroxetine	
Quinidine	
MODERATE	
Abiraterone	
Doxepin	
Duloxetine	
Halofantrine	
Lorcaserin	
Moclobemide	
Rolapitant	
Terbinafine	
WEAK	
Amiodarone	
Celecoxib	
Cimetidine	
Citalopram	
Clobazam	
Clomipramine	
Diphenhydramine	
Escitalopram	
Levomepromazine	
Sertraline	