

Online Resource 3

to

Effect of CYP2D6 pharmacogenetic phenotype and phenoconversion on serum concentrations of antidepressants and antipsychotics - a retrospective cohort study

Maïke Scherf-Clavel, PhD¹, Amelie Frantz², Andreas Eckert, MD², Heike Weber, PhD^{1,2}, Stefan Unterecker, MD¹, Jürgen Deckert, MD¹, Andreas Reif, MD², Martina Hahn, PhD^{2,3}

¹ Department of Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University Hospital of Würzburg, 97080 Würzburg, Germany

² Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt, 60528 Frankfurt, Germany

³ Department of mental health, varisano Hospital Frankfurt Hoechst, Germany

Corresponding author:

Dr. rer. nat. Maïke Scherf-Clavel
Department of Psychiatry, Psychosomatics and Psychotherapy
University Hospital of Würzburg
Margarete-Höppel-Platz 1
97080 Würzburg, Germany
Tel.: +49/931/201 77546
Fax: +49/931/201 77262
E-Mail: Scherf_M@ukw.de

Table 3 Number (N) of administered psychiatric drugs with serum concentration determinations (TDM) in the combined sample.

ANTIDEPRESSANTS	N	ANTIPSYCHOTICS	N	ANTIEPILEPTICS	N
Venlafaxine	117	Quetiapine	125	Pregabalin	25
Amitriptyline	100	Risperidone	73	Pipamperone	20
Mirtazapine	85	Aripiprazole	32	Valproic Acid	15
Sertraline	64	Olanzapine	20	Lamotrigine	13
Escitalopram	52	Cariprazine	10	Oxcarbazepine	9
Bupropion	38	Clozapine	8	Gabapentine	3
Trazodon	32	Chlorprothixene	7	Carbamazepine	2
Duloxetine	28	Amisulpride	5	Topiramate	2
Clomipramine	27	Haloperidol	5	Levetiracetam	1
Milnacipran	19	Perazine	4		
Doxepin	16	Melperone	3		
Trimipramine	5	Benperidol	2		
Fluoxetine	4	Flupentixol	2		
Moclobemid	3	Fluphenazine	1		
Citalopram	2				
Maprotiline	1				
Opipramole	1				