

Evaluating the quality, feasibility and patient satisfaction of medication history taking by telephone for patients with planned admissions to two gastroenterology wards

– Supplement E –

Theresa Terstegen^a, Janina Bittmann^a, Luise Kauk^a, Marietta Kirchner^b, Sebastian Krug^c, Annika Gauss^c, Ute Chiriac^d, Benedict Morath^d, Walter E. Haefeli^a, Hanna M. Seidling^a

^aHeidelberg University, Medical Faculty Heidelberg / Heidelberg University Hospital, Internal Medicine IX, Clinical Pharmacology and Pharmacoepidemiology, Cooperation Unit Clinical Pharmacy, Im Neuenheimer Feld 410, 69120 Heidelberg, Germany.

^bHeidelberg University, Medical Faculty Heidelberg / Heidelberg University Hospital, Institute of Medical Biometry, Im Neuenheimer Feld 103.3, 69120 Heidelberg, Germany.

^cHeidelberg University, Medical Faculty Heidelberg / Heidelberg University Hospital, Internal Medicine IV, Department of Gastroenterology, Infectiology and Toxicology, Im Neuenheimer Feld 410, 69120 Heidelberg, Germany.

^dHeidelberg University, Medical Faculty Heidelberg / Heidelberg University Hospital, Hospital Pharmacy, Im Neuenheimer Feld 670, 69120 Heidelberg, Germany.

International Journal of Clinical Pharmacy

Corresponding Author

Prof. Dr. sc. hum. Hanna M. Seidling

Heidelberg University, Medical Faculty Heidelberg / Heidelberg University Hospital, Internal Medicine IX, Clinical Pharmacology and Pharmacoepidemiology, Cooperation Unit Clinical Pharmacy, Im Neuenheimer Feld 410, 69120, Heidelberg, Germany. hanna.seidling@med.uni-heidelberg.de

Supplement Ea: Overview of subtypes of changes in the category “changed medicines”.

Type of update in medicines	Intervention (n=76)			Control (n=75)			Overall (n=151)		
	Mean (SD)	Range	N (%)	Mean (SD)	Range	N (%)	Mean (SD)	Range	N (%)
Overall changes	2.43 (± 2.68)	0–13	185 (100)	1.27 (± 1.98)	0–11	95 (100)	1.85 (± 2.42)	0–13	280 (100)
Dose	1.00 (± 1.29)	0–6	76 (41.1)	0.68 (± 1.04)	0–5	51 (53.7)	0.84 (± 1.2)	0–6	127 (45.4)
Frequency	0.80 (± 1.18)	0–6	61 (33.0)	0.40 (± 0.82)	0–5	30 (31.6)	0.60 (± 1.0)	0–6	91 (32.5)
Time of intake	0.21 (± 0.47)	0–2	16 (8.6)	0.11 (± 0.31)	0–1	8 (8.4)	0.16 (± 0.40)	0–2	24 (8.6)
Dosage form	0.13 (± 0.44)	0–3	10 (5.4)	0.04 (± 0.20)	0–1	3 (3.2)	0.09 (± 0.35)	0–3	13 (4.6)
Paused/resumed medicine	0.16 (± 0.49)	0–3	12 (6.5)	0.01 (± 0.12)	0–1	1 (1.1)	0.09 (± 0.36)	0–3	13 (4.6)
Single-agent products/fixed-dose combination product	0.09 (± 0.33)	0–2	7 (3.8)	-	-	0 (0)	0.05 (± 0.24)	0–2	7 (2.5)
Other	0.04 (± 0.20)	0–1	3 (1.6)	0.03 (± 0.16)	0–1	2 (2.1)	0.03 (± 0.18)	0–1	5 (1.7)

N = number; SD = standard deviation.

Supplement Eb: Fractions of subtypes in the category “changed medicines”.

