

		shoot length day 2		shoot length day 3		shoot length day 4		shoot length day 5		shoot length day 6		shoot length day 8	
	<i>d.f.</i>	F	<i>p</i>	F	<i>p</i>	F	<i>p</i>	F	<i>p</i>	F	<i>p</i>	F	<i>p</i>
Amoebae (AM)	1	0.74	0.39	3.84	0.052	6.83	0.01	9.2	0.003	9.31	0.003	7.86	0.005
Pythium (PY)	1	1.13	0.288	0.12	0.725	0.02	0.879	0.02	0.877	0.41	0.525	0.03	0.863
Bacterial Isolate													
(BI)	8	0.31	0.961	0.55	0.819	0.65	0.736	0.72	0.671	0.74	0.66	0.252	0.98
AM x PY	1	0.16	0.694	2.32	0.129	4.99	0.027	7.66	0.006	11.11	0.001	10.353	0.002
AM x BI	8	0.34	0.949	0.88	0.536	0.96	0.473	1.41	0.195	1.29	0.25	0.952	0.475
BI x PY	8	0.39	0.926	0.9	0.515	<i>1.91</i>	<i>0.061</i>	2.385	0.018	2.7	0.008	<i>1.985</i>	<i>0.051</i>
AM x PY x BI	8	1.22	0.288	<i>1.71</i>	<i>0.1</i>	2.02	0.047	2.75	0.007	2.99	0.004	2.464	0.0148

Significant effects are highlighted in bold, tendencies are highlighted in italic

Protozoa stimulate the plant beneficial activity of rhizospheric pseudomonads, Plant & Soil,

Simone Weidner*, Ellen Latz, Betina Agaras, Claudio Valverde, Alexandre Jousset,

*corr. author: Ecology and Biodiversity, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

S.Weidner@uu.nl