

Th17 cytokines and factors modulating their activity in patients with pernicious anemia




Fig. 6S The concentration of IL-33 in the blood of PA patients and healthy controls (C)
The difference between the groups is statistically significant. The median and quartiles 25% and 75% are presented (Mann-Whitney *U* test).

¹Dariusz Kajdaniuk*, ²Wanda Foltyn, ³Elżbieta Morawiec-Szymonik, ⁴Zenon Czuba, ⁵Ewa Szymonik, ²Beata Kos-Kudła, ¹Bogdan Marek

¹Department of Pathophysiology, Chair of Pathophysiology and Endocrinology, Medical University of Silesia, Katowice, Poland

²Department of Endocrinology and Neuroendocrine Tumors, Chair of Pathophysiology and Endocrinology, Medical University of Silesia, Katowice, Poland

³Department of Internal Medicine and Oncological Chemotherapy, Andrzej Mielęcki Independent Public Clinical Hospital, Katowice, Poland

⁴Department of Microbiology and Immunology, Medical University of Silesia, Katowice, Poland

⁵Department of Anesthesiology and Intensive Care, Stanisław Szyszko Independent Public Clinical Hospital No. 1, Zabrze, Poland

*corresponding author:

dkajdaniuk@sum.edu.pl