

Classification of Raw Açaí (*Euterpe oleracea* Mart.) Fruits from Upland and Lowland

Environments Using Portable Near-Infrared Spectroscopy

Hervé Rogez ^a; João Paulo Silva e Silva ^a; Risaldo Amaral Junior ^a; Lorena Modesto Caldeira ^a; Vincent

Baeten ^b; Neirivaldo Cavalcante da Silva ^{*a}

^a Center of Valorization of Amazonian Bioactive Compounds (CVACBA), Federal University of Pará, Av. Perimetral da Ciência, km 01, CEP: 66095-780, Belém, PA, Brazil.

^b Walloon Agricultural Research Centre (CRA-W), Valorisation of Agricultural Products Department, Food and Feed Quality Unit (U15), ‘Henseval Building’, Chaussée de Namur 24, 5030 Gembloux, Belgium.

*Corresponding author: ncs@ufpa.br, tel. +55 91 9 84519782

Supplementary material

Food and Bioprocess Technology

Additional figures showing the loading vectors of all latent variable–based classification models (PLS-DA, SIMCA, DD-SIMCA, and OCPLS) developed using the full spectral range and variable-selected datasets (SPA, iPLS, and VIP).

Fig.S1




Fig.S2




Fig.S3




Fig.S4

