

Preferred term codes included in searches for different safety topics.

CMQ 'overdose'

'multiple drug overdose', 'multiple drug overdose accidental', 'multiple drug overdose intentional', 'accidental overdose', 'intentional overdose', 'overdose', 'drug level above therapeutic', 'analgesic drug level above therapeutic', 'chemotherapeutic drug level above therapeutic', 'antasthmatic drug level above therapeutic', 'cardioactive drug level above therapeutic', 'antipsychotic drug level above therapeutic', 'anticonvulsant drug level above therapeutic', 'antidepressant drug level above therapeutic', 'anticoagulation drug level above therapeutic', 'muscle relaxant drug level above therapeutic', 'antiviral drug level above therapeutic', and 'antibiotic level above therapeutic'.

SMQ 'acute central respiratory depression'

'acute respiratory distress syndrome', 'acute respiratory failure', 'apnoea', 'apnoeic attack', 'bradypnoea', 'breath holding', 'breath sounds abnormal', 'breath sounds absent', 'central-alveolar hypoventilation', 'hypopnoea', 'hypoventilation', 'hypoventilation neonatal', 'infantile apnoea', 'lung hypoinflation', 'neonatal respiratory arrest', 'neonatal respiratory depression', 'neonatal respiratory failure', 'postoperative respiratory failure', 'respiratory arrest', 'respiratory depression', 'respiratory depth decreased', 'respiratory failure', 'respiratory paralysis', 'respiratory rate decreased', and 'severe acute respiratory syndrome'.

SMQ 'convulsion'

'acquired epileptic aphasia', 'acute encephalitis with refractory, repetitive partial seizures', 'alcoholic seizure', 'atonic seizures', 'atypical benign partial epilepsy', 'automatism epileptic', 'autonomic seizure', 'Baltic myoclonic epilepsy', 'benign familial neonatal convulsions', 'benign rolandic epilepsy', 'biotinidase deficiency', 'change in seizure presentation', 'clonic convulsion', 'complex partial seizures', 'convulsion in childhood', 'convulsion neonatal', 'convulsions local', 'convulsive threshold lowered', 'déjà vu', 'double cortex syndrome', 'dreamy state', 'drug withdrawal convulsions', 'early infantile epileptic encephalopathy with burst-suppression', 'eclampsia', 'epilepsy', 'epileptic aura', 'epileptic psychosis', 'febrile convulsion', 'frontal lobe epilepsy', 'generalised non-convulsive epilepsy', 'generalised tonic-clonic seizure', 'hyperglycaemic seizure', 'hypocalcaemic seizure', 'hypoglycaemic seizure', 'hyponatraemic seizure', 'idiopathic generalised epilepsy', 'infantile spasms', 'juvenile myoclonic epilepsy', 'Lafora's myoclonic epilepsy', 'Lennox-Gastaut syndrome', 'migraine-triggered seizure', 'molybdenum cofactor deficiency', 'myoclonic epilepsy', 'myoclonic epilepsy and ragged-red fibres', 'partial seizures', 'partial seizures with secondary generalisation', 'petit mal epilepsy', 'polymicrogyria', 'post stroke epilepsy', 'post stroke seizure', 'post-traumatic epilepsy', 'postictal headache', 'postictal paralysis', 'postictal psychosis', 'postictal state', 'psychomotor seizures', 'schizencephaly', 'seizure', 'seizure anoxic', 'seizure cluster', 'seizure like phenomena', 'severe myoclonic epilepsy of infancy', 'simple partial seizures', 'status epilepticus', 'sudden unexplained death in epilepsy', 'temporal lobe epilepsy', 'temporal lobe epilepsy', 'tonic clonic movements', 'tonic convulsion', 'tonic posturing', 'topectomy', and 'uncinate fits'.