

The peripheral neuropathy prevalence and characteristics are comparable in people with obesity and long duration type 1 diabetes

Lim JZM^{1,2}, Burgess J¹, Ooi CG², Ponirakis G³, Malik RA³, *Wilding JPH^{1,2}, *Alam, U^{1,2,4}

1 Department of Cardiovascular and Metabolic Medicine, Institute of Life Course and Medical Sciences, University of Liverpool, UK.

2 Division of Diabetes & Endocrinology, Aintree University Hospital, Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK.

3 Division of Medicine, Qatar Foundation, Weill Cornell Medicine-Qatar, Doha, Qatar

4 Department of Diabetes, Endocrinology & Gastroenterology, Faculty of Biology, Medicine and Health, University of Manchester, UK.

Corresponding author:

Dr Uazman Alam,

Department of Cardiovascular and Metabolic Medicine,

Institute of Life Course and Medical Sciences,

University of Liverpool,

Liverpool, UK.

E: uazman.alam@liverpool.ac.uk

Figure S1. Scatter Plot of VPT vs VAS for pain. Linear Regression ($R^2 = 0.579$) with line of best fit. VPT: vibration perception threshold; VAS: visual analogue score for pain

